

Spectral covers, algebraically completely integrable, Hamiltonian systems, and moduli of bundles

Ron Donagi ¹

University of Pennsylvania

and

Eyal Markman ²

University of Michigan

Contents

1	Introduction	3
2	Basic Notions	6
2.1	Symplectic Geometry	6
2.2	Integrable Systems	9
2.3	Algebraically Completely Integrable Hamiltonian Systems	11
2.4	Moment Maps and Symplectic Reduction	12
2.5	Finite dimensional Poisson loop group actions	17
2.5.1	Finite dimensional approximations	17
2.5.2	Type loci	19
3	Geodesic flow on an ellipsoid	22
3.1	Integrability	22
3.2	Algebraic integrability	24
3.3	The flows	26
3.4	Explicit parametrization	28
4	Spectral curves and vector bundles	31
4.1	Vector Bundles on a Curve	31
4.2	Spectral Curves and the Hitchin System	35
4.3	Polynomial Matrices	38
4.3.1	Explicit Equations for Jacobians of Spectral Curves with a Cyclic Ramification Point	40
4.3.2	Geodesic Flow on Ellipsoids via 2×2 Polynomial Matrices	41

¹Partially supported by NSA Grant MDA904-92-H3047 and NSF Grant DMS 95-03249

²Partially supported by a Rackham Fellowship, University of Michigan, 1993

5	Poisson structure via levels	44
5.1	Level structures	44
5.2	The cotangent bundle	44
5.3	The Poisson structure	45
5.4	Linearization	46
5.5	Hamiltonians and flows in $T^*\mathcal{U}_D$	48
6	Spectral flows and KP	49
6.1	The hierarchies	49
6.2	Krichever maps	51
6.3	Compatibility of hierarchies	54
6.3.1	Galois covers and relative Krichever maps	55
6.3.2	Compatibility of stratifications	57
6.3.3	The compatibility theorem, the smooth case	58
6.3.4	The compatibility theorem, singular cases	59
6.3.5	Elliptic Solitons	62
6.3.6	Outline of the proof of the compatibility theorem	65
7	The Cubic Condition and Calabi-Yau threefolds	68
7.1	Families of Tori	68
7.2	The Cubic Condition	70
7.3	An Integrable System for Calabi-Yau Threefolds	73
8	The Lagrangian Hilbert scheme and its relative Picard	84
8.1	Introduction	84
8.2	Lagrangian Hilbert Schemes	86
8.3	The construction of the symplectic structure	89
8.4	Partial compactifications: a symplectic structure on the moduli space of Lagrangian sheaves	96
8.5	Examples	98
8.5.1	Higgs Pairs	98
8.5.2	Fano Varieties of Lines on Cubic Fourfolds	100
9	Spectral covers	101
9.1	Algebraic extensions	101
9.2	Decomposition of spectral Picards	103
9.2.1	The question	103
9.2.2	Decomposition of spectral covers	105
9.2.3	Decomposition of spectral Picards	106
9.2.4	The distinguished Prym	107
9.3	Abelianization	108
9.3.1	Abstract vs. K -valued objects	108
9.3.2	The regular semisimple case: the shift	109
9.3.3	The regular case: the twist along the ramification	111
9.3.4	Adding values and representations	113
9.3.5	Irregulars?	114

1 Introduction

The purpose of these notes is to present an algebro-geometric point of view on several interrelated topics, all involving integrable systems in symplectic-algebro-geometric settings. These systems range from some very old examples, such as the geodesic flow on an ellipsoid, through the classical hierarchies of KP - and KdV -types, to some new systems which are often based on moduli problems in algebraic geometry.

The interplay between algebraic geometry and integrable systems goes back quite a way. It has been known at least since Jacobi that many integrable systems can be solved explicitly in terms of *theta functions*. (There are numerous examples, starting with various *spinning tops* and the *geodesic flow on an ellipsoid*.) Geometrically, this often means that the system can be mapped to the total space of a family of Jacobians of some curves, in such a way that the flows of the system are mapped to linear flows along the Jacobians. In practice, these curves tend to arise as the spectrum (hence the name ‘*spectral*’ curves) of some parameter-dependent operator; they can therefore be represented as branched covers of the parameter space, which in early examples tended to be the Riemann sphere \mathbf{CP}^1 .

In *Hitchin’s system*, the base \mathbf{CP}^1 is replaced by an arbitrary (compact, non-singular) Riemann surface Σ . The cotangent bundle $T^*\mathcal{U}_\Sigma$ to the moduli space \mathcal{U}_Σ of stable vector bundles on Σ admits two very different interpretations: on the one hand, it parametrizes certain *Higgs bundles*, or vector bundles with a (canonically) twisted endomorphism; on the other, it parametrizes certain *spectral data*, consisting of torsion-free sheaves (generically, line bundles) on spectral curves which are branched covers of Σ . In our three central chapters (4,5,6) we study this important system, its extensions and variants. All these systems are linearized on Jacobians of spectral curves.

We also study some systems in which the spectral curve is replaced by a higher-dimensional geometric object: a *spectral variety* in Chapter 9, an algebraic *Lagrangian subvariety* in Chapter 8, and a *Calabi-Yau manifold* in Chapter 7. Our understanding of some of these wild systems is much less complete than in the case of the curve-based ones. We try to explain what we know and to point out some of what we do not. The Calabi-Yau systems seem particularly intriguing. Not only are the tori (on which these systems are linearized) not Jacobians of curves, they are in general not even abelian varieties. There are some suggestive relations between these systems and the conjectural mirror-symmetry for Calabi-Yaus.

The first three chapters are introductory. In Chapter 2 we collect the basic notions of *symplectic geometry* and *integrable systems* which will be needed, including some information about *symplectic reduction*. (An excellent further reference is [AG].) In Chapter 3 we work out in some detail the classical theory of geodesic flow on an ellipsoid, which is integrable via hyperelliptic theta functions. We think of this both as a beautiful elementary and explicit example and as an important special case of the much more powerful results which follow. (Our presentation follows [Kn, Re, D5]). Some of our main algebro-geometric objects of study are introduced in Chapter 4: vector bundles and their moduli spaces, spectral curves, and the ‘*spectral systems*’ constructed from them. In particular, we consider the *polynomial matrix system* [AHH, B1] (which contains the geodesic flow on an ellipsoid as special case) and *Hitchin’s system* [H1, H2].

Each of the remaining five chapters presents in some detail a recent or current re-

search topic. Chapter 5 outlines constructions (from [Ma1, Bn, Ty1]) of the Poisson structure on the spectral system of curves. This is possible whenever the twisting line bundle K is a non-negative twist $\omega_\Sigma(D)$ of the canonical bundle ω_Σ , and produces an algebraically completely integrable Hamiltonian system. Following [Ma1] we emphasize the deformation-theoretic construction, in which the Poisson structure on an open subset of the system is obtained via symplectic reduction from the cotangent bundle $T^*\mathcal{U}_{\Sigma,D}$ of the moduli space $\mathcal{U}_{\Sigma,D}$ of stable bundles with a *level- D structure*.

In Chapter 6 we explore the relation between these spectral systems and the KP -hierarchy and its variants (multi-component KP , Heisenberg flows, and their KdV -type subhierarchies). These hierarchies are, of course, a rich source of geometry: The Krichever construction (e.g. [SW]) shows that any Jacobian can be embedded in KP -space, and these are the only finite-dimensional orbits [Mul, AdC, Sh]. Following [AB, LM1] we describe some “multi-Krichever” constructions which take spectral data to the spaces of the KP , $mcKP$ and Heisenberg systems. Our main new result is that the flows on the spectral system which are obtained by pulling back the $mcKP$ or Heisenberg flows via the corresponding Krichever maps are *Hamiltonian* with respect to the Poisson structure constructed in Chapter 5. In fact, we write down explicitly the Hamiltonians for these KP flows on the spectral system, as residues of traces of meromorphic matrices. (Some related results have also been obtained recently in [LM2].)

The starting point for Chapter 7 is an attempt to understand the condition for a given family of complex tori to admit a symplectic structure and thus become an ACIHS. We find that the condition is a symmetry on the derivatives of the period map, which essentially says that the periods are obtained as partials of some field of symmetric cubic tensors on the base. In the rest of this Chapter we apply this idea to an analytically (not algebraically) integrable system constructed from any family of Calabi-Yau 3-folds. Some properties of this system suggest that it may be relevant to a purely hodge-theoretic reformulation of the mirror-symmetry conjectures. (This chapter is based on [DM].)

Chapter 8 is devoted to the construction of symplectic and Poisson structures in some inherently non-linear situations, vastly extending the results of Chapter 5. The basic space considered here is the moduli space parametrizing line-bundle-like sheaves supported on (variable) subvarieties of a given symplectic space X . It is shown that when the subvarieties are Lagrangian, the moduli space itself becomes symplectic. The spectral systems considered in Chapter 5 can be recovered as the case where X is the total space of $T^*\Sigma$ and the Lagrangian subvarieties are the spectral curves. (A fuller version of these results will appear in [Ma2].)

In the final chapter we consider extensions of the spectral system to allow a higher-dimensional base variety S , an arbitrary reductive group G , an arbitrary representation $\rho : G \rightarrow \text{Aut}V$, and values in an arbitrary vector bundle K . (Arbitrary reductive groups G were considered, over a curve $S = \Sigma$ with $K = \omega_\Sigma$, by Hitchin [H2], while the case $K = \Omega_S$ over arbitrary base S is Simpson’s [Sim1]). We replace spectral curves by various kinds of spectral covers, and introduce the cameral cover, a version of the Galois-closure of a spectral cover which is independent of K and ρ . It comes with an action of W , the Weyl group of G . We analyze the decomposition, under the action of W , of the cameral and spectral Picard varieties, and identify the distinguished Prym in there. This is shown to correspond, up to certain shifts and twists, to the fiber of the Hitchin map in this general setting, i.e. to moduli of Higgs bundles with a given \tilde{S} . Combining this

with some obvious remarks about existence of Poisson structures, we find that the moduli spaces of K -valued Higgs bundles support algebraically completely integrable systems. Our presentation closely follows that of [D4]

It is a pleasure to express our gratitude to the organizers, Mauro Francaviglia and Silvio Greco, for the opportunity to participate in the CIME meeting and to publish these notes here. During the preparation of this long work we benefited from many enjoyable conversations with M. Adams, M. Adler, A. Beauville, R. Bryant, C. L. Chai, I. Dolgachev, L. Ein, B. van Geemen, A. Givental, M. Green, P. Griffiths, N. Hitchin, Y. Hu, S. Katz, V. Kanev, L. Katzarkov, R. Lazarsfeld, P. van Moerbeke, D. Morrison, T. Pantev, E. Previato and E. Witten.

2 Basic Notions

We gather here those basic concepts and elementary results from symplectic and Poisson geometry, completely integrable systems, and symplectic reduction which will be helpful throughout these notes. Included are a few useful examples and only occasional proofs or sketches. To the reader unfamiliar with this material we were hoping to impart just as much of a feeling for it as might be needed in the following chapters. For more details, we recommend the excellent survey [AG].

2.1 Symplectic Geometry

Symplectic structure

A symplectic structure on a differentiable manifold M of even dimension $2n$ is given by a non-degenerate closed 2-form σ . The non degeneracy means that either of the following equivalent conditions holds.

- σ^n is a nowhere vanishing volume form.
- Contraction with σ induces an isomorphism $\lrcorner\sigma : TM \rightarrow T^*M$
- For any non-zero tangent vector $v \in T_mM$ at $m \in M$, there is some $v' \in T_mM$ such that $\sigma(v, v') \neq 0$.

Examples 2.1

1. Euclidean space

The standard example of a symplectic manifold is Euclidean space \mathbb{R}^{2n} with $\sigma = \sum dp_i \wedge dq_i$, where $p_1, \dots, p_n, q_1, \dots, q_n$ are linear coordinates. Darboux's theorem says that any symplectic manifold is locally equivalent to this example (or to any other).

2. Cotangent bundles

For any manifold X , the cotangent bundle $M := T^*X$ has a natural symplectic structure. First, M has the tautological 1-form α , whose value at $(x, \theta) \in T^*X$ is θ pulled back to T^*M . If q_1, \dots, q_n are local coordinates on X , then locally $\alpha = \sum p_i dq_i$ where the p_i are the fiber coordinates given by $\partial/\partial q_i$. The differential

$$\sigma := d\alpha$$

is then a globally defined closed (even exact) 2-form on M . It is given in local coordinates by $\sum dp_i \wedge dq_i$, hence is non-degenerate.

3. Coadjoint orbits

Any Lie group G acts on its Lie algebra \mathfrak{g} (adjoint representation) and hence on the dual vector space \mathfrak{g}^* (coadjoint representation). Kostant and Kirillov noted that for any $\xi \in \mathfrak{g}^*$, the coadjoint orbit $\mathcal{O} = G\xi \subset \mathfrak{g}^*$ has a natural symplectic structure. The tangent space to \mathcal{O} at ξ is given by $\mathfrak{g}/\mathfrak{g}_\xi$, where \mathfrak{g}_ξ is the stabilizer of ξ :

$$\mathfrak{g}_\xi := \{x \in \mathfrak{g} \mid ad_x^* \xi = 0\} = \{x \in \mathfrak{g} \mid (\xi, [x, y]) = 0 \quad \forall y \in \mathfrak{g}\}.$$

Now ξ determines an alternating bilinear form on \mathfrak{g}

$$x, y \longmapsto (\xi, [x, y]),$$

which clearly descends to $\mathfrak{g}/\mathfrak{g}_\xi$ and is non-degenerate there. Varying ξ we get a non-degenerate 2-form σ on \mathcal{O} . The Jacobi identity on g translates immediately into closedness of σ .

Hamiltonians

To a function f on a symplectic manifold (M, σ) we associate its *Hamiltonian vector field* v_f , uniquely determined by

$$v_f \lrcorner \sigma = df.$$

A vector field v on M is Hamiltonian if and only if the 1-form $v \lrcorner \sigma$ is exact. We say v is *locally Hamiltonian* if $v \lrcorner \sigma$ is closed. This is equivalent to saying that the flow generated by v preserves σ . Thus on a symplectic surface ($n = 1$), the locally Hamiltonian vector fields are the area-preserving ones.

Example: (Geodesic flow)

A Riemannian metric on a manifold X determines an isomorphism of $M := TX$ with T^*X ; hence we get on M a natural symplectic structure together with a C^∞ function $f =$ (squared length). The geodesic flow on X is the differential equation, on M , given by the Hamiltonian vector field v_f . Its integral curves are the geodesics on M .

Poisson structures

The association $f \mapsto v_f$ gives a map of sheaves

$$v : C^\infty(M) \longrightarrow V(M) \tag{1}$$

from C^∞ functions on the symplectic manifold M to vector fields. Now $V(M)$ always has the structure of a Lie algebra, under commutation of vector fields. The symplectic structure on M determines a Lie algebra structure on $C^\infty(M)$ such that v becomes a morphism of (sheaves of) Lie algebras. The operation on $C^\infty(M)$, called *Poisson bracket*, is

$$\{f, g\} := (df, v_g) = -(dg, v_f) = \frac{ndf \wedge dg \wedge \sigma^{n-1}}{\sigma^n}.$$

More generally, a *Poisson structure* on a manifold M is a Lie algebra bracket $\{, \}$ on $C^\infty(M)$ which acts as a derivation in each variable:

$$\{f, gh\} = \{f, g\}h + \{f, h\}g, \quad f, g, h \in C^\infty(M).$$

Since the value at a point m of a given derivation acting on a function g is a linear function of $d_m g$, we see that a Poisson structure on M determines a global 2-vector

$$\psi \in H^0(M, \overset{2}{\wedge} TM).$$

or equivalently a skew-symmetric homomorphism

$$\Psi : T^*M \longrightarrow TM.$$

Conversely, any 2-vector ψ on M determines an alternating bilinear bracket on $C^\infty(M)$, by

$$\{f, g\} := (df \wedge dg, \psi),$$

and this acts as a derivation in each variable. An equivalent way of specifying a Poisson structure is thus to give a global 2-vector ψ satisfying an integrability condition (saying that the above bracket satisfies the Jacobi identity, hence gives a Lie algebra).

We saw that a symplectic structure σ determines a Poisson bracket $\{, \}$. The corresponding homomorphism Ψ is just $(\lrcorner\sigma)^{-1}$; the closedness of σ is equivalent to integrability of ψ . Thus, a Poisson structure which is (i.e. whose 2-vector is) everywhere non-degenerate, comes from a symplectic structure.

A general Poisson structure can be degenerate in two ways: first, there may exist non-constant functions $f \in C^\infty(M)$, called *Casimirs*, satisfying

$$0 = df \lrcorner \psi = \Psi(df),$$

i.e.

$$\{f, g\} = 0 \text{ for all } g \in C^\infty(M).$$

This implies that the rank of Ψ is less than maximal everywhere. In addition, or instead, rank Ψ could drop along some strata in M . For even r , let

$$M_r := \{m \in M \mid \text{rank}(\Psi) = r\}.$$

Then a basic result [We] asserts that the M_r are submanifolds, and they are canonically foliated into *symplectic leaves*, i.e. r -dimensional submanifolds $Z \subset M_r$ which inherit a symplectic structure. (This means that the restriction $\psi|_Z$ is the image, under the inclusion $Z \hookrightarrow M_r$, of a two-vector ψ_Z on Z which is everywhere nondegenerate, hence comes from a symplectic structure on Z .) These leaves can be described in several ways:

- The image $\Psi(T^*M_r)$ is an involutive subbundle of rank r in TM_r ; the Z are its integral leaves.
- The leaf Z through $m \in M_r$ is $Z = \{z \in M_r \mid f(m) = f(z) \text{ for all Casimirs } f \text{ on } M_r\}$.
- Say that two points of M are ψ -connected if there is an integral curve of some Hamiltonian vector field passing through both. The leaves are the equivalence classes for the equivalence relation generated by ψ -connectedness.

Example 2.2 The Kostant-Kirillov symplectic structures on coadjoint orbits of a Lie algebra \mathfrak{g} extend to a Poisson structure on the dual vector space \mathfrak{g}^* . For a function $F \in C^\infty(\mathfrak{g}^*)$ we identify its differential $d_\xi F$ at $\xi \in \mathfrak{g}^*$ with an element of $\mathfrak{g} = \mathfrak{g}^{**}$. We then set:

$$\{F, G\}(\xi) := (\xi, [d_\xi F, d_\xi G]).$$

This is a Poisson structure, whose symplectic leaves are precisely the coadjoint orbits. The rank of \mathfrak{g} is, by definition, the smallest codimension ℓ of a coadjoint orbit. The Casimirs are the ad-invariant functions on \mathfrak{g}^* . Their restrictions to the largest stratum $\mathfrak{g}_{\dim \mathfrak{g} - \ell}^*$ foliate this stratum, the leaves being the *regular* (i.e. largest dimensional) coadjoint orbits.

2.2 Integrable Systems

We say that two functions h_1, h_2 on a Poisson manifold (M, ψ) *Poisson commute* if their Poisson bracket $\{h_1, h_2\}$ is zero. In this case the integral flow of the Hamiltonian vector field of each function h_i , $i = 1, 2$ is tangent to the level sets of the other. In other words, h_2 is a conservation law for the Hamiltonian h_1 and the Hamiltonian flow of h_2 is a symmetry of the Hamiltonian system associated with (M, ψ, h_1) (the flow of the Hamiltonian vector field v_{h_1} on M).

A map $f : M \rightarrow B$ between two Poisson manifolds is a *Poisson map* if pullback of functions is a Lie algebra homomorphism with respect to the Poisson bracket

$$f^*\{F, G\}_B = \{f^*F, f^*G\}_M.$$

Equivalently, if $df(\psi_M)$ equals $f^*(\psi_B)$ as sections of $f^*(\wedge^2 T_B)$. If $H : M \rightarrow B$ is a Poisson map with respect to the trivial (zero) Poisson structure on B we will call H a *Hamiltonian map*. Equivalently, H is Hamiltonian if the Poisson structure ψ vanishes on the pullback $H^*(T^*B)$ of the cotangent bundle of B (regarding the latter as a subbundle of (T^*M, ψ)). In particular, the rank of the differential dH is less than or equal to $\dim M - \frac{1}{2} \text{rank}(\psi)$ at every point. A Hamiltonian map pulls back the algebra of functions on B to a commutative Poisson subalgebra of the algebra of functions on M .

The study of a Hamiltonian system (M, ψ, h) simplifies tremendously if one can extend the Hamiltonian function h to a Hamiltonian map $H : M \rightarrow B$ of maximal rank $\dim M - \frac{1}{2} \text{rank}(\psi)$. Such a system is called a completely integrable Hamiltonian system. The Hamiltonian flow of a completely integrable system can often be realized as a linear flow on tori embedded in M . The fundamental theorem in this case is Liouville's theorem (stated below).

Definition 2.3 1. Let V be a vector space, $\sigma \in \wedge^2 V^*$ a (possibly degenerate) two form.

A subspace $Z \subset V$ is called *isotropic (coisotropic)* if it is contained in (contains) its symplectic complement. Equivalently, Z is isotropic if σ restricts to zero on Z . If σ is nondegenerate, a subspace $Z \subset V$ is called *Lagrangian* if it is both isotropic and coisotropic. In this case V is even (say $2n$) dimensional and the Lagrangian subspaces are the n dimensional isotropic subspaces.

2. Let (M, σ) be a symplectic manifold. A submanifold Z is *isotropic* (respectively *coisotropic*, *Lagrangian*) if the tangent subspaces $T_z Z$ are, for all $z \in Z$.

Example 2.4 For every manifold X , the fibers of the cotangent bundle T^*X over points of X are Lagrangian submanifolds with respect to the standard symplectic structure. A section of T^*X over X is Lagrangian if and only if the corresponding 1-form on X is closed.

We will extend the above definition to Poisson geometry:

- Definition 2.5** 1. Let U be a vector space, ψ an element of $\overset{2}{\wedge} U$. Let $V \subset U$ be the image of the contraction $\lrcorner \psi : U^* \rightarrow U$. Let $W \subset U^*$ be its kernel. W is called the null space of ψ . ψ is in fact a nondegenerate element of $\overset{2}{\wedge} V$ giving rise to a symplectic form $\sigma \in \overset{2}{\wedge} V^*$ (its inverse). A subspace $Z \subset U$ is *Lagrangian* with respect to ψ if Z is a Lagrangian subspace of $V \subset U$ with respect to σ . Equivalently, Z is Lagrangian if $(U/Z)^*$ is both an isotropic and a coisotropic subspace of U^* with respect to $\psi \in \overset{2}{\wedge} U \cong \overset{2}{\wedge} (U^*)^*$.
2. Let (M, ψ) be a Poisson manifold, assume that ψ has constant rank (this condition will be relaxed in the complex analytic or algebraic case). A submanifold $Z \subset M$ is *Lagrangian* if the tangent subspaces $T_z Z$ are, for all $z \in Z$. Notice that the constant rank assumption implies that each connected component of Z is contained in a single symplectic leaf.

Theorem 2.6 (*Liouville*). *Let M be an m -dimensional Poisson manifold with Poisson structure ψ of constant rank $2g$. Suppose that $H : M \rightarrow B$ is a proper submersive Hamiltonian map of maximal rank, i.e., $\dim B = m - g$. Then*

- i) *The null foliation of M is induced locally by a foliation of B (globally if H has connected fibers).*
- ii) *The connected components of fibers of H are Lagrangian compact tori with a natural affine structure.*
- iii) *The Hamiltonian vector fields of the pullback of functions on B by H are tangent to the level tori and are translation invariant (linear).*

Remark 2.7 : If H is not proper, but the Hamiltonian flows are complete, then the fibers of H are generalized tori (quotients of a vector space by a discrete subgroup, not necessarily of maximal rank).

Sketch of proof of Liouville's theorem:

- i) Since H is a proper submersion the connected components of the fibers of H are smooth compact submanifolds. Since H is a Hamiltonian map of maximal rank $m - g$, the pullback $H^*(T_B^*)$ is isotropic and coisotropic and hence H is a Lagrangian fibration. In particular, each connected component of a fiber of H is contained in a single symplectic leaf.
- ii),iii) Let A_b be a connected component of the fiber $H^{-1}(b)$. Let $0 \rightarrow T_{A_b} \rightarrow T_{M|_{A_b}} \xrightarrow{dH} (T_b B) \otimes \mathcal{O}_{A_b} \rightarrow 0$ be the exact sequence of the differential of H . Part i) implies that the null subbundle $W_{|_{A_b}} := \text{Ker}[\Psi : T^*M \rightarrow TM]_{|_{A_b}}$ is the pullback of a subspace W_b of $T_b^* B$. Since H is a Lagrangian fibration, the Poisson structure

induces a surjective homomorphism $\phi_b : H^*(T_b^*B) \rightarrow T_{A_b}$ inducing a trivialization $\bar{\phi}_b : (T_b^*B/W_b) \otimes \mathcal{O}_{A_b} \xrightarrow{\sim} T_{A_b}$.

A basis of the vector space T_b^*B/W_b corresponds to a frame of global independent vector fields on the fiber A_b which commute since the map H is Hamiltonian. Hence A_b is a compact torus.

□

2.3 Algebraically Completely Integrable Hamiltonian Systems

All the definitions and most of the results stated in this chapter for C^∞ -manifolds translate verbatim and hold in the complex analytic and complex algebro-geometric categories replacing the real symplectic form by a holomorphic or algebraic $(2, 0)$ -form (similarly for Poisson structures). The (main) exception listed below is due to the differences between the Zariski topology and the complex or C^∞ topologies. A Zariski open subset is the complement of the zero locus of a system of polynomial equations. It is hence always a dense open subset.

The (local) foliation by symplectic leaves exists only local analytically. For example, a rank 2 translation invariant section $\psi \in H^0(A, \wedge^2 TA)$ on a 3 dimensional abelian variety A which is simple (does not contain any abelian subvariety) is an algebraic Poisson structure with a non algebraic null foliation.

We will relax the definitions of a Lagrangian subvariety and integrable system in the algebro-geometric category:

Definition 2.8 Let (M, ψ) be a Poisson smooth algebraic variety. An irreducible and reduced subvariety $Z \subset M$ is *Lagrangian* if the tangent subspace $T_z Z \subset T_z M$ is Lagrangian for a generic point $z \in Z$.

Definition 2.9 An *algebraically completely integrable Hamiltonian system* consists of a proper flat morphism $H : M \rightarrow B$ where (M, ψ) is a smooth Poisson variety and B is a smooth variety such that, over the complement $B \setminus \Delta$ of some proper closed subvariety $\Delta \subset B$, H is a Lagrangian fibration whose fibers are isomorphic to abelian varieties.

Multiples of a theta line bundle embed an abelian variety in projective spaces with the coordinates being theta functions. Thus, a priori, the solutions of an algebraically completely integrable Hamiltonian system can be expressed in terms of theta functions. Finding explicit formulas is usually hard. In the next chapter we will study one example, the geodesic flow on ellipsoids, in some detail. Later we will encounter certain equations of Kdv type, the Hitchin system, and a few other examples. Other classical integrable systems include various Euler-Arnold systems, spinning tops, the Neumann system of evolution of a point on the sphere subject to a quadratic potential.

Most of these systems are the complexification of real algebraic systems. Given a real algebraic symplectic variety (M, σ) and an algebraic Hamiltonian h on M we say that the system is *algebraically completely integrable* if its complexification $(M_{\mathbb{C}}, \sigma_{\mathbb{C}}, h_{\mathbb{C}})$ is. A real completely integrable system (M, σ, h) need not be algebraically completely integrable even if (M, σ, h) are algebraic:

A Counter Example: Let (M, σ) be $(\mathbb{R}^2, dx \wedge dy)$ and $h : \mathbb{R}^2 \rightarrow \mathbb{R}$ a polynomial of degree d whose level sets are nonsingular. The system is trivially completely integrable, but it is algebraically completely integrable if and only if $d = 3$ because in all other cases the generic fiber of the complexification is a complex affine plane curve of genus $\frac{(d-1)(d-2)}{2} \neq 1$.

Action Angle Coordinates:

Let (M, σ) be a $2n$ -dimensional symplectic manifold, $H : M \rightarrow B$ a Lagrangian fibration by compact connected tori.

Theorem 2.10 (*real action angle coordinates*).

In a neighborhood of a fiber of $H : M \rightarrow B$ one can introduce the structure of a direct product $(\mathbb{R}^n/\mathbb{Z}^n) \times \mathbb{R}^n$ with action coordinates $(I_1 \cdots I_n)$ on the factor \mathbb{R}^n and angular coordinates $(\phi_1, \cdots \phi_n)$ on the torus $(\mathbb{R}^n/\mathbb{Z}^n)$ in which the symplectic structure has the form $\sum_{k=1}^n dI_k \wedge d\phi_k$.

The Local action coordinates on B are canonical up to affine transformation on \mathbb{R}^n with differential in $SL(n, \mathbb{Z})$. The angle coordinates depend canonically on the action coordinates and a choice of a Lagrangian section of $H : M \rightarrow B$.

Remarks:

1. In action angle coordinates the equations of the Hamiltonian flow of a function h on B becomes: $\dot{I}_k = 0, \dot{\phi}_k = c_k(I_1, \cdots, I_n)$ where the slopes c_k are $c_k = \frac{\partial h}{\partial I_k}$.
2. In the polarized complex analytic case, we still have local holomorphic action coordinates. They depend further on a choice of a Lagrangian subspace of the integral homology $H_1(A_b, \mathbb{Z})$ with respect to the polarization (a section of $\wedge^2 H^1(A_b, \mathbb{Z})$).

2.4 Moment Maps and Symplectic Reduction

Poisson Actions

An action ρ of a connected Lie group G on a manifold M determines an *infinitesimal action*

$$d\rho : \mathfrak{g} \longrightarrow V(M),$$

which is a homomorphism from the Lie algebra of G to the Lie algebra of C^∞ vector fields on M . When (M, σ) is symplectic, we say that the action ρ is *symplectic* if

$$(\rho(g))^* \sigma = \sigma, \quad \text{all } g \in G,$$

or equivalently if the image of $d\rho$ consists of locally Hamiltonian vector fields.

We say that the action ρ is *Poisson* if it factors through the Lie algebra homomorphism (1) $v : C^\infty(M) \rightarrow V(M)$ and a Lie algebra homomorphism

$$H : \mathfrak{g} \longrightarrow C^\infty(M).$$

This imposes two requirements on ρ , each of a cohomological nature: the locally Hamiltonian fields $d\rho(X)$ should be globally Hamiltonian, $d\rho(X) = v(H(X))$; and it must be possible to choose the $H(X)$ consistently so that

$$H([X, Y]) = \{H(X), H(Y)\}.$$

(a priori the difference between the two terms is a constant function, since its v is zero, so the condition is that it should be possible to make all these constants vanish simultaneously.)

Moment Maps

Instead of specifying the Hamiltonian lift

$$H : \mathfrak{g} \longrightarrow C^\infty(M)$$

for a Poisson action of G on (M, σ) , it is convenient to consider the equivalent data of the *moment map*

$$\mu : M \longrightarrow \mathfrak{g}^*$$

defined by

$$(\mu(m), X) := H(X)(m).$$

It is a Poisson map with respect to the Kostant-Kirillov Poisson structure on \mathfrak{g} (example 2.2), and is G -equivariant.

Examples 2.11 1. Any action of G on a manifold X lifts to an action on $M := T^*X$. This action is Poisson. The corresponding moment map $T^*X \rightarrow \mathfrak{g}^*$ is the dual of the infinitesimal action $\mathfrak{g} \rightarrow \Gamma(TX)$. It can be identified with the pullback of differential forms from X to G via the action.

2. The coadjoint action of G on \mathfrak{g}^* is Poisson, with the identity as moment map.

Symplectic Reduction

Consider a Poisson action of G on (M, σ) for which a reasonable quotient G/M exists. (We will remain vague about this for now, and discuss the properties of the quotient on a case-by-case basis. A general sufficient condition for the quotient to be a manifold is that the action is proper and free.) The Poisson bracket on M then descends to give a Poisson structure on M/G . The moment map,

$$\mu : M \longrightarrow \mathfrak{g}^*,$$

determines the symplectic leaves of this Poisson structure: let $\xi = \mu(m)$, let \mathcal{O} be the coadjoint orbit through ξ and let G_ξ be the stabilizer of ξ . Assume for simplicity that $\mu^{-1}(\xi)$ is connected and μ is submersive at $\mu^{-1}(\xi)$. Then, the leaf through m is

$$\mu^{-1}(\mathcal{O}_\xi)/G \approx \mu^{-1}(\xi)/G_\xi.$$

These symplectic leaves are often called the Marsden-Weinstein reductions M_{red} of M .

As an example, consider a situation where G acts on X with nice quotient X/G . The lifted action of G on $M = T^*X$ is Poisson, and has a quotient M/G which is a vector

bundle over X/G . The cotangent $T^*(X/G)$ sits inside $(T^*X)/G$ as the symplectic leaf over the trivial orbit $\mathcal{O}_0 = \{0\} \subset \mathfrak{g}^*$.

In contrast, the action of G on \mathfrak{g}^* does not in general admit a reasonable quotient. Its action on the dense open subset \mathfrak{g}_{reg}^* of regular elements (cf. example 2.2) does have a quotient, which is a manifold. The Poisson structure on the quotient is trivial, so the symplectic leaves are points, in one-to-one correspondence with the regular orbits. We refer to this quotient simply as \mathfrak{g}^*/G . The map $\pi_{reg} : \mathfrak{g}_{reg}^* \rightarrow \mathfrak{g}^*/G$ extends to $\pi : \mathfrak{g}^* \rightarrow \mathfrak{g}^*/G$, and there is a sense in which \mathfrak{g}^*/G really is the quotient of all \mathfrak{g}^* . Each coadjoint orbit \mathcal{O} is contained in the closure of a unique regular orbit \mathcal{O}' and $\pi(\mathcal{O}) = \pi_{reg}(\mathcal{O}')$.

A Diagram of Quotients

In the general situation of Poisson action (with a nice quotient π) of G on a symplectic manifold (M, σ) , there is another, larger, Poisson manifold \bar{M} , which can also be considered as a reduction of M by G . Everything fits together in the commutative diagram of Poisson maps:

$$\begin{array}{ccc}
 & M & \\
 & \downarrow & \\
 \pi & \bar{M} & \mu \\
 \swarrow & & \searrow \\
 M/G & & \mathfrak{g}^* \\
 \swarrow & & \searrow \\
 & \mathfrak{g}^*/G & \\
 & \downarrow & \\
 & (0) &
 \end{array} \tag{2}$$

\bar{M} may be described in several ways:

- \bar{M} is the quotient of M by the equivalence relation $m \sim gm$ if $g \in G_{\mu(m)}$, i.e., if $g(\mu(m)) = \mu(m)$.
- \bar{M} is the fiber product $\bar{M} = (M/G) \times_{(\mathfrak{g}^*/G)} \mathfrak{g}^*$.
- \bar{M} is the dual realization to the realization $M \rightarrow \mathfrak{g}^*/G$.

A *realization* of a Poisson manifold P is defined to be a Poisson map from a symplectic manifold M to P (see [We]). The realization will be called *full* if it is submersive. A pair of realizations $P_2 \xleftarrow{f_2} M \xrightarrow{f_1} P_1$ is called a dual pair if functions on one induce vector fields along the fibers of the other (i.e., the two opposite foliations are symplectic complements of each other).

We note that in the diagram of quotients, any two opposite spaces are a dual pair of realizations.

Given a full dual pair with connected fibers, the symplectic leaf foliations on P_1 and P_2 induce the same foliation on M (P_1 and P_2 have the “same” Casimir functions). The bijection between symplectic leaves on P_1 and P_2 is given by

$$P_1 \supset S_1 \mapsto f_2(f_1^{-1}(S_1)) = f_2(f_1^{-1}(x)) \quad \forall x \in S_1.$$

Returning to moment maps, we have over a coadjoint orbit $\mathcal{O} \subset \mathfrak{g}^*$:

- $\mu^{-1}(\mathcal{O})$ is coisotropic in M
- $\pi(\mu^{-1}(\mathcal{O}))$ is a symplectic leaf M_{red} in M/G
- $\bar{\mu}^{-1}(\mathcal{O})$ is also a symplectic leaf in \bar{M} . It is isomorphic to $\mu^{-1}(\mathcal{O})/(\text{null})$, or to $\mu^{-1}(\mathcal{O})/\sim$, or to $M_{red} \times \mathcal{O}$.

Example 2.12 Take M to be the cotangent bundle T^*G of a Lie group G . Denote by $\mu_L : T^*G \rightarrow \mathfrak{g}^*$ the moment map for the lifted left action of G . The quotient $\pi : M \rightarrow M/G$ is just the moment map $\mu_R : T^*G \rightarrow \mathfrak{g}^*$ for the lifted right action, and \bar{M} is the fiber product $\mathfrak{g}^* \times_{(\mathfrak{g}^*/G)} \mathfrak{g}^*$.

Example 2.13 If G is a connected commutative group T , the pair of nodes \mathfrak{t}^* and \mathfrak{t}^*/T coincide. Consequently, so do M/T and \bar{M} . The diagram of quotients degenerates to

$$\begin{array}{ccc} M & & \\ \downarrow \pi & & \\ M/T & & \\ \downarrow \bar{\mu} & & (3) \\ \mathfrak{t}^* & & \\ \downarrow & & \\ (0) & & \end{array}$$

Example 2.14 Consider two Poisson actions on (M, σ) of two groups G, T with moment maps μ_G, μ_T with connected fibers. Assume that

i) The actions of G and T commute.

It follows that $\mu_T : M \rightarrow \mathfrak{t}^*$ factors through M/G and $\mu_G : M \rightarrow \mathfrak{g}^*$ factors through M/T . Assume moreover

ii) T is commutative,

iii) $M \rightarrow \mathfrak{g}^*/G$ factors through \mathfrak{t}^*

Then $\bar{\mu}_G : \bar{M}_G \rightarrow \mathfrak{g}^*$ factors through M/T and the two quotient diagrams fit nicely together:

$$\begin{array}{ccccc}
 & & M & & \\
 & & \downarrow & & \\
 & & \bar{M}_G & & \\
 & \swarrow & & \searrow & \\
 M/G & & & & M/T \\
 & \searrow & & \swarrow & \\
 & & \mathfrak{t}^* & & \mathfrak{g}^* \\
 & & \swarrow & \searrow & \\
 & & \mathfrak{g}^*/G & & \\
 & & \downarrow & & \\
 & & (0) & &
 \end{array} \tag{4}$$

Remark 2.15 Note that condition iii in example 2.14 holds whenever $M/G \rightarrow \mathfrak{t}^*$ is a completely integrable system (with connected fibers). In that case the map $M/G \rightarrow \mathfrak{t}^*$ pulls back $C^\infty(\mathfrak{t}^*)$ to a *maximal* commutative subalgebra \mathcal{I}_T of $(C^\infty(M/G), \{, \})$. The map $M/G \rightarrow \mathfrak{g}^*/G$ pulls back $C^\infty(\mathfrak{g}^*/G)$ to a commutative Lie subalgebra \mathcal{I}_G of $(C^\infty(M/G), \{, \})$. As the two group actions commute so do the subalgebras \mathcal{I}_G and \mathcal{I}_T . By maximality, \mathcal{I}_T contains \mathcal{I}_G and consequently $M/G \rightarrow \mathfrak{g}^*/G$ factors through \mathfrak{t}^* .

The diagram of quotients for a Poisson action (diagram 2) generalizes to an analogous diagram for any full dual pair of realizations $P_2 \xleftarrow{f_2} M \xrightarrow{f_1} P_1$. Denote by \bar{M} the image of M in the Poisson manifold $P_1 \times P_2$ under the diagonal Poisson map $f_1 \times f_2 : M \rightarrow P_1 \times P_2$. The realization dual to $f_1 \times f_2 : M \rightarrow \bar{M}$ is the pullback of the symplectic leaf foliations on P_1 or P_2 (they pull back to the same foliation of M).

The following is the analogue of Example 2.14 replacing the commutative T -action by a realization:

Example 2.16 Let $M/G \xleftarrow{\pi} M \xrightarrow{\mu} \mathfrak{g}^*$ be the full dual pair associated to a Poisson action of G on M and $N \xleftarrow{\ell} M \xrightarrow{h} B$ a full dual pair of realizations with connected fibers where:

- (i) h is G -invariant
- (ii) $h : M \rightarrow B$ is a Hamiltonian map (B is endowed with the trivial Poisson structure) and
- (iii) The composition $M \xrightarrow{\mu} \mathfrak{g}^* \rightarrow \mathfrak{g}^*/G$ factors through $h : M \rightarrow B$.

Then we get a diagram analogous to the one in example 2.14:

$$\begin{array}{ccc}
 & M & \\
 & \downarrow & \\
 & \bar{M}_G & \\
 \swarrow \bar{\pi} & & \searrow N \\
 M/G & & \mathfrak{g}^* \\
 \searrow & \swarrow \bar{h} & \searrow \\
 & B & \\
 \searrow & & \swarrow \\
 & \mathfrak{g}^*/G & \\
 \downarrow & & \\
 & (0) &
 \end{array} \tag{5}$$

It follows that the Poisson map $M \xrightarrow{h \times \mu} B \times_{(\mathfrak{g}^*/G)} \mathfrak{g}^*$ into the fiber product space factors through the realization $M \xrightarrow{\ell} N$ dual to $h : M \rightarrow B$. If, moreover, $M/G \rightarrow B$ is a Lagrangian fibration, then $M \xrightarrow{h \times \mu} B \times_{(\mathfrak{g}^*/G)} \mathfrak{g}^*$ is itself a realization dual to $h : M \rightarrow B$.

2.5 Finite dimensional Poisson loop group actions

We present in this section two elementary constructions related to finite dimensional symplectic leaves in the Poisson quotient Q_∞ of an infinite dimensional symplectic space M by subgroups of loop groups. The material in this section will only be used in section 6.3 so the reader may prefer to read it in conjunction with that section.

We will not construct the quotient Q_∞ . The spaces involved are constructed independently. Rather, we will analyze the relationship between the Poisson action of the loop group on the infinite dimensional spaces and its descent to the finite dimensional symplectic leaves of Q_∞ . In fact, our main purpose in this section is to provide the terminology needed in order to study the Poisson loop group action in the finite dimensional setting (convention 2.20 and corollary 2.22).

In section 2.5.1 we note that the infinitesimal Hamiltonian actions of elements of the loop group descend to Hamiltonian vector fields on finite dimensional symplectic approximations $M_{(l,l)}$. The $M_{(l,l)}$'s dominate finite dimensional Poisson subvarieties Q_l of Q_∞ with positive dimensional fibers. In section 2.5.2 the action of certain maximal tori in the loop group further descends to finite Galois covers of certain (type) loci in Q and we examine the sense in which it is Hamiltonian.

2.5.1 Finite dimensional approximations

The loop group G_∞ is the group $GL(n, \mathbb{C}((z)))$. The level infinity group G_∞^+ is its positive part $GL(n, \mathbb{C}[[z]])$. Let (M, σ) be a symplectic variety with a Poisson loop group action whose moment map is

$$\mu : M \rightarrow \mathfrak{g}_\infty^*.$$

In section 6.3 M will be the cotangent bundle of a projective (inverse) limit of finite dimensional smooth algebraic varieties (the cotangent bundles of the moduli spaces of vector bundles with level structure). It is thus the inductive (direct) limit of projective limits of finite dimensional varieties. All constructions (morphisms, group actions, symplectic structures etc ...) can be made precise as limits of the standard constructions on finite dimensional approximations. We will omit the technical details as our point is to transfer the discussion back to the finite dimensional symplectic leaves of the Poisson quotient $Q_\infty := M/G_\infty^+$.

Let $G_\infty^{>l}$, $l \geq -1$, be the subgroup of G_∞^+ of elements equal to 1 up to order l . Denote by $\mu_{G_\infty^{>l}}$ its moment map. We assume that the subquotients

$$M_{(l,k)} := \mu_{G_\infty^{>l}}^{-1}(0)/G_\infty^{>k}, \quad k \geq l,$$

are smooth, finite dimensional and that they approximate M :

$$M = \lim_{l \rightarrow \infty} \lim_{\infty \leftarrow k} M_{(l,k)}.$$

Notice that $M_{(l,l)}$ is a symplectic reduction, hence symplectic.

Let a be an element of the loop algebra \mathfrak{g}_∞ with poles of order at most l_0 . The Hamiltonian vector field ξ_a on M is an infinite double sequence of Hamiltonian vector fields on $M_{(l,k)}$, $l \geq 0$, $k \geq \max\{l, l_0\}$ compatible with respect to projections and inclusions (by a Hamiltonian vector field on $M_{(l,k)} \subset M_{(k,k)}$ we mean, the restriction of a Hamiltonian vector field on $M_{(k,k)}$ which is tangent to $M_{(l,k)}$).

The quotient $Q_\infty := M/G_\infty^+$ is the direct limit $\lim_{l \rightarrow \infty} Q_l$ of the finite dimensional Poisson varieties

$$Q_l := \mu_{G_\infty^{>l}}^{-1}(0)/G_\infty^+ = M_{(l,k)}/G_k = M_{(l,l)}/G_l$$

where $G_k := G_\infty^+/G_\infty^{>k}$ is the finite dimensional level- k group (we assume that the quotients Q_l are smooth).

Example 2.17 The homogeneous G_∞^+ -space $\mathcal{U}_\infty := G_\infty^+/GL(n, \mathbb{C})$ is endowed with a canonical infinitesimal G_∞ -action via its embedding as the degree-0 component of the homogeneous G_∞ -space $G_\infty/GL(n, \mathbb{C}[[z^{-1}]])$

$$G_\infty^+/GL(n, \mathbb{C}) \hookrightarrow G_\infty/GL(n, \mathbb{C}[[z^{-1}]])$$

(the degree of $a \in G_\infty$ is the signed order of the pole/zero of $\det(a)$). Let M be an open subset of the cotangent bundle $T^*\mathcal{U}_\infty$ for which the regularity assumptions on the approximating quotients $M_{(l,k)}$ hold. This will be made precise in section 4.3 and the quotients Q_l will be the spaces of conjugacy classes of polynomial matrices studied in that section.

Unfortunately, the action of $a \in \mathfrak{g}_\infty$ above is not defined on Q_l . It is well defined only when we retain at least the l_0 -level structure, i.e., on $M_{(l,k)}$, $k \geq l_0$. In section 2.5.2 we will see that the action of certain maximal tori in G_∞ descends to *finite* Galois covers of certain loci in Q .

2.5.2 Type loci

Let (M, σ) be a smooth symplectic variety endowed with an infinitesimal Poisson action $\mu_G^* : \mathfrak{g} \rightarrow [\Gamma(M, \mathcal{O}_M), \{, \}]$ of a group G . Consider a subgroup $G^+ \subset G$, a commutative subgroup $T \subset G$, and their intersection $T^+ := T \cap G^+$. Assume further that the following conditions hold:

- i) The infinitesimal G^+ -action integrates to a free action on M ,
- ii) T^+ is a maximal commutative subgroup whose Weyl group $W_{T^+} := N_{G^+}(T^+)/T^+$ is finite.

Definition 2.18 The *type* τ of T is the class of all commutative subgroups T' of G which are conjugate to T via an element of G^+ .

Let $W := [N_{G^+}(T^+) \cap N(T)]/T^+$ be the corresponding subgroup of both W_{T^+} and W_T . Denote by

$$\mathfrak{g}_\tau^* \subset \mathfrak{g}^*$$

(respectively, $\mathfrak{g}_T^* \subset \mathfrak{g}^*$) the subset of elements whose stabilizer (with respect to the coadjoint action) is a torus of type τ (respectively, precisely T).

Example 2.19 Let G be the loop group, G^+ the level infinity group and $T \subset G$ a maximal torus of type \underline{n} determined by a partition of the integer n (see section 6.1). In this case G^+ and T generate G . It follows that $W = W_T = W_{T^+}$ and the type τ is invariant throughout a coadjoint orbit in \mathfrak{g}^* .

Assume that a “nice” (Poisson) quotient $Q := M/G^+$ exists. Let

$$M^\tau := \mu_G^{-1}(\mathfrak{g}_\tau^*), \quad \text{and} \quad Q^\tau := M^\tau/G^+ \subset Q$$

be the loci of type τ . Note that for each T of type τ there is a canonical isomorphism

$$\mu_G^{-1}(\mathfrak{g}_T^*)/[N_{G^+}(T^+) \cap N(T)] \xrightarrow{\cong} Q^\tau \subset Q.$$

In particular, a choice of T of type τ determines a canonical W -Galois cover of Q^τ

$$\tilde{Q}^T := \mu_G^{-1}(\mathfrak{g}_T^*)/T^+. \tag{6}$$

All the \tilde{Q}^T of type τ are isomorphic (not canonically) to a fixed abstract W -cover \tilde{Q}^τ . Note that \tilde{Q}^T is a subset of M/T^+ . We get a canonical “section” (the inclusion)

$$s_T : \tilde{Q}^T \hookrightarrow M/T^+ \tag{7}$$

into a T -invariant subset. Consequently, we get an induced T -action on the Galois cover \tilde{Q}^T . The moment map μ_T is T -invariant, hence, descends to M/T^+ . Restriction to $s^T(\tilde{Q}^T)$ gives rise to a canonical map

$$\bar{\mu}_T : \tilde{Q}^T \rightarrow \mathfrak{t}^*. \tag{8}$$

The purpose of this section is to examine *the extent to which $\bar{\mu}_T$ is the moment map of the T -action with respect to the Poisson structure on Q* . In general, the G^+ -equivariant projection

$$j : \mathfrak{g}^* \twoheadrightarrow (\mathfrak{g}^+)^*$$

might *forget the type*. Coadjoint orbits $S \subset (\mathfrak{g}^+)^*$ may intersect nontrivially the images $j(\mathfrak{g}_\tau^*)$ of several types (e.g., take $S = 0$ in example 2.19 and observe that the kernel of j intersects coadjoint orbits of all types). Consequently, symplectic leaves Q_S of Q would intersect nontrivially several type loci Q^τ . If $Q_S^{\tau^{open}}$ is an open subvariety of Q_S of type τ (e.g., if Q is the disjoint union of finitely many type loci and τ is a *generic type*) then the corresponding open subvariety $\tilde{Q}_S^{\tau^{open}}$ of \tilde{Q}_S^T will be a symplectic variety. In this case the T -action on $\tilde{Q}_S^{\tau^{open}}$ is Poisson whose moment map $\bar{\mu}_T$ is given by (8).

The Galois W -covers \tilde{Q}_S^T of the nongeneric type loci in Q_S are not symplectic. Nevertheless, motivated by the fact that $\bar{\mu}_T$ can be extended canonically to M/T^+

$$\begin{array}{ccc}
 M & \xrightarrow{\mu_T} & \mathfrak{t}^* \\
 \downarrow & \nearrow & \nearrow \\
 M/T^+ & \xleftarrow{\supset} & \tilde{Q}_S^T \\
 \downarrow & & \downarrow \\
 Q & \xleftarrow{\supset} & Q_S^\tau
 \end{array} \tag{9}$$

we will adopt the:

Convention 2.20 i) Given an element h of \mathfrak{t} we will say that the corresponding vector field $\bar{\xi}_h$ on \tilde{Q}_S^T is the *Hamiltonian vector field of h* (even if the type τ of T is not generic in Q_S). ii) We will refer to the pair $(\bar{\mu}_T, \mu_T)$ as the moment map of the T -action on \tilde{Q}_S^T .

Remarks 2.21 Let G be the loop group and M, G^+, T as in section 2.5.1,

1. Diagram (9) has an obvious finite dimensional approximation in which Q_S^τ, \tilde{Q}_S^T and T stay the same but with M replaced by $M_{(l,l)}$ and Q by Q_l . By μ_T we mean in this context a linear homomorphism $\mu_T^* : \mathfrak{t} \rightarrow \mathfrak{t}/\mathfrak{t}_\infty^{>l} \rightarrow \Gamma(M_{(l,l)}, \mathcal{O}_{M_{(l,l)}})$.
2. (Relation with the diagram of quotients (2)) Let S be a coadjoint orbit of level l , i.e., $S \subset \mathfrak{g}_l^* := (\mathfrak{g}_\infty^+/\mathfrak{g}_\infty^{>l})^* \subset (\mathfrak{g}_\infty^+)^*$. There is a rather subtle relationship between the Galois cover $\tilde{Q}_S^T \rightarrow Q_S^\tau$ and the space $\bar{M}_{(l,l)}$ dual to \mathfrak{g}_l^*/G_l from the diagram of quotients (2) of level l . The Galois cover $\tilde{Q}_S^T \rightarrow Q_S^\tau$ factors canonically through an intermediate subspace \tilde{Q}_S^T/\sim of $\bar{M}_{(l,l)}$. Note that the loop group moment map μ_{G_∞} descends to a map

$$\bar{\mu}_{G_\infty} : \tilde{Q}_S^T \rightarrow (\mathfrak{g}_\infty^*)_T \subset \mathfrak{g}_\infty^*.$$

Two points $\tilde{x}_1, \tilde{x}_2 \in \tilde{Q}_S^T$ in a fiber over $x \in Q_S^\tau$ are identified in \tilde{Q}_S^T/\sim if and only if $\bar{\mu}_{G_\infty}(\tilde{x}_1)$ and $\bar{\mu}_{G_\infty}(\tilde{x}_2)$ project to the same point in $S \subset (\mathfrak{g}_\infty^+)^*$. The relation \sim is a geometric realization of the partial type-forgetfulness of the projection $j : \mathfrak{g}_\infty^* \rightarrow (\mathfrak{g}_\infty^+)^*$. The loci in $\bar{M}_{(l,l)}$ at which the type is not forgotten are precisely the loci to which the moment map of the infinitesimal *loop group* action descends (from $M^T := \mu_{G_\infty}^{-1}((\mathfrak{g}_\infty^*)_T)$). Note that the moment map of the level infinity subgroup descends by definition of the quotient $\bar{M}_{(l,l)}$. In particular, the infinitesimal action of the maximal torus \mathfrak{t} integrates to a Poisson action in these loci. (See section 6.3.2 for examples of such loci.)

Assume further that we have a “nice” quotient $B := M/G$ and that G is generated by T and G^+ . We get the type loci $B^\tau := M^\tau/G$. Fixing T of type τ we get the W -cover $\tilde{B}^T := M^T/T$. The restriction of the moment map μ_T to M^T descends further to $\phi_T : \tilde{B}^T \rightarrow \mathfrak{t}^*$ and we get the commutative diagram

$$\begin{array}{ccc}
 & M^T & \\
 \swarrow & & \searrow \mu_G \\
 \tilde{Q}^T & & \mathfrak{g}_T^* \\
 \swarrow \tilde{h} & & \searrow \\
 \tilde{B}^T & \xrightarrow{\phi_T} & \mathfrak{t}^*
 \end{array} \tag{10}$$

Corollary 2.22 *i) The T -action on M descends to a canonical action on the W -cover \tilde{Q}^T of the type locus $Q^\tau \subset Q$. ii) Its moment map, in the sense of convention 2.20, is $(\phi_T \circ \tilde{h}, \mu_T)$. iii) If the type τ of T is the generic type in a symplectic leaf $Q_S \subset Q$ and $Q_S^{\tau \text{ open}} \subset Q_S$ is an open subvariety, then the corresponding W -cover $\tilde{Q}_S^{\tau \text{ open}}$ is symplectic and $\phi_T \circ \tilde{h}$ is the moment map of the T -action in the usual sense.*

3 Geodesic flow on an ellipsoid

Consider the geodesic flow on an ellipsoid

$$E = \{(x_1, \dots, x_{n+1}) \mid \sum_{i=1}^{n+1} \frac{1}{a_i} x_i^2 = 1\} \subset \mathbb{R}^{n+1},$$

where the metric is induced from the standard one on \mathbb{R}^{n+1} , and where the a_i are distinct positive numbers, say

$$0 < a_1 < \dots < a_{n+1}.$$

For $n = 1$, the problem is to compute arc length on an ellipse. It amounts to computing the integral

$$s = \int \sqrt{\frac{a_1^2 + (a_2 - a_1)x^2}{a_1(a_1 - x^2)}} dx.$$

(Hence the name *elliptic* for this and similar integrals.)

For $n = 2$, the problem was solved by Jacobi. Each geodesic γ on E determines a hyperboloid E' , intersecting E in a pair of ovals. The geodesic γ oscillates in the band between these ovals, meeting them tangentially. In fact, each tangent line of γ is also tangent to the hyperboloid E' . The solutions can be parametrized explicitly in terms of hyperelliptic theta functions.

The geodesic flow on an n -dimensional ellipsoid is integrable, in fact algebraically integrable. We will see this, first using some elementary geometric techniques to describe the geodesics concretely, and then again using the algebraic description of hyperelliptic jacobians which will be extended later to all spectral curves.

3.1 Integrability

The geodesic flow on the $2n$ -dimensional symplectic manifold $TE \approx T^*E$ is given by the Hamiltonian function $h = \text{length}^2$. We need $n - 1$ farther, commuting, independent, Hamiltonians.

Consider the family of quadrics confocal to E :

$$E_\lambda : \sum_{i=1}^{n+1} \frac{x_i^2}{a_i - \lambda} = 1,$$

depending on a parameter λ . (The name makes sense only when $n = 1$: we get the family of ellipses ($\lambda < a_1$), hyperbolas ($a_1 < \lambda < a_2$), and empty (real) conics ($\lambda > a_2$), with fixed foci.)

Here is an intrinsic way to think of this family. Start with a linear pencil

$$Q_\lambda = Q_0 + \lambda Q_\infty, \quad (\lambda \in \mathbb{P}^1)$$

of quadrics in general position in projective space \mathbb{P}^{n+1} . By “general position” we mean that there are exactly $n + 2$ values of $\lambda \in \mathbb{P}^1$ such that Q_λ is singular, and for those λ , Q_λ is a cone (i.e. its singular locus, or vertex, is a single point).

Lemma. *A generic linear subspace $L \approx \mathbb{P}^{k-1}$ in \mathbb{P}^{n+1} is tangent to Q_λ for k values of λ . The points of tangency p_λ are pairwise harmonic with respect to each of the quadrics Q_μ .*

Proof: Four points of \mathbb{P}^1 are harmonic if their cross ratio is -1 ; e.g. $0, \infty, a, -a$. Two points $p_1, p_2 \in \mathbb{P}^1$ are harmonic with respect to a quadric Q if the set $\{p_1, p_2\} \cup (Q \cap \mathbb{P}^1)$ is harmonic. For example, two points on the line at infinity in \mathbb{P}^2 (i.e. two directions in the affine plane) are harmonic with respect to some (hence every) circle, iff the directions are perpendicular.

Since Q is tangent to L if and only if $Q \cap L$ is singular, the first part of the lemma follows by restriction of the pencil to L . The second part follows by restricting to the line $p_{\lambda_1}, p_{\lambda_2}$, where in appropriate coordinates $Q_{\lambda_1} = x^2$ and $Q_{\lambda_2} = y^2$, so the points of tangency are $0, \infty$ and the quadric Q_λ vanishes at $\pm a$, where $a^2 = \lambda$.

□

We choose the parameter λ so that Q_∞ is one of the singular quadrics. The dual Q_λ^* of a non-singular Q_λ is a non-singular quadric in $(\mathbb{P}^{n+1})^*$. The dual of Q_∞ is a hyperplane $H_\infty \subset (\mathbb{P}^{n+1})^*$ (corresponding to the vertex of Q_∞), with a non-singular quadric $Q_\infty^* \subset H_\infty$. We get a family of confocal quadrics by restriction to the affine space $\mathbb{R}^{n+1} := (\mathbb{P}^{n+1})^* \setminus H_\infty$:

$$E_\lambda := Q_\lambda^*|_{\mathbb{R}^{n+1}}.$$

If we choose coordinates so that

$$\begin{aligned} Q_\infty &= \sum_{i=1}^{n+1} x_i^2 \\ Q_0 &= \sum_{i=0}^n a_i x_i^2 \end{aligned}$$

(where $a_0 = -1$ and the other a_i are as above), we retrieve the original E_λ . (Euclidean geometry in \mathbb{R}^{n+1} is equivalent, in the sense of Klein's program, to the geometry of \mathbb{P}^{n+1} with a distinguished "light-cone" Q_∞ . In this equivalence, $E = E_0$ corresponds to Q_0 , which determines the pencil $\{Q_\lambda\}$, which corresponds to the confocal family $\{E_\lambda\}$.)

Dualizing the lemma, for $k = n + 1, n$, gives the following properties of the confocal family. (The reader is invited to amuse herself by drawing the case $n = 1$ in the plane.)

- (1) Through a generic point x of \mathbb{R}^{n+1} pass $n + 1$ of the E_λ .
- (2) These $n + 1$ quadrics intersect perpendicularly at x .
- (3) A generic line ℓ in \mathbb{R}^{n+1} is tangent to n of the E_λ .
- (4) The tangent hyperplanes to these n quadrics (at their respective points of tangency to ℓ) are perpendicular.

By property (3) we can associate to a generic line ℓ in \mathbb{R}^{n+1} an unordered set of n values $\lambda_i, 1 \leq i \leq n$, such that Q_{λ_i} is tangent to ℓ . When ℓ comes from a point of TE , one of these, say λ_n , equals 0. The remaining $n - 1$ values λ_i (or rather, their symmetric functions) give $n - 1$ independent functions on TE ; in fact, they can take an arbitrary $(n - 1)$ -tuple of values. These functions descend to the projectivized tangent bundle $\mathbb{P}(TE)$; so together with the original Hamiltonian h (= length squared) they give n independent functions on TE . The key to integrability is:

Chasles' Theorem. *The λ_i are flow invariants, i.e. they are constant along a geodesic $\gamma = \gamma(t)$.*

Proof. For any curve $\gamma(t)$ in \mathbb{R}^{n+1} , the family of tangent lines $\ell(t)$ gives a curve Λ in the Grassmannian $Gr(1, \mathbb{R}^{n+1})$ of affine lines in \mathbb{R}^{n+1} . This curve is developable, i.e. its tangent line $T_{\ell(t)}\Lambda$ is given by the pencil of lines through $\gamma(t)$ in the osculating plane of γ at t . When γ is a geodesic, this plane is the span of $\ell(t)$ and the normal vector $n(t)$ to E at $\gamma(t)$. Write $\lambda_i(t)$ for the value of λ_i at $\ell(t)$.

Let Z_i be the hypersurface in $Gr(1, \mathbb{R}^{n+1})$ parametrizing lines tangent to $E_{\lambda_i(t)}$, for some fixed t . The tangent space $T_{\ell(t)}Z_i$ contains all lines through $\gamma(t)$ in the tangent hyperplane $T_{p_i(t)}E_{\lambda_i(t)}$, and this hyperplane contains the normal $n(t)$, by property (4). Hence:

$$T_{\ell(t)}\Lambda \subset \{\text{lines through } \gamma(t), \text{ in } T_{p_i(t)}E_{\lambda_i(t)}\} \subset T_{\ell(t)}Z_i \quad i = 1, \dots, n-1.$$

If the family Λ of tangent lines to a geodesic meets Z_i , it must therefore stay in it.

□

3.2 Algebraic integrability

Since a line ℓ determines two (opposite) tangent vectors of given non-zero length, we have identified the fiber of the geodesic flow as a double cover \tilde{K} of

$$K := \{\ell \in Gr(1, \mathbb{R}^{n+1}) \mid \ell \text{ is tangent to } E_{\lambda_1}, \dots, E_{\lambda_n}\}.$$

Next we want to interpret this in terms of the real points of a complex abelian variety. We follow Knörrer's approach [Kn], which in turn is based on [Mo],[Re] and [D5].

Start with the pencil of quadrics in \mathbb{P}^{2n+1} (over \mathbb{C}):

$$Y_\lambda := Y_0 + \lambda Y_\infty$$

with

$$\begin{aligned} Y_0 &= \sum_{i=1}^{2n+1} a_i x_i^2 - x_0^2 \\ Y_\infty &= \sum_{i=1}^{2n+1} x_i^2. \end{aligned}$$

The base locus $X = Y_0 \cap Y_\infty$ is non-singular if the a_i are distinct. We set $a_0 = \infty$. The family of linear subspaces \mathbb{P}^n contained in a fixed quadric Y_λ consists of two connected components, or rulings, for the non-singular Y_λ ($\lambda \notin \{a_i\}$), and of a single ruling for $\lambda = a_i$. We thus get a double cover

$$\pi : C \rightarrow \mathbb{P}^1$$

of the λ -line, parametrizing the rulings. (More precisely, one considers the variety of pairs

$$\mathcal{P} = \{(A, \lambda) \mid A \text{ is a } \mathbb{P}^n \text{ contained in } Q_\lambda\},$$

and takes the Stein factorization of the second projection.) Explicitly, C is the hyperelliptic curve of genus n :

$$C : s^2 = \prod_{i=1}^{2n+1} (t - a_i).$$

Miles Reid [Re] showed that the Jacobian $J(C)$ is isomorphic to the variety

$$F := \{A \in Gr(n-1, \mathbb{P}^{2n+1}) \mid A \subset X\}$$

of linear subspaces in the base locus. An explicit group law on F is given in [D5], and corresponding results for rank 2 vector bundles on C are in [DR]. Since we are interested in a family of varieties F with varying parameters, we need some information about the isomorphism. Let $Pic^d(C)$ denote the variety parametrizing isomorphism classes of degree- d line bundles on C . Then $Pic^0(C) = J(C)$ is a group, $Pic^1(C)$ is a torser (= principal homogeneous space) over it, but, these two have no natural identification; while $Pic^2(C) \approx J(C)$ canonically, using the hyperelliptic bundle on C . It turns out that F is isomorphic to $Pic^0(C)$ and to $Pic^1(C)$, but neither isomorphism is canonical. Rather, we may think of F as “ $Pic^{\frac{1}{2}}(C)$ ”: it is a torser over $J(C)$, and has a natural torser map

$$F \times F \rightarrow Pic^1(C).$$

All of this is based on the existence of a natural morphism

$$j : F \times C \rightarrow F.$$

The ruling p (on the quadric $Y_{\pi(p)}$) contains a unique subspace \mathbb{P}^n which contains a given \mathbb{P}^{n-1} -subspace $A \in F$. This \mathbb{P}^n intersects X in the union of A and another element of F , which we call $j(A, p)$. We can also think of j as a family of involutions of F , indexed by $p \in C$. This extends to a map

$$F \times J(C) \rightarrow F$$

which gives the torser structure on F . Once F is thus identified with $J(C)$, the map j becomes

$$j(A, p) = p - A,$$

up to an additive constant. Since this is well defined globally, points $A \in F$ must behave as line bundles on C of “degree $\frac{1}{2}$ ”. In particular, we have for $0 \leq i \leq 2n+1$ the involution

$$\begin{aligned} j_i : F &\rightarrow F \\ A &\mapsto j(A, a_i), \end{aligned}$$

where we set $a_i = \infty$ and identify the $a_i \in \mathbb{P}^1$ with the $2n+2$ Weierstrass points $\pi^{-1}(a_i) \in C$. Explicitly, each j_i is induced by the linear involution

$$\bar{j}_i : \mathbb{P}^{2n+1} \rightarrow \mathbb{P}^{2n+1}$$

flipping the sign of the i -th coordinate.

Consider the linear projection

$$\begin{aligned} \rho : \mathbb{P}^{2n+1} &\rightarrow \mathbb{P}^{n+1} \\ (x_0, \dots, x_{2n+1}) &\mapsto (x_0, \dots, x_{n+1}), \end{aligned}$$

which commutes with the \bar{j}_i , $n+2 \leq i$. Recall that in \mathbb{P}^{n+1} we have the pencil of quadrics Q_λ , with dual quadrics E_λ in $(\mathbb{P}^{n+1})^*$.

Proposition.

(i) The projection ρ maps F to

$$F' := \{B \in Gr(n-1, \mathbb{P}^{n+1}) \mid B \text{ is tangent to } Q_{n+2}, \dots, Q_{2n+1}\}.$$

(ii) The induced $\rho : F \rightarrow F'$ is a finite morphism of degree 2^n , and can be identified with the quotient of F by the group $G \approx (\mathbb{Z}/2\mathbb{Z})^n$ generated by the involutions j_i , $n+2 \leq i \leq 2n+1$.

(iii) Duality takes F' isomorphically to the variety K of lines in $(\mathbb{P}^{n+1})^*$ tangent to E_{λ_i} , $\lambda_i = a_{n+1+i}$, $1 \leq i \leq n$.

We omit the straightforward proof. Let $\tilde{G} \subset G$ be the index-2 subgroup generated by the products $j_{i_1} \circ j_{i_2}$, and set

$$\tilde{K} := F/\tilde{G}.$$

We obtain natural commuting maps, whose degrees are indicated next to the arrows:

$$\begin{array}{ccccc} F & \xrightarrow{2^{n-1}} & \tilde{K} & \xrightarrow{2^{n+1}} & Pic^1(C) \\ & & \downarrow^2 & & \downarrow^2 \\ & & K & \xrightarrow{2^{n+1}} & Kummer^1(C). \end{array}$$

Here $Kummer^d(C)$ stands for the quotient of $Pic^d C$ by the involution

$$L \mapsto dH - L,$$

where H is the hyperelliptic bundle $\in Pic^2(C)$. The composition of the maps in the top row is multiplication by 2:

$$F \approx Pic^{\frac{1}{2}}(C) \xrightarrow{\cdot 2} Pic^1(C).$$

In conclusion, the fiber of the geodesic flow on $E = E_0$ with invariants $h = 1$ (say) and $\lambda_i = a_{n+1+i}$, $1 \leq i \leq n$ can be identified with the real locus in $\tilde{K} = \tilde{K}(\lambda_1, \dots, \lambda_n)$. The latter is a 2^{n+1} -sheeted cover of $Pic^1(C)$, so up to translation by some points of order 2, it is an abelian variety, isomorphic to a 2^{n+1} sheeted cover of the hyperelliptic Jacobian $J(C)$.

3.3 The flows

Two details of the above story are somewhat unsatisfactory: First, the asymmetry between the $n-1$ Hamiltonians λ_i and the remaining Hamiltonian H (length squared). And second, the fact that the complexified total space TE of the system is not quite symplectic. Indeed, for an arbitrary algebraic hypersurface $M \subset \mathbb{C}^{n+1}$, given by $f = 0$, the complexified metric on \mathbb{C}^{n+1} induces bundle maps

$$TM \hookrightarrow T\mathbb{C}^{n+1}|_M \xrightarrow{\sim} T^*\mathbb{C}^{n+1}|_M \twoheadrightarrow T^*M,$$

but the composition is not an isomorphism; rather, it is degenerate at points where

$$0 = (\nabla f)^2 = \sum_{k=1}^{n+1} \left(\frac{\partial f}{\partial x_k} \right)^2.$$

For an ellipsoid $\sum x_k^2/a_k = 1$ (other than a sphere) there will be an empty real, but non-empty complex degeneracy locus, given by the equation $\sum (x_k/a_k)^2 = 0$.

Both of these annoyances disappear if we replace the total space by the tangent bundle TS of the sphere

$$S = \{(x_1, \dots, x_{n+1}) \in \mathbb{C}^{n+1} \mid \sum x_k^2 = 1\},$$

i.e.

$$TS = \{(x, y) \in \mathbb{C}^{2n+2} \mid \sum x_k^2 = 1, \sum x_k y_k = 0\}.$$

This is globally symplectic, and the n (unordered) Commuting Hamiltonians can be taken to be the values λ_i , $1 \leq i \leq n$, such that the line

$$\ell_{x,y} := (\text{line through } y \text{ in direction } x)$$

is tangent to E_{λ_i} . The original system TE can be recovered as a \mathbb{C}^* -bundle (where \mathbb{C}^* acts by rescaling the tangent direction x) over the hypersurface $\lambda = 0$ in TS .

Here is the explicit equation of the hypersurface:

$$\begin{aligned} \lambda = 0 &\Leftrightarrow \ell_{x,y} \text{ is tangent to } E = \{\sum \frac{x_k^2}{a_k} = 1\} \\ &\Leftrightarrow -1 + \sum \frac{(y_k + tx_k)^2}{a_k} = 0 \text{ has a unique solution } t \\ &\Leftrightarrow (-1 + \sum \frac{y_k^2}{a_k} + 2t(\sum \frac{x_k y_k}{a_k}) + t^2(\sum \frac{x_k^2}{a_k})) = 0 \text{ has a unique solution} \\ &\Leftrightarrow 0 = (\sum \frac{x_k y_k}{a_k})^2 - (\sum \frac{x_k^2}{a_k})(-1 + \sum \frac{y_k^2}{a_k}). \end{aligned}$$

More generally, this computation shows that $\ell_{x,y}$ is tangent to E_λ if and only if

$$0 = (\sum \frac{x_k y_k}{a_k - \lambda})^2 - (\sum \frac{x_k^2}{a_k - \lambda})(-1 + \sum \frac{y_k^2}{a_k - \lambda}) = \sum_k \frac{x_k^2}{a_k - \lambda} + \sum_{k \neq l} \frac{x_k y_k x_l y_l - x_k^2 y_l^2}{(a_k - \lambda)(a_l - \lambda)}.$$

As a function of λ , the last expression has first order poles at $\lambda = a_k$, $1 \leq k \leq n+1$, so it can be rewritten as

$$\sum \frac{1}{a_k - \lambda} F_k(x, y)$$

where the F_k are found by taking residue at $\lambda = a_k$:

$$F_k(x, y) := x_k^2 + \sum_{\ell \neq k} \frac{(x_k y_\ell - x_\ell y_k)^2}{a_k - a_\ell}.$$

We see that fixing the $n+1$ values $F_k(x, y)$, $1 \leq k \leq n+1$, subject to the condition

$$\sum_{k=1}^{n+1} F_k = 1,$$

is equivalent to fixing the n (unordered) values λ_i , $1 \leq i \leq n$.

This determines the hyperelliptic curve

$$C : s^2 = \prod_{k=1}^{n+1} (t - a_k) \cdot \prod_{i=1}^n (t - \lambda_i),$$

and the corresponding abelian variety

$$\widetilde{K} = \widetilde{K}(\lambda_1, \dots, \lambda_n) = J(C)/\widetilde{G} \approx \{(x, y) | \ell_{x,y} \text{ is tangent to } E_{\lambda_i}, \quad 1 \leq i \leq n\}.$$

Theorem.

- (1) *Geodesic flow on the quadric E_{λ_i} is the Hamiltonian vector field on TS given by the (local) Hamiltonian λ_i . On \widetilde{K} it is a constant vector field in the direction of the Weierstrass point $\lambda_i \in C$.*
- (2) *The Hamiltonian vector field on TS with Hamiltonian F_k is constant on \widetilde{K} , in the direction of the Weierstrass point $a_k \in C$.*

The direction at $\ell \in \widetilde{K}$ of geodesic flow on E_{λ_i} was described in the proof of Chasles' theorem. The direction given by the Weierstrass point λ_i is given at $A \in F$ as the tangent vector at λ_i to the curve

$$p \mapsto j(A_i, p) \quad (\text{where } A_i = j(A, \lambda_i)).$$

The proof of (1) amounts to unwinding the definitions to see that these two directions agree. (For details, see [Kn] and [D5].) Since the level sets of the λ_i and the F_k are the same, the Hamiltonian vector field of F_k evolves on the same \widetilde{K} , and is constant there. A monodromy argument on the family of hyperelliptic curves then shows that its direction must agree with a_k . Mumford gives an explicit computation for this in [Mum2], Theorem 4.7, following Moser [Mo].

We have identified the flows corresponding to $2n + 1$ of the Weierstrass points. The remaining one, at $\lambda = \infty$, corresponds to the Hamiltonian

$$H = \frac{1}{2} \sum_{k=1}^{n+1} a_k F_k = \frac{1}{2} \sum a_k x_k^2 + \frac{1}{2} \sum y_k^2,$$

giving Neumann's system, which is the starting point for the analysis in [Mo] and [Mum2].

3.4 Explicit parametrization

Fix the hyperelliptic curve of genus n

$$C : s^2 = f(t) := \prod_{i=1}^{2n+1} (t - a_i),$$

with projection

$$\begin{aligned} \pi : C &\rightarrow \mathbb{P}^1 \\ (t, s) &\mapsto t \end{aligned}$$

and involution

$$\begin{aligned} i : C &\rightarrow C \\ (t, s) &\mapsto (t, -s). \end{aligned}$$

We identify the various components $Pic^d(C)$ by means of the base point ∞ , which is a Weierstrass point. The affine open subset $J(C) \setminus \Theta$ can be described geometrically, by Riemann's theorem:

$$\begin{aligned} J(C) \setminus \Theta &\approx \\ \{L \in Pic^{n-1}(C) | h^0(L) = 0\} &\approx \\ \{(p_1, \dots, p_n) \in Sym^n C | p_i \neq \infty, p_i \neq i(p_j)\}, & \end{aligned}$$

where the last identification sends L to the unique effective divisor of $L(\infty)$. Mumford [Mum2] gives an explicit algebraic parametrization of the same open set, which he attributes to Jacobi: to the n -tuple $D = (p_1, \dots, p_n)$ he associates three polynomials of a single variable t :

(i) $U(t) := \prod_{i=1}^n (t - t(p_i)).$

(ii) $V(t)$ is the unique polynomial of degree $\leq n - 1$ such that the meromorphic function

$$V \circ \pi - s : C \rightarrow \mathbb{P}^1$$

vanishes on the divisor $D \subset C$. It is obtained by Lagrange interpolation of the expansions of s at the p_i , e.g. when the p_i are all distinct,

$$V(t) = \sum_{i=1}^n s(p_i) \prod_{j \neq i} \frac{t - t(p_j)}{t(p_i) - t(p_j)}.$$

(iii) $W(t) = \frac{f(t) - V(t)^2}{U(t)}$; the definition of V and the equation $s^2 = f(t)$ guarantee that this is a monic polynomial of degree $n + 1$.

Conversely, the polynomials U, V, W determine the values $t(p_i), s(p_i)$, hence the divisor D . By reading off the coefficients, we obtain an embedding:

$$(U, V, W) : \mathcal{J}(C) \setminus \Theta \hookrightarrow \mathbb{C}^{3n+1}.$$

The image is

$$\{(U, V, W) | V^2 + UW = f\}.$$

This description fits beautifully with the integrable system on TS representing geodesic flow on the E_λ . We can rephrase our previous computation as:

$$\ell_{x,y} \text{ is tangent to } E_{\lambda_1}, \dots, E_{\lambda_n} \Leftrightarrow f_1(t)f_2(t) = UW + V^2,$$

where, for $(x, y) \in TS$, we set:

$$\begin{aligned} f_1(t) &= \prod_{k=1}^{n+1} (t - a_k) && \text{(this is independent of } x, y) \\ f_2(t) &= f_1(t) \cdot \sum_k \frac{F_k(x, y)}{t - a_k} && \text{(this varies with } x, y; \text{ the roots are the } \lambda_i) \\ U(t) &= f_1(t) \left(\sum_k \frac{x_k^2}{t - a_k} \right) \\ W(t) &= f_1(t) \left(1 + \sum_k \frac{y_k^2}{t - a_k} \right) \\ V(t) &= \sqrt{-1} \cdot f_1(t) \cdot \left(\sum_k \frac{x_k y_k}{t - a_k} \right) \end{aligned}$$

The entire system TS is thus mapped to \mathbb{C}^{3n+1} . Each abelian variety $\widetilde{K} = \widetilde{K}(\lambda_1, \dots, \lambda_n)$ is mapped to $\mathcal{J}(C) \setminus \Theta$ embedded in \mathbb{C}^{3n+1} as before, where C is defined by $s^2 = f(t)$, and $f = f_1 \cdot f_2$, with f_1 fixed (of degree $n+1$) and f_2 variable (of degree n). On each \widetilde{K} the map is of degree 2^{n+1} ; the group $(\mathbb{Z}/2\mathbb{Z})^{n+1}$ operates by sending

$$(x_k, y_k) \mapsto (\epsilon_k x_k, \epsilon_k y_k), \quad \epsilon_k = \pm 1.$$

4 Spectral curves and vector bundles

We review in this chapter a general construction of an integrable system on the moduli space of Higgs pairs (E, φ) consisting of a vector bundle E on a curve and a meromorphic 1-form valued endomorphism φ (theorem 4.8). These moduli spaces admit a natural foliation by Jacobians of spectral curves. The spectral curves are branched covers of the base curve arising from the eigenvalues of the endomorphisms φ .

We concentrate on two examples:

- The Hitchin system supported on the cotangent bundle of the moduli space of vector bundles on a curve (section 4.2), and
- An integrable system on the moduli space of conjugacy classes of polynomial matrices (section 4.3).

The latter is then used to retrieve the Jacobi-Moser-Mumford system which arose in chapter 3 out of the geodesic flow on an ellipsoid.

Both examples are endowed with a natural symplectic or Poisson structure. The general construction of the Poisson structure on the moduli spaces of Higgs pairs is postponed to chapter 5.

We begin with a short survey of some basic facts about vector bundles on a curve.

4.1 Vector Bundles on a Curve

We fix a (compact, non-singular) curve Σ of genus g . A basic object in these lectures will be the moduli space of stable (or semistable) vector bundles on Σ of given rank r and degree d . To motivate the introduction of this object, let us try to describe a “general” vector bundle on Σ . One simple operation which produces vector bundles from line bundles is the direct image: start with an r -sheeted branched covering $\pi : C \rightarrow \Sigma$, ramified at points of some divisor R in the non-singular curve C . Then any line bundle $L \in Pic C$ determines a rank- r vector bundle $E := \pi_* L$ on Σ . As a locally free sheaf of \mathcal{O}_Σ -modules of rank r , this is easy to describe

$$\Gamma(\mathcal{U}, \pi_* L) := \Gamma(\pi^{-1}\mathcal{U}, L),$$

for open subsets $\mathcal{U} \subset \Sigma$. As a vector bundle, the description is clear only at unbranched points of Σ : if $\pi^{-1}(p)$ consists of r distinct points p_1, \dots, p_r then the fiber of E at p is naturally isomorphic to the direct sum of the fibers of L :

$$E_p \approx \bigoplus_{i=1}^r L_{p_i}.$$

At branch points of π , E_p does not admit a natural decomposition, but only a filtration. This is reflected in a drop in the degree. Indeed, the Grothendieck-Riemann-Roch theorem says in our (rather trivial) case that

$$\chi(\pi_* L) = \chi(L),$$

where χ is the holomorphic Euler characteristic,

$$\chi(E) := \deg E - (g - 1) \text{rank } E = \deg E + \chi(\mathcal{O}) \cdot \text{rank } E.$$

Using Hurwitz' formula:

$$\chi(\mathcal{O}_C) = r \chi(\mathcal{O}_\Sigma) - \frac{1}{2} \deg R,$$

this becomes:

$$\deg E = \deg L - \frac{1}{2} \deg R.$$

Example 4.1 Consider the double cover

$$\begin{aligned} \pi : \mathbb{P}^1 &\longrightarrow \mathbb{P}^1 \\ w &\longrightarrow z = w^2 \end{aligned}$$

branched over $0, \infty$. The direct image of the structure sheaf is:

$$\pi_* \mathcal{O} \approx \mathcal{O} \oplus \mathcal{O}(-1).$$

We can think of this as sending a regular function $f = f(w)$ (on some invariant open set upstairs) to the pair $(f_+(z), f_-(z))$ downstairs, where

$$f(w) = f_+(w^2) + w f_-(w^2).$$

In the image we get all pairs with f_+ regular (i.e. a section of \mathcal{O}) and f_- regular and vanishing at ∞ (i.e. a section of $\mathcal{O}(-1)$). (Similar considerations show that

$$\pi_* \mathcal{O}(-1) \approx \mathcal{O}(-1) \oplus \mathcal{O}(-1)$$

and more generally:

$$\pi_* \mathcal{O}(d) \approx \mathcal{O}(\lfloor \frac{d}{2} \rfloor) \oplus \mathcal{O}(\lfloor \frac{d-1}{2} \rfloor).$$

Note that this has degree $d-1$, as expected). The structure of $\pi_* L$ near the branch point $z = 0$ can be described, in this case, by the action on the local basis a, b (of even, odd sections) of multiplication by the section w upstairs:

$$a \mapsto b, \quad b \mapsto za,$$

i.e. w is represented by the matrix

$$\begin{pmatrix} 0 & z \\ 1 & 0 \end{pmatrix}$$

whose square is $z \cdot I$. At a branch point where k sheets come together, the corresponding action (in terms of a basis indexed by the k -th roots of unity) is given by the matrix:

$$P_k := \begin{pmatrix} 0 & & 0 & z \\ 1 & & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & & \cdot & \cdot \\ \cdot & & \cdot & 0 \\ 0 & & 1 & 0 \end{pmatrix} \quad (11)$$

whose k -th power is $z \cdot I$.

Example 4.2 Now consider a 2-sheeted branched cover $\pi : C \rightarrow \mathbb{P}^1$, where $g(C) > 0$. If we take $\chi(L) = 0$, i.e. $\deg L = g - 1$, we get $\deg(\pi_*L) = -2$. The equality

$$\ell := h^0(C, L) = h^0(\mathbb{P}^1, \pi_*L)$$

implies

$$\pi_*L \approx \mathcal{O}(\ell - 1) \oplus \mathcal{O}(-\ell - 1).$$

In particular, we discover a very disturbing phenomenon: as the line bundle L varies continuously, in $\text{Pic}^{g-1}C$, so should presumably π_*L ; but if we consider a 1-parameter family of line bundles L_t such that

$$\begin{aligned} L_0 &\in \Theta \\ L_t &\notin \Theta, \quad t \neq 0, \end{aligned}$$

we see that the vector bundle π_*L_t jumps from its generic value, $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ to $\mathcal{O} \oplus \mathcal{O}(-2)$ at $t = 0$. Similar jumps can clearly be forced on a rank- r bundle by considering r -sheeted branched covers.

The moral of these examples is that if we want a moduli space parametrizing the “general” vector bundle on a curve and having a reasonable (say, separated) topology, we cannot consider *all* bundles. In the case of \mathbb{P}^1 , we will end up with only the balanced bundles such as $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$, thus avoiding the possibility of a discontinuous jump.

The slope $\mu(E)$ of a vector bundle E is defined by:

$$\mu(E) := \frac{\deg E}{\text{rank } E}.$$

A bundle E is called stable (resp., semistable) if for every subbundle $F \subset E$ (other than $0, E$),

$$\mu(F) < \mu(E), \quad (\text{resp. } \mu(F) \leq \mu(E)).$$

The basic result due to Mumford and Seshadri [Se], is that reasonable (coarse) moduli spaces $\mathcal{U}_\Sigma^s(r, d) \subset \mathcal{U}_\Sigma(r, d)$ exist, with the following properties:

- $\mathcal{U}_\Sigma^s(r, d)$ is smooth; its points parametrize isomorphism classes of stable bundles of rank r and degree d on Σ ; it is an open subset of $\mathcal{U}_\Sigma(r, d)$.
- $\mathcal{U}_\Sigma(r, d)$ is projective; its points parametrize equivalence classes of semistable bundles, where two bundles are equivalent, roughly, if they admit filtrations by semistable subbundles (of constant slope) with isomorphic graded pieces.
- Both are coarse moduli spaces; this means that any “family”, i.e. vector bundle on a product $S \times \Sigma$, where S is any scheme, whose restrictions E_s to copies $s \times \Sigma$ of Σ are (semi) stable of rank r and degree d , determines a unique morphism of S to $\mathcal{U}_\Sigma^s(r, d)$ (respectively, $\mathcal{U}_\Sigma(r, d)$) which sends each $s \in S$ to the isomorphism (resp. equivalence) class of E_s , and has the obvious functoriality properties.

Examples

$g = 0$. The stable bundles are the line bundles $\mathcal{O}(d)$. The semi-stable bundles are the balanced vector bundles, $\mathcal{O}(d)^{\oplus r}$. Thus $\mathcal{U}_{P^1}(r, d)$ is a point if $r \mid d$, empty otherwise, while the stable subset is empty when $r \neq 1$.

$g = 1$. Let $h := \gcd(r, d)$. Atiyah [At] shows that $\mathcal{U}_{\Sigma}(r, d)$ is isomorphic to the symmetric product $S^h \Sigma$, and that each semistable equivalence class contains a unique decomposable bundle $E = \bigoplus_{i=1}^h E_i$, where each E_i is stable of rank r/h and degree d/h . (Other bundles in this equivalence class are filtered, with the E_i as subquotients.) Thus when $h = 1$, $\mathcal{U}_{\Sigma}^s = \mathcal{U}_{\Sigma}$, and when $h > 1$, \mathcal{U}_{Σ}^s is empty.

The possibilities for semistable bundles are illustrated in the case $r = 2, d = 0$: given two line bundles $L_1, L_2 \in \text{Pic}^0 \Sigma$, the possible extensions are determined, up to non zero scalars, by elements of

$$\text{Ext}_{\mathcal{O}_{\Sigma}}^1(L_1, L_2) \approx H^1(L_2 \otimes L_1^{-1}).$$

The direct sum is thus the only extension when $L_1 \not\approx L_2$, while if $L_1 \approx L_2 \approx L$ there is, up to isomorphism, also a unique non-trivial extension, say E_L . There is, again, a jump phenomenon: by rescaling the extension class we get a family of vector bundles with generic member isomorphic to E_L and special member $L \oplus L$. This explains why there cannot exist a moduli space parametrizing *isomorphism* classes of semistable bundles; neither $L \oplus L$ nor E_L is excluded, and the point representing the former is in the closure of the latter, so they must be identified, i.e. E_L and $L \oplus L$ must be declared to be equivalent.

Higher Genus. The only other cases where an explicit description of $\mathcal{U}_{\Sigma}(r, d)$ is known are when $r = 2$ and $g = 3$ [NR] or $r = 2$ and Σ is hyperelliptic of any genus [DR]. In the latter case, the moduli space $\mathcal{U}_{\Sigma}(2, \xi)$ of rank 2 vector bundles with a fixed determinant line bundle ξ of odd degree is isomorphic to the family of linear spaces \mathbb{P}^{g-2} in the intersection of the two quadrics in \mathbb{P}^{2g+1} used in Chapter 3. In the even degree case, $\mathcal{U}_{\Sigma}(2, \xi)$ can also be described in terms of the same two quadrics; when $g = 2$, it turns out to be isomorphic to \mathbb{P}^3 , in which the locus of semistable but non-stable points is the Kummer surface $K := \mathcal{J}(\Sigma)/\pm 1$, with its classical embedding in \mathbb{P}^3 as a quadric with 16 nodes.

Elementary deformation theory lets us make some general statements about $\mathcal{U} := \mathcal{U}_{\Sigma}(r, d)$ and $\mathcal{U}^s := \mathcal{U}_{\Sigma}^s(r, d)$:

Lemma 4.3 *For $g \geq 2$:*

1. $\dim \mathcal{U} = 1 + r^2(g - 1)$, and \mathcal{U}^s is a dense open subset.
2. Stable bundles E are simple, i.e. the only (global) endomorphisms of E are scalars.
3. Stable bundles are non-singular points of \mathcal{U} .
4. At points of \mathcal{U}^s there are canonical identifications

$$\begin{aligned} T_E \mathcal{U}^s &\approx H^1(\text{End } E) \\ T_E^* \mathcal{U}^s &\approx H^0(w_{\Sigma} \otimes \text{End } E). \end{aligned}$$

The proof of (2) is based on the observation that any nonzero $\alpha : E \rightarrow E$ must be invertible, otherwise either $\ker \alpha$ or $\text{im } \alpha$ would violate stability. Therefore $H^0(\text{End } E)$ is a finite dimensional division algebra containing \mathbb{C} , hence equal to it. Since a vector bundle E on Σ is determined by a 1-cocycle with values in $GL(r, \mathcal{O}_\Sigma)$ (= transition matrices), a first order deformation of E is given by a 1-cocycle with values in the associated bundle of Lie algebras, i.e. (up to isomorphism) by a class in $H^1(\text{End } E)$. The functoriality property of \mathcal{U} (“coarse moduli space”) implies that this is the Zariski tangent space, $T_E \mathcal{U}$. By Riemann-Roch

$$h^1(\text{End } E) = r^2(g - 1) + h^0(\text{End } E).$$

so the minimal value is obtained at the simple points, and equals $1 + r^2(g - 1)$ as claimed in (1). The identification of $T_E^* \mathcal{U}^s$ follows from that of $T_E \mathcal{U}^s$ by Serre duality. □

4.2 Spectral Curves and the Hitchin System

The relation between vector bundles and finite dimensional integrable systems arises from Hitchin’s amazing result.

Theorem 4.4 [H1, H2] *The cotangent bundle to the moduli space of semistable vector bundles supports a natural ACIHS.*

At the heart of Hitchin’s theorem is a construction of a spectral curve associated to a 1-form valued endomorphism of a vector bundle. The spectral construction allows a uniform treatment of a wide variety of algebraically completely integrable Hamiltonian systems. We will concentrate in this section on the algebro-geometric aspects of these systems leaving their symplectic geometry to Chapter 8. We work with vector bundles over curves, other structure groups will be treated in Chapter 9. The reader is referred to [BNR] and [H2] for more details.

The total space of the cotangent bundle $T^* \mathcal{U}_\Sigma^s(r, d)$ of the moduli space of stable vector bundles parametrizes pairs (E, φ) consisting of a stable vector bundle E and a covector φ in $H^1(\Sigma, \text{End } E)^* \simeq H^0(\Sigma, \text{End } E \otimes \omega_\Sigma)$, i.e., a 1-form valued endomorphism of E .

Consider more generally a pair (E, φ) of a rank r vector bundle E and a section $\varphi \in \text{Hom}(E, E \otimes K)$ where K is a line bundle on Σ . The i -th coefficient b_i of the characteristic polynomial of (E, φ) is a homogeneous polynomial of degree i on K^{-1} , hence a section of $H^0(\Sigma, K^{\otimes i})$. In fact $b_i = (-1)^i \cdot \text{trace}(\wedge^i \varphi)$.

The Hamiltonian map of the Hitchin system is the characteristic polynomial map

$$H : T^* \mathcal{U}_\Sigma(r, d) \longrightarrow B_\omega := \bigoplus_{i=1}^r H^0(\Sigma, \omega^{\otimes i}).$$

The fibers of the Hitchin map H turn out to be Jacobians of curves associated canonically to characteristic polynomials.

Going back to the general K -valued pair (E, φ) , notice that a characteristic polynomial $\text{char } (\varphi) = y^r - \text{tr}(\varphi)y^{r-1} + \dots + (-1)^r \det \varphi$ in $B_K := \bigoplus_{i=1}^r H^0(\Sigma, K^{\otimes i})$ defines a morphism from the line bundle K to $K^{\otimes r}$. The inverse image C of the zero section in $K^{\otimes r}$ under a

polynomial P in B_K is called a spectral curve. If P is the characteristic polynomial of a pair (E, φ) then indeed the fibers of $\pi : C \rightarrow \Sigma$ consist of eigenvalues of φ . If $K^{\otimes r}$ has a section without multiple zeroes (e.g., if it is very ample) then the generic spectral curve is smooth.

Lagrange interpolation extends a function on the inverse image $\pi^{-1}(U) \subset C$ of an open set U in Σ to a unique function on the inverse image of U in the surface K which is a polynomial of degree $\leq r - 1$ on each fiber. It follows that the direct image $\pi_*\mathcal{O}_C$ is isomorphic to $\mathcal{O}_\Sigma \oplus K^{-1} \oplus \dots \oplus K^{1-r}$. Assuming that $K^{\otimes i}$ has no sections for $i < 0$, the genus $h^1(C, \mathcal{O}_C) = h^1(\Sigma, \pi_*\mathcal{O}_C)$ of C is equal to $\deg(K) \cdot r(r - 1)/2 + r(g - 1) + 1$. In particular, when $K = \omega_\Sigma$, the genus of C is equal to half the dimension of the cotangent bundle. The data (E, φ) determines moreover a sheaf L on the spectral curve which is a line bundle if the curve is smooth. Away from the ramification divisor R in C , L is the tautological eigenline subbundle of the pullback π^*E . More precisely, the homomorphism $(\pi^*(\varphi) - y \cdot I) : \pi^*E \rightarrow \pi^*(E \otimes K)$, where $y \in H^0(C, \pi^*K)$ is the tautological eigenvalue section, has kernel $L(-R)$.

Conversely, given a spectral curve C and a line bundle L on it we get a pair $(\pi_*L, \pi_*(\otimes y))$ of a rank r vector bundle on Σ and a K valued endomorphism (see example 4.1). The two constructions are the inverse of each other.

Proposition 4.5 [H2, BNR] *If C is an irreducible and reduced spectral curve there is a bijection between isomorphism classes of*

- Pairs (E, φ) with spectral curve C .
- Rank 1 torsion free sheaves L on C .

Under this correspondence, line bundles on C correspond to endomorphisms φ which are regular in every fiber, i.e., whose centralizer in each fiber is an r -dimensional subspace of the corresponding fiber of $\text{End } E$. (This notion of regularity agrees with the one in Example 2.2.)

We conclude that the fiber of the Hitchin map $H : T^*U_\Sigma^s(r, d) \rightarrow B_\omega$ over a characteristic polynomial $b \in B_\omega$ is precisely the open subset of the Jacobian $J_C^{d+r(1-g_\Sigma)+g_C-1}$ consisting of the line bundles L whose direct image is a stable vector bundle. Moreover, the construction of the characteristic polynomial map and a similar description of its fibers applies to moduli spaces of pairs with K -valued endomorphism where K need not be the canonical line bundle (Theorem 4.8).

The missing line bundles in the fibers of the Hitchin map indicate that we need to relax the stability condition for the pair (E, φ) .

Definition: A pair (E, φ) is stable (semistable) if the slope of every φ -invariant subbundle of E is less than (or equal) to the slope of E .

As in the case of vector bundles we can define an equivalence relation for semistable pairs, where two bundles are equivalent, roughly, if they admit φ -invariant filtrations by semistable pairs (of constant slope) with isomorphic graded pieces. Two stable pairs are equivalent if and only if they are isomorphic.

Theorem 4.6 [H1, Sim1, Nit] *There exists an algebraic coarse moduli scheme $\text{Higgs}_K := \text{Higgs}_\Sigma(r, d, K)$ parametrizing equivalence classes of semistable K -valued pairs.*

The characteristic polynomial map $H : \text{Higgs}_K \rightarrow B_K$ is a proper algebraic morphism.

A deeper reason for working with the above definition of stability is provided by the following theorem from nonabelian Hodge theory:

Theorem 4.7 [H1, Sim2] *There is a canonical real analytic diffeomorphism between*

- *The moduli space of conjugacy classes of semisimple representations of the fundamental group $\pi_1(\Sigma)$ in $GL(r, \mathbb{C})$ and*
- *The moduli space of semistable ω -valued (Higgs) pairs (E, φ) of rank r and degree 0.*

In the case of Hitchin's system ($K = \omega_\Sigma$), the symplectic structure of the cotangent bundle extends to the stable locus of the moduli space of Higgs pairs giving rise to an integrable system $H : \text{Higgs}_\Sigma(r, d, \omega_\Sigma) \rightarrow B_\omega$ whose generic fiber is a complete Jacobian of a spectral curve.

We will show in Chapter 6 that the Hitchin system is, in fact, the lowest rank symplectic leaf of a natural infinite dimensional Poisson variety $\text{Higgs}_\Sigma(r, d)$ obtained as an inductive limit of the moduli spaces $\text{Higgs}_\Sigma^{sm}(r, d, \omega(D))$ of $\omega(D)$ -valued pairs as D varies through all effective divisors on Σ . The basic fact, generalizing the results of [H2, BNR, B1] is:

Theorem 4.8 [Bn, Ma1] *Let D be an effective divisor (not necessarily reduced) on a smooth algebraic curve Σ of genus g . Assume that $[\omega(D)]^{\otimes r}$ is very ample and if $g = 0$ assume further that $\deg(D) > \max(2, \rho)$ where $0 \leq \rho < r$ is the residue of d mod r . Then*

1. *The moduli space $\text{Higgs}_\Sigma^s(r, d, \omega(D))$ of stable rank r and degree d $\omega(D)$ -valued Higgs pairs has a smooth component $\text{Higgs}_\Sigma^{sm}(r, d, \omega(D))$ of top dimension $r^2(2g - 2 + \deg(D)) + 1 + \epsilon_{D=0}$, where $\epsilon_{D=0}$ is 1 if $D = 0$ and zero if $D > 0$. $\text{Higgs}_\Sigma^{sm}(r, d, \omega(D))$ is the unique component which contains Higgs pairs supported on irreducible and reduced spectral curves.*
2. *$\text{Higgs}_\Sigma^{sm}(r, d, \omega(D))$ has a canonical Poisson structure.*
3. *The characteristic polynomial map $H : \text{Higgs}_\Sigma^{sm}(r, d, \omega(D)) \rightarrow B_{\omega(D)}$ is an algebraically completely integrable Hamiltonian system. The generic (Lagrangian) fiber is a complete Jacobian of a smooth spectral curve of genus $r^2(g - 1) + 1 + (\deg D)(\frac{r(r-1)}{2})$.*
4. *The foliation of $\text{Higgs}_\Sigma^{sm}(r, d, \omega(D))$ by closures of top dimensional symplectic leaves is induced by the cosets of*

$$H^0 \left(\Sigma, \left[\bigoplus_{i=1}^r \omega_\Sigma(D)^{\otimes i} \right] (-D) \right) \text{ in } B_{\omega(D)}.$$

Definition 4.9 *As in the theorem, we will denote by $\text{Higgs}_\Sigma^{sm}(r, d, \omega(D))$ the unique component which contains Higgs pairs supported on irreducible and reduced spectral curves.*

In Chapter 6 we will discuss the relationship of these integrable systems with flows of KdV type. In the next section we will discuss the example of geodesic flow on the ellipsoid as a Hamiltonian flow of a symplectic leaf of one of these spaces. See [B1, Ma1] for more examples.

The Hitchin system has been useful in the study of the geometry of the moduli space of vector bundles. The main technique is to reduce questions about vector bundles to questions about spectral Jacobians. Hitchin used these ideas to compute the cohomology groups $H^i(\mathcal{U}, S^k T)$, $i = 0, 1$, of the symmetric products of the tangent bundle of the moduli space \mathcal{U} of rank 2 and odd degree stable vector bundles. In [BNR] these techniques provided the first mathematical proof that the dimensions of the space of sections of the generalized theta line bundle are

$$\begin{aligned} h^0(\mathcal{U}_\Sigma(n, n(g-1)), \Theta) &= 1, \\ h^0(\mathcal{S}\mathcal{U}_\Sigma(n), \Theta) &= n^g, \end{aligned}$$

where $\mathcal{S}\mathcal{U}_\Sigma(n)$ denotes the moduli space of vector bundles with trivial determinant line bundle. (This of course is now subsumed in the Verlinde Formula for sections of powers of theta bundles.) These ideas were proven useful in the proof of the existence of a projectively flat connection on the bundles of level k theta sections over the moduli space \mathcal{M}_g of curve of genus g [H3], an important fact in conformal field theory. Kouvidakis and Pantev applied these ideas to the study of automorphisms of the moduli space of vector bundles [Ko-P].

4.3 Polynomial Matrices

Theorem 4.8 has a concrete description when the base curve Σ is \mathbb{P}^1 . Let K be the line bundle $\mathcal{O}_{\mathbb{P}^1}(d)$. Consider the moduli space $\text{Higgs}_K := \text{Higgs}_K^{sm}(-r, r)$ of pairs (E, φ) consisting of a vector bundle E of rank r and degree $-r$ with a K -valued endomorphism $\varphi : E \rightarrow E \otimes K$ (we also follow the notation of definition 4.9 singling out a particular component). Choose a coordinate x on $\mathbb{P}^1 - \{\infty\}$. The space B_K of characteristic polynomials becomes

$$\{P(x, y) = y^r + b_1(x)y^{r-1} + \cdots + b_r(x) \mid b_i(x) \text{ is a polynomial in } x \text{ of degree } \leq i \cdot d\}.$$

The total space of the line bundle $\mathcal{O}_{\mathbb{P}^1}(d \cdot \infty)$ restricted to the affine line $\mathbb{P}^1 - \{\infty\}$ is isomorphic to the affine plane, and under this isomorphism $P(x, y)$ becomes the equation of the spectral curve as an affine plane curve.

Denote by $B^0 \subset B_K$ the subset of smooth spectral curves. Let $Q := Q_r(d)$ be the subset of Higgs_K parametrizing pairs (E, φ) with a smooth spectral curve and a vector bundle E isomorphic to $E_0 := \oplus^r \mathcal{O}_{\mathbb{P}^1}(-1)$. Q is a Zariski open (dense) subset of Higgs_K because:

- i) by definition 4.9 $\text{Higgs}_K^{sm}(r, -r)$ is irreducible,
- ii) E_0 is the unique semistable rank r vector bundle of degree $-r$ on \mathbb{P}^1 and semistability is an open condition.

The bundle $\text{End } E_0$ is the trivial Lie algebra bundle $\mathfrak{gl}_r(\mathbb{C}) \otimes \mathcal{O}_{\mathbb{P}^1}$. Hence, every point in Q is represented by an element $\varphi \in M_r(d) := H^0(\mathbb{P}^1, \mathfrak{gl}_r(\mathbb{C}) \otimes \mathcal{O}_{\mathbb{P}^1}(d \cdot \infty))$, i.e., by an $r \times r$ matrix φ with polynomial entries of degree $\leq d$. Denote the inverse image of B^0 in $M_r(d)$ by $M_r^0(d)$. The subset $Q \subset \text{Higgs}_K$ is simply the quotient of $M_r^0(d)$ by the conjugation action of $PGL_r(\mathbb{C})$.

$$\begin{array}{ccc}
M_r^0(d) & & \\
\downarrow & & \\
Q & \longrightarrow & \text{Higgs}_K \\
\downarrow & & \downarrow \\
B^0 & \longrightarrow & B_K
\end{array}$$

In this setting, Theorem 4.8 specializes to the following theorem of Beauville and Adams-Harnad-Hurtubise-Previato generalizing results of Mumford and Moser [Mum2] in rank 2:

Theorem 4.10 [B1, AHH]

1. *The quotient Q of the action of $PGL_r(\mathbb{C})$ by conjugation on $M_r^0(d)$ is a smooth variety.*
2. *The fiber of the characteristic polynomial maps $H : Q \rightarrow B^0$ over the polynomial of a spectral curve C is the complement $J_C^{g-1} - \Theta$ of the theta divisor in the Jacobian of line bundles on C of degree $g - 1$ ($g = \text{genus of } C$).*
3. *The choice of $d + 2$ points a_1, \dots, a_{d+2} on \mathbb{P}^1 determines a Poisson structure on Q . The characteristic polynomial map $H : Q \rightarrow B^0$ is an algebraically completely integrable Hamiltonian system with respect to each of these Poisson structures.*
4. *The symplectic leaves of Q are obtained by fixing the values (of the coefficients) of the characteristic polynomials at the points $\{a_i\}_{i=1}^{d+2}$.*

We note that in [B1] the Poisson structure on Q was obtained as the reduction of a Poisson structure on $M_r^0(d)$. The latter was the pullback of the Kostant-Kirillov Poisson structure via the embedding

$$M_r(d) \hookrightarrow \mathfrak{gl}_r(\mathbb{C})^{d+2}$$

by Lagrange interpolation at a_1, \dots, a_{d+2} . This embedding will be used in section 4.3.2 where geodesic flow on ellipsoids is revisited.

A choice of a divisor $D = a_1 + \dots + a_{d+2}$ of degree $d+2$ on \mathbb{P}^1 determines an isomorphism of $\mathcal{O}_{\mathbb{P}^1}(d \cdot \infty)$ with $\omega_{\mathbb{P}^1}(D)$. For example, if a_1, \dots, a_i are finite, $a_{i+1} = a_{i+2} = \dots = a_{d+2} = \infty$ then we send a polynomial $f(x)$ of degree $\leq d$ to the meromorphic 1-form

$$\frac{f(x)}{\prod_{j=1}^i (x - a_j)} dx.$$

When the $d + 2$ points are distinct, Lagrange interpolation translates to the embedding

$$\text{Res} : M_r(d) = H^0(\mathbb{P}^1, \mathfrak{gl}_r(\mathbb{C}) \otimes \omega_{\mathbb{P}^1}(D)) \hookrightarrow \mathfrak{gl}_r(\mathbb{C})^{d+2}$$

via the residues of meromorphic 1-form valued matrices at the points a_i (if a_i has multiplicity 2 or higher, we replace the i -th copy of $\mathfrak{gl}_r(\mathbb{C})$ by its tangent bundle or higher order infinitesimal germs at a_i of sections of the trivial bundle $\mathfrak{gl}_r(\mathbb{C}) \otimes \mathcal{O}_{\mathbb{P}^1}$).

4.3.1 Explicit Equations for Jacobians of Spectral Curves with a Cyclic Ramification Point

A further simplification occurs for matrices with a nilpotent leading coefficient (nilpotent at ∞). The projection $M_r^0(d) \rightarrow Q$ has a natural section over the image $N \subset Q$ of this locus. So N can be described concretely as a space of polynomials (rather than as a quotient of such a space).

As a consequence we obtain explicit equations in $M_r(d)$ for the complement $J_C - \Theta$ of the theta divisor in the Jacobian of every irreducible and reduced r -sheeted spectral curve over \mathbb{P}^1 which is totally ramified and smooth at ∞ (generalizing the equations for hyperelliptic curve (case $r = 2$) obtained in [Mum2]).

Lemma 4.11 *Let $A = A_d x^d + \dots + A_1 x + A_0$ be an $r \times r$ traceless matrix with polynomial entries of degree $\leq d$*

- i) *whose spectral curve in $\mathcal{O}_{\mathbb{P}^1}(d \cdot \infty)$ is irreducible and reduced and smooth over ∞ , and*
- ii) *whose leading coefficient A_d is a nilpotent (necessarily regular) matrix.*

Then there exists a unique element $g_0 \in PGL_r(\mathbb{C})$ conjugating A to a matrix $A' = x^d \cdot J + \sum_{i=0}^{d-1} A'_i x^i$ of the form:

$$A' = x^d \begin{pmatrix} 0 & & 0 & 0 \\ 1 & & 0 & 0 \\ 0 & \cdot & 0 & 0 \\ 0 & & 0 & 0 \\ 0 & & 0 & 0 \\ 0 & & 1 & 0 \end{pmatrix} + x^{d-1} \begin{pmatrix} \star & \dots & \star & \beta_r \\ \star & \dots & \star & 0 \\ \vdots & \vdots & \vdots & \vdots \\ \star & \dots & \star & 0 \end{pmatrix} + \sum_{i=0}^{d-2} x^i A'_i \quad (12)$$

where $(-1)^{r+1} \beta_r$ is the (leading) coefficient of x^{dr-1} in the determinant $b_r(x)$ of $A(x)$.

Remark: Notice that the coefficients $b_i(x)$ in the characteristic polynomial $P(x, y) = y^r + b_1(x)y^{r-1} + \dots + b_r(x)$ of $A(x)$ satisfy degree $b_i(x) \leq d \cdot i - 1$ since A is nilpotent at ∞ , and degree $b_r(x) = dr - 1$ since the spectral curve is smooth over ∞ . Thus $\beta_r \neq 0$.

Proof (of lemma 4.11): Let J be the nilpotent regular constant matrix appearing as the leading coefficient of $A'(x)$ in the normalized form (12). Let $\mathbb{C}[J]$ be the algebra of polynomials in J with constant coefficients. The proof relies on the elementary fact that \mathbb{C}^r , as a left $\mathbb{C}[J]$ -module, is free. Any vector with non zero first entry is a generator. A_d is conjugate to J . Thus we may assume that $A_d = J$ and it remains to show that there exists a unique element in the stabilizer of J in $PGL_r(\mathbb{C})$ conjugating $A(x)$ to the normal form (12).

Since $A_d = J$, the first entry in the right column R of A_{d-1} is β_r . Thus R is a generator of \mathbb{C}^r as a $\mathbb{C}[J]$ -module. Any element $g \in PGL_r(\mathbb{C})$ in the commutator subgroup of J is an invertible element in $\mathbb{C}[J]$ and can be written (up to scalar multiple) in the form $g = I + N$, N nilpotent. The right column of $gA_{d-1}g^{-1}$ is $R + NR$ and there exists a

unique nilpotent $N \in \mathbb{C}[J]$ such that $NR = \begin{pmatrix} \beta_r \\ 0 \\ \vdots \\ 0 \end{pmatrix} - R$. Thus g is unique up to a scalar

factor.

□

Denote the affine subvariety of $M_d^0(r)$ of matrices satisfying the $r^2 + r - 1$ equations (12) by \tilde{N} . The subvariety \tilde{N} is a section of the principal $PGL_r(\mathbb{C})$ bundle $M_r^0(d) \rightarrow Q$ over the locus N of conjugacy classes of polynomial matrices with a nilpotent leading coefficient. N is a Poisson subvariety of Q with respect to any Poisson structure on Q determined by a divisor D as in theorem 4.10, provided that D contains the point at infinity $\infty \in \mathbb{P}^1$.

Choose a characteristic polynomial $P(x, y) = y^r + b_1(x)y^{r-1} + \dots + b_r(x)$ in $B_{\mathbb{C}\mathbb{P}^1}(d)$ of a smooth spectral curve C with degree $b_i(x) \leq id - 1$, $b_r(x)$ of degree $rd - 1$ with leading coefficient $(-1)^{r+1}\beta_r$. Theorem 4.10 implies that the equations

a) $A_d = J$,

b) The r -th column of A_{d-1} is $\begin{pmatrix} \beta_r \\ 0 \\ \vdots \\ 0 \end{pmatrix}$,

c) $\text{char}(A(x)) = P(x, y)$

define a subvariety of $M_r(d)$ isomorphic to the complement $J_C^{g-1} - \Theta_C$ of the theta divisor in the Jacobian of C .

4.3.2 Geodesic Flow on Ellipsoids via 2×2 Polynomial Matrices

We use polynomial matrices to retrieve the Jacobi-Moser-Mumford system which arose in chapter 3 out of the geodesic flow on an ellipsoid. Our presentation follows [B1].

Consider a spectral polynomial $P(x, y)$ in $B_{\mathbb{C}\mathbb{P}^1}^0(d, \infty)$ of the form

(i) $P(x, y) = y^2 - f(x)$ where $f(x)$ is monic of degree $2d - 1$.

The corresponding spectral curve C is smooth, hyperelliptic of genus $g = d - 1$ and ramified over ∞ . Theorem 4.10 implies that the fiber

$$H^{-1}(P(x, y)) = \left\{ \begin{pmatrix} V & U \\ W & -V \end{pmatrix} \mid V^2 + UW = f(x) \right\} / PGL_2(\mathbb{C})$$

of the characteristic polynomial map is isomorphic to the complement $J_C^{g-1} - \Theta$ of the theta divisor.

Lemma 4.11 specializes in our case to the following statement (note that $\beta_r = 1$ since f is taken to be monic):

The $PGL_2(\mathbb{C})$ orbit of a matrix $\begin{pmatrix} V & U \\ W & -V \end{pmatrix}$ over $H^{-1}(P(x, y)) \cong J_C^{g-1} - \Theta$ contains a unique matrix satisfying

- (ii) W is monic of degree d ,
 U is monic of degree $d - 1$ and
 $\deg V \leq d - 2$.

In other words, condition (ii) and

- (iii) $V^2 + UW = f(x)$

are the equations of $J_C^{g-1} - \Theta$ as an affine subvariety of the subspace of traceless matrices in $M_2(d)$. In fact, condition (ii) defines a section $\varphi : N \rightarrow M_2(d)$ over the locus N in Q of conjugacy classes with characteristic polynomial satisfying condition (i).

Recall that the Jacobi-Moser-Mumford integrable system linearizing the geodesic flow of the ellipsoid $\sum_{i=1}^d a_i^{-1} x_i^2 = 1$ is supported on the tangent bundle TS of the sphere $S \subset \mathbb{R}^d$. Our discussion ended by describing the quotient of TS by the group $G \simeq (\mathbb{Z}/2\mathbb{Z})^d$ of involutions. We will describe in the next three steps an isomorphism of this quotient with a symplectic leaf X of Q .

Step I: (Identification of the symplectic leaf X). Assume that the points $a_1, \dots, a_d \in \mathbb{P}^1 - \{\infty\}$ are distinct and let $a_{d+1} = a_{d+2} = \infty$. Let $X \subset Q$ be the symplectic leaf over the subspace of characteristic polynomials $P(x, y) = y^2 - f(x)$ satisfying

$$f(a_i) = 0, \quad 1 \leq i \leq d, \quad \deg f = 2d - 1 \quad \text{and } f \text{ is monic.}$$

The spectral curves of matrices in the leaf X have genus $d - 1$, and are branched over the fixed $g + 2$ points a_1, \dots, a_d, ∞ and g varying points.

Step II: (Embedding of X in the product \mathcal{N}^d of the regular nilpotent orbit). The isomorphism $\mathcal{O}_{\mathbb{P}^1}(d \cdot \infty) \xrightarrow{\sim} \omega_{\mathbb{P}^1}(\sum_{i=1}^d a_i + 2 \cdot \infty)$ sending $F(x)$ to $\frac{F(x)dx}{\prod_{i=1}^d (x - a_i)}$ translates the matrix $\begin{pmatrix} V(x) & U(x) \\ W(x) & -V(x) \end{pmatrix}$ to a matrix φ of meromorphic 1-forms. The residues of φ satisfy:

$$R_\infty := \text{Res}_\infty(\varphi) = \begin{pmatrix} 0 & -1 \\ s & 0 \end{pmatrix} \text{ for some } s \in \mathbb{C} \quad (\text{condition (ii)}),$$

$$R_i := \text{Res}_{a_i}(\varphi) = \begin{pmatrix} V(a_i) & U(a_i) \\ W(a_i) & -V(a_i) \end{pmatrix} \frac{1}{\prod_{\substack{j=1 \\ j \neq i}}^d (a_i - a_j)}.$$

The residues at the finite points a_i can be calculated using Lagrange interpolation of polynomials of degree d at the $d + 1$ points a_1, \dots, a_d, ∞ given by the formula

$$F(x) = \sum_{i=1}^d F(a_i) \frac{\prod_{j=1, j \neq i}^d (x - a_j)}{\prod_{j=1, j \neq i}^d (a_i - a_j)} + F(\infty) \prod_{j=1}^d (x - a_j) \quad (13)$$

where $F(\infty)$ is the leading coefficient of $F(x)$.

The residues R_∞, R_i are nilpotent regular 2×2 matrices and the residue theorem implies that $R_\infty = -\sum_{i=1}^d R_i$. The residue map $Res : X \rightarrow \mathcal{N}^d$ defines a symplectic embedding $\varphi \mapsto (R_1, \dots, R_d)$ of the symplectic leaf X of Q in the Cartesian product of d copies of the regular nilpotent orbit \mathcal{N} in $\mathfrak{gl}_2(\mathbb{C})$.

Step III (The 2^d covering $TS \rightarrow X$). Endow \mathbb{C}^2 with the symplectic structure $2dx \wedge dy$. The map $\mathbb{C}^2 - \{(0,0)\} \rightarrow \mathcal{N}$ sending (x, y) to $\begin{pmatrix} xy & -x^2 \\ y^2 & -xy \end{pmatrix}$ is a symplectic $SL_2(\mathbb{C})$ -equivariant double cover of the regular nilpotent orbit \mathcal{N} (where $SL_2(\mathbb{C}) \cong Sp_2(\mathbb{C}, 2dx \wedge dy)$ acts on \mathbb{C}^2 via the standard representation). We obtain a 2^d -covering $\tau : (\mathbb{C}^2 - \{(0,0)\})^d \rightarrow \mathcal{N}^d$. The residue theorem translates to the fact that the image $Res(X) \subset \mathcal{N}^d$ is covered by

$$\left\{ (\bar{x}, \bar{y}) = ((x_1, y_1), \dots, (x_d, y_d)) \mid \sum x_i y_i = 0 \text{ and } \sum_{i=1}^d x_i^2 = 1 \right\}.$$

This is exactly the tangent bundle $TS \subset (\mathbb{C}^2)^d$ of the sphere $S \subset \mathbb{C}^d$.

5 Poisson structure via levels

We construct a Poisson structure on the moduli space of meromorphic Higgs pairs in two steps (following [Ma1]):

- First we realize a dense open subset of moduli as the orbit space of a Poisson action of a group on the cotangent bundle of the moduli space of vector bundles with level structures (sections 5.1, 5.2 and 5.3).
- Next we exhibit a 2-vector on the smooth locus of moduli, using a cohomological construction (section 5.4). On the above dense open set this agrees with the Poisson structure, so it is a Poisson structure everywhere.

We summarize the construction in section 5.5 in a diagram whose rotational symmetry relates dual pairs of realizations.

5.1 Level structures

Fix a curve Σ , an effective divisor $D = \sum p_i$ in Σ , and a rank r vector bundle E on Σ . A level D structure on E is an \mathcal{O}_D -isomorphism $\eta : E \otimes \mathcal{O}_D \xrightarrow{\sim} \mathcal{O}_D^{\oplus r}$. Seshadri [Se] constructs a smooth, quasi-projective moduli space $\mathcal{U}_\Sigma(r, d, D)$ parametrizing stable pairs (E, η) . Here stability means that for any subbundle $F \subset E$,

$$\frac{\deg F - \deg D}{\text{rank} F} < \frac{\deg E - \deg D}{\text{rank} E}.$$

The level- D group is the projectivized group of \mathcal{O}_D -algebra automorphisms,

$$G_D := \mathbb{P}\text{Aut}_{\mathcal{O}_D}(\mathcal{O}_D^{\oplus r}).$$

(i.e. the automorphism group modulo complex scalars \mathbb{C}^* .) It acts on $\mathcal{U}_\Sigma(r, d, D)$: an element $g \in G_D$ sends

$$[(E, \eta)] \mapsto [(E, \bar{g} \circ \eta)],$$

where $\bar{g} \in \text{Aut}_{\mathcal{O}_D}(\mathcal{O}_D^{\oplus r})$ lifts g , and $[\cdot]$ denotes the isomorphism class of a pair. The open set $\mathcal{U}_\Sigma^{\circ}(r, d, D)$, parametrizing pairs (E, η) where E itself is stable, is a principal G_D -bundle over $\mathcal{U}_\Sigma^s(r, d)$. The Lie algebra \mathfrak{g}_D of G_D is given by $\mathfrak{gl}_r(\mathcal{O}_D)/\text{scalars}$.

5.2 The cotangent bundle

We compute deformations of a pair

$$(E, \eta) \in \mathcal{U}_D := \mathcal{U}_\Sigma(r, d, D)$$

as we did for the single vector bundle

$$E \in \mathcal{U} := \mathcal{U}_\Sigma(r, d).$$

Namely, E is given (in terms of an open cover of Σ) by a 1-cocycle with values in the sheaf of groups $GL_r(\mathcal{O}_\Sigma)$. Differentiating this cocycle with respect to parameters gives

a 1-cocycle with values in the corresponding sheaf of Lie algebras, so we obtain the identification

$$T_E \mathcal{U} \approx H^1(\text{End}E).$$

Similarly, the pair (E, η) is given by a 1-cocycle with values in the subsheaf

$$GL_{r,D}(\mathcal{O}_\Sigma) := \{f \in GL_r(\mathcal{O}_\Sigma) \mid f - 1 \in \mathfrak{gl}_r(\mathcal{I}_D)\}.$$

Differentiating, we find the natural isomorphism

$$T_{(E,\eta)} \mathcal{U}_D \approx H^1(\mathcal{I}_D \otimes \text{End}E),$$

so by Serre duality,

$$T_{(E,\eta)}^* \mathcal{U}_D \approx H^0(\text{End}E \otimes \omega(D)).$$

(we identify $\text{End}E$ with its dual via the trace.) We will denote a point of this cotangent bundle by a triple (E, φ, η) , where $(E, \eta) \in \mathcal{U}_D$ and φ is a D -Higgs field,

$$\varphi : E \rightarrow E \otimes \omega(D).$$

5.3 The Poisson structure

The action of the level group G_D on \mathcal{U}_D lifts naturally to an action of G_D on $T^* \mathcal{U}_D$. Explicitly, an element $g \in G_D$ with lift $\bar{g} \in GL_r(\mathcal{O}_D)$ sends

$$(E, \varphi, \eta) \mapsto (E, \varphi, \bar{g} \circ \eta).$$

The lifted action has the following properties:

- (1) It is Poisson with respect to the standard symplectic structure on $T^* \mathcal{U}_D$ (holds for any lifted action, see example 2.2).
- (2) The moment map

$$\mu : T^* \mathcal{U}_D \rightarrow \mathfrak{g}_D^*$$

is given by

$$\mu(E, \varphi, \eta) : A \mapsto \text{Res Trace}(A \cdot \varphi^\eta), \tag{14}$$

where

$$\begin{aligned} A &\in \mathfrak{g}_D = (\mathfrak{gl}_r(\mathcal{O}_D)) / (\text{scalars}) \approx (\mathfrak{gl}_r(\mathcal{O}_D))_{\text{traceless}}, \\ \varphi^\eta &:= \eta \circ \varphi \circ \eta^{-1} \in H^0(\mathfrak{gl}_r(\omega(D) \otimes \mathcal{O}_D)), \end{aligned}$$

and the residue map

$$\text{Res} : H^0(\omega(D) \otimes \mathcal{O}_D) \rightarrow H^1(\omega) \approx \mathbb{C}$$

is the coboundary for the restriction sequence

$$0 \rightarrow \omega \rightarrow \omega(D) \rightarrow \omega(D) \otimes \mathcal{O}_D \rightarrow 0$$

(cf. [Ma1] Proposition 6.12).

- (3) G_D acts freely on the open subset $(T^* \mathcal{U}_D)^\circ$ parametrizing triples (E, φ, η) where (E, η) is stable and (E, φ) is a stable Higgs bundle, since such bundles are simple. This makes $(T^* \mathcal{U}_D)^\circ$ into a principal G_D -bundle over an open subset Higgs_D° of Higgs_D^s .

We conclude that the symplectic structure on $(T^* \mathcal{U}_D)^\circ$ induces a Poisson structure on Higgs_D° . The symplectic leaves will then be the inverse images under μ of coadjoint orbits in \mathfrak{g}_D^* .

5.4 Linearization

The main remaining task is to find a two-vector on the non-singular locus Higgs_D^{ns} whose restriction to Higgs_D° is the above Poisson structure. This two vector is then automatically Poisson. The algebraic complete integrability of the component Higgs_D^{sm} (see definition 4.9) would then follow: The spectral curve C_b , for generic $b \in B_D$, is non-singular, so its Jacobian $\mathcal{J}(C)$ is contained in Higgs_D^{sm} . Thus any Hamiltonian vector field must be constant on the generic fiber $\mathcal{J}(C)$, hence on all fibers.

A natural two-vector defined over all of Higgs_D^{ns} can be given in several ways. One [Ma1] is to identify the tangent spaces to Higgs_D (and related spaces) at their smooth points as hypercohomologies, \mathbb{H}^1 , of appropriate complexes:

<u>space</u>	<u>at</u>	<u>complex</u>
\mathcal{U}	E	$\text{End}E$
\mathcal{U}_D	(E, η)	$\text{End}E(-D)$
Higgs_D	(E, φ)	$\overline{\mathcal{K}} = [\text{End}E \xrightarrow{ad\varphi} \text{End}E \otimes \omega(D)]$
$T^*\mathcal{U}_D$	(E, φ, η)	$\mathcal{K} := [\text{End}E(-D) \xrightarrow{ad\varphi \otimes i} \text{End}E \otimes \omega(D)]$

where $i : \mathcal{O}(-D) \hookrightarrow \mathcal{O}$ is the natural inclusion. These identifications are natural, and differentials of maps between these spaces are realized by maps of complexes. For example, the fibration $T^*\mathcal{U}_D \rightarrow \mathcal{U}_D$, with fiber $T_{(E,\eta)}^*\mathcal{U}_D$, gives the sequence

$$\begin{array}{ccccccc}
 0 & \rightarrow & T_{(E,\eta)}^*\mathcal{U}_D & \rightarrow & T_{(E,\varphi,\eta)}(T^*\mathcal{U}_D) & \rightarrow & T_{(E,\eta)}\mathcal{U}_D & \rightarrow & 0 \\
 & & \parallel & & \parallel & & \parallel & & \\
 0 & \rightarrow & H^0(\text{End}E \otimes \omega(D)) & \rightarrow & \mathbb{H}^1(\mathcal{K}) & \rightarrow & H^1(\text{End}E(-D)) & \rightarrow & 0
 \end{array}$$

derived from the short exact sequence of complexes,

$$0 \rightarrow \text{End}E \otimes \omega(D)[-1] \rightarrow K \rightarrow \text{End}E(-D) \rightarrow 0,$$

while the (rational) map $T^*\mathcal{U}_D \rightarrow \text{Higgs}_D$ gives

$$\begin{array}{ccccccc}
 0 & \rightarrow & \mathfrak{g}_D & \rightarrow & T_{(E,\varphi,\eta)}(T^*\mathcal{U}_D) & \rightarrow & T_{(E,\varphi)}\text{Higgs}_D & \rightarrow & 0 \\
 & & \parallel & & \parallel & & \parallel & & \\
 0 & \rightarrow & \frac{H^0(\text{End}E \otimes \mathcal{O}_D)}{H^0(\text{End}E)} & \rightarrow & \mathbb{H}^1(\mathcal{K}) & \rightarrow & \mathbb{H}^1(\overline{\mathcal{K}}) & \rightarrow & 0
 \end{array}$$

which derives from:

$$0 \rightarrow \mathcal{K} \rightarrow \overline{\mathcal{K}} \rightarrow \text{End}E \otimes \mathcal{O}_D \rightarrow 0.$$

The dual of a complex $\mathcal{L} : A \rightarrow B$ of vector bundles is the complex

$$\mathcal{L}^\vee : B^* \otimes \omega \rightarrow A^* \otimes \omega.$$

Grothendieck duality in this case gives a natural isomorphism

$$\mathbb{H}^1(\mathcal{L}) \approx \mathbb{H}^1(\mathcal{L}^\vee)^*.$$

We note that \mathcal{K} is self-dual, in the sense that there is a natural isomorphism of complexes, $J : \mathcal{K}^\vee \xrightarrow{\sim} \mathcal{K}$. For $\overline{\mathcal{K}}$ we obtain a natural isomorphism of complexes, $\overline{\mathcal{K}}^\vee \xrightarrow{\sim} \overline{\mathcal{K}} \otimes \mathcal{O}(-D)$, hence (composing with i) a morphism

$$I : \overline{\mathcal{K}}^\vee \rightarrow \overline{\mathcal{K}}.$$

Combining with duality, we get maps

$$\mathbb{H}^1(\mathcal{K})^* \approx \mathbb{H}^1(\mathcal{K}^\vee) \xrightarrow{J} \mathbb{H}^1(\mathcal{K})$$

and

$$\mathbb{H}^1(\overline{\mathcal{K}})^* \approx \mathbb{H}^1(\overline{\mathcal{K}}^\vee) \xrightarrow{I} \mathbb{H}^1(\overline{\mathcal{K}}).$$

These give elements of $\otimes^2 \mathbb{H}^1(\mathcal{K})$ and $\otimes^2 \mathbb{H}^1(\overline{\mathcal{K}})$. Both are skew symmetric (since ad_φ , and hence I, J , are), so we obtain global two-vectors on $T^*\mathcal{U}_D^{ns}$ and Higgs_D^{ns} . At stable points these agree with (the dual of) the symplectic form and its reduction modulo G_D , which is what we need.

Another way to find the two-vector on Higgs_D is based on the interpretation of Higgs_D as a moduli space of sheaves on the total space S of $\omega(D)$. At such a simple sheaf \mathcal{E} , with support on some spectral curve C , Mukai [Mu1] identifies the tangent space to moduli with

$$\text{Ext}_{\mathcal{O}_S}^1(\mathcal{E}, \mathcal{E}),$$

and notes that any two-form $\sigma \in H^0(\omega_S)$ determines an alternating bilinear map:

$$\text{Ext}_{\mathcal{O}_S}^1(\mathcal{E}, \mathcal{E}) \times \text{Ext}_{\mathcal{O}_S}^1(\mathcal{E}, \mathcal{E}) \rightarrow \text{Ext}_{\mathcal{O}_S}^2(\mathcal{E}, \mathcal{E}) \xrightarrow{\text{tr}} H^2(\mathcal{O}_S) \xrightarrow{\sigma} H^2(\omega_S) \approx \mathbb{C},$$

hence a two-form on moduli. Mukai uses this argument to produce symplectic structures on the moduli spaces of sheaves on $K3$ and abelian surfaces. The same argument works, of course, for sheaves on $T^*\Sigma$; this reconstructs the symplectic form on Hitchin's system.

Our surface S (the total space of $\omega(D)$) is related to $T^*\Sigma$ by a birational morphism $\alpha : T^*\Sigma \rightarrow S$. The symplectic form σ does not descend to S , but its inverse σ^{-1} does give a two vector on S which is non-degenerate away from D and is closed there (since it is locally equivalent to the Poisson structure on $T^*\Sigma$).

Tyurin notes [Ty1] that a variant of this argument produces a two-vector on moduli from a two-vector on S . Now the birational morphism $T^*\Sigma \rightarrow S$ takes the Poisson structure on $T^*\Sigma$ to one on S , so the Mukai-Tyurin argument gives the desired two-vector on Higgs_D . In chapter 8 this approach is generalized to higher dimensional varieties.

A third argument for the linearization is given by Bottacin [Bn]. He produces an explicit two-vector at stable points using a deformation argument as above, and then makes direct, local computations to check closedness of the Poisson structures and linearity of the flows.

5.5 Hamiltonians and flows in $T^*\mathcal{U}_D$

We saw that the level group G_D acts on $T^*\mathcal{U}_D$, inducing the Poisson structure on the quotient Higgs_D , and that the moment map is

$$\begin{aligned} \mu : T^*\mathcal{U}_D &\rightarrow \mathfrak{g}_D^* \\ \mu(E, \varphi, \eta)(A) &:= \text{Res Trace}(A \cdot \varphi^\eta). \end{aligned}$$

The characteristic polynomial map $\tilde{h} : T^*\mathcal{U}_D \rightarrow B_D$ is a composition of the Poisson map $T^*\mathcal{U}_D \rightarrow \text{Higgs}_D$ with the Hamiltonian map $h : \text{Higgs}_D \rightarrow B_D$. Hence \tilde{h} is also Hamiltonian. Clearly, \tilde{h} is G_D -invariant.

The composition $T^*\mathcal{U}_D \xrightarrow{\mu} \mathfrak{g}_D^* \rightarrow \mathfrak{g}_D^*/G_D$ is a G_D -invariant Hamiltonian morphism and hence factors through Higgs_D . It follows that it factors also through B_D since $h : \text{Higgs}_D \rightarrow B_D$ is a Lagrangian fibration whose generic fiber is connected (see remark 2.15). The conditions of example 2.16 in section 2.4 are satisfied and we get a diagram with a 180° rotational symmetry in which opposite spaces are dual pairs of realizations. The realization dual to $T^*\mathcal{U}_D \rightarrow \mathfrak{g}_D^*/G_D$ is the rational morphism $T^*\mathcal{U}_D \rightarrow G\text{Higgs}_D := \text{Higgs}_D \times_{(\mathfrak{g}_D^*/G_D)} \mathfrak{g}_D^*$ to the fiber product. The one dual to $\tilde{h} : T^*\mathcal{U}_D \rightarrow B_D$ is the morphism $T^*\mathcal{U}_D \rightarrow GB_D := B_D \times_{(\mathfrak{g}_D^*/G_D)} \mathfrak{g}_D^*$ to the fiber product. We write down the spaces and typical elements in them:

$$\begin{array}{ccc}
 T^*\mathcal{U}_D & & (E, \varphi, \eta) \\
 \downarrow & & \downarrow \\
 GHiggs_D & & (E, \varphi, \varphi^\eta) \\
 \swarrow \quad \searrow & & \swarrow \quad \searrow \\
 Higgs_D & & (\text{char}\varphi, \varphi^\eta) \\
 \swarrow \quad \searrow & & \swarrow \quad \searrow \\
 B_D & & \text{char}\varphi \quad \varphi^\eta \\
 \swarrow \quad \searrow & & \swarrow \quad \searrow \\
 \mathfrak{g}_D^*/G_D & & \text{char}\varphi^\eta \\
 \downarrow & & \downarrow \\
 (0) & & (0)
 \end{array} \tag{15}$$

6 Spectral flows and KP

Our aim in this section is to relate the general spectral system which we have been considering to the KP and multi-component KP hierarchies. We start by reviewing these hierarchies and their traditional relationship to curves and bundles via the Krichever map. We then reinterpret these flows as coming from Hamiltonians on the limit T^*U_∞ of our previous symplectic spaces. We show that Higgs_∞ can be partitioned into a finite number of loci, each of which maps naturally to one of the $mcKP$ -spaces in a way which intertwines isospectral flows with KP flows. As an example we consider the Elliptic solitons studied by Treibich and Verdier.

6.1 The hierarchies

KP

Following the modern custom (initiated by Sato, explained by Segal-Wilson [SW], and presented elegantly in [AdC, Mul, LM1] and elsewhere), we think of the KP hierarchy as given by the action of an infinite-dimensional group on an infinite-dimensional Grassmannian: set

$$\begin{aligned} K &:= \mathbb{C}((z)) = \text{field of formal Laurent series in a variable } z \\ Gr &:= \{\text{subspaces } W \subset K \mid \text{projection } W \rightarrow K/\mathbb{C}[[z]]z \text{ is Fredholm}\} \\ &= \{\text{subspaces "comparable to } \mathbb{C}[z^{-1}]\text{"}\}. \end{aligned}$$

This can be given an algebraic structure which allows us to talk about vector fields on Gr , finite-dimensional algebraic subvarieties, etc. Every $a \in K$ determines a vector field KP_a on Gr , whose value at $W \in Gr$ is the map

$$W \hookrightarrow K \xrightarrow{a} K \rightarrow K/W,$$

considered as an element of

$$\text{Hom}(W, K/W) \approx T_W Gr.$$

The (double) KP hierarchy on Gr is just this collection of commuting vector fields. For $a \in \mathbb{C}[[z]]$, this vector field comes from the action on Gr of the one-parameter subgroup $\exp(ta)$ in $\mathbb{C}[[z]]^*$, which we consider trivial. The KP hierarchy itself thus consists of the vector fields KP_a , for $a \in \mathbb{C}[z^{-1}]z^{-1}$, on the quotient $Gr/(\mathbb{C}[[z]]^*)$. This quotient is well-behaved: the action of \mathbb{C}^* is trivial, and the unipotent part $1 + z\mathbb{C}[[z]]$ acts freely and with transversal slices. One restricts attention to the open subset of this quotient ("the big cell") parametrizing W of fixed index (the index of W is the index of the Fredholm projection) and satisfying a general position condition with respect to the standard subspace $W_0 := \mathbb{C}[z^{-1}]$. Sato's construction identifies this subset with the space Ψ of pseudo differential operators of the form

$$\mathcal{L} = D + \sum_{i=1}^{\infty} u_i D^{-i}$$

where

$$u_i = u_i(t_1, t_2, \dots)$$

and $D = \partial/\partial t_1$. The resulting flows on Ψ have the familiar Lax form:

$$\frac{\partial \mathcal{L}}{\partial t_i} = [(\mathcal{L}^i)_+, \mathcal{L}],$$

where $(\mathcal{L}^i)_+$ is the differential operator part of \mathcal{L}^i .

multi component KP

The k^{th} multi-component KP hierarchy ($mcKP$) is obtained by considering instead the Grassmannian Gr_k of subspaces of $K^{\oplus k}$ comparable to $(\mathbb{C}[z^{-1}])^{\oplus k}$. The entire “loop algebra” $gl(k, K)$ acts here, but to obtain commuting flows we need to restrict to a commutative subalgebra. For the k -th multi-component KP we take the simplest choice, of diagonal matrices, i.e. we consider the action of $(\mathbb{C}[z^{-1}]z^{-1})^{\oplus k}$ on the quotient $Gr_k/(\mathbb{C}[[z]])^k$. There is a big cell $\Psi_k \subset Gr_k/(\mathbb{C}[[z]])^k$, consisting as before of subspaces in general position with respect to a reference subspace W_0 , on which the flow is given by a Lax equation (for vector-valued operators).

Heisenberg flows

More generally, for a partition

$$\underline{n} = (n_1, \dots, n_k)$$

of the positive integer n , we can consider, following [AB], the maximal torus $Heis_{\underline{n}}$ of type \underline{n} in $GL(n, K)$, as well as $heis_{\underline{n}}$, the corresponding Lie subalgebra in $gl(n, K)$. These consist of matrices in block-diagonal form, where the i^{th} block is a formal power series in the $n_i \times n_i$ matrix

$$P_{n_i} := \begin{pmatrix} 0 & & 0 & z \\ 1 & & 0 & 0 \\ 0 & \cdot & 0 & 0 \\ 0 & & 0 & 0 \\ 0 & & \cdot & 0 & 0 \\ 0 & & & 1 & 0 \end{pmatrix} \quad (16)$$

We recall that this matrix arises naturally when we consider a vector bundle which is the direct image of a line bundle, near a point where n_i sheets come together: in terms of a natural local basis of the vector bundle, it expresses multiplication by a coordinate upstairs (see (11)). The $\underline{n}^{\text{th}}$ $mcKP$ (or “Heisenberg flows” of type \underline{n}) lives on the quotient of Gr_n by the non-negative powers of the P_{n_i} , and a basis for the surviving flows is indexed by k -tuples (d_1, \dots, d_k) , $d_i > 0$. Again, this can all be realized by Lax equations on an appropriate space $\Psi_{\underline{n}}$ of pseudo differential operators. When $\underline{n} = (1, \dots, 1)$ we recover the n^{th} $mcKP$. When $\underline{n} = (n)$, the flows are pulled back from the standard KP flows on Gr , via the mixing map

$$m_n : Gr_n \rightarrow Gr$$

sending

$$\widetilde{W} \subset \mathbb{C}((\tilde{z}))^{\oplus n}$$

to

$$W := \left\{ \sum_{i=0}^{n-1} a_i(z^n) z^i \mid (a_0(\tilde{z}), \dots, a_{n-1}(\tilde{z})) \in \widetilde{W} \right\} \subset \mathbb{C}((z)).$$

An arbitrary k -part partition \underline{n} of n determines a map

$$m_{\underline{n}} : Gr_n \rightarrow Gr_k,$$

and the \underline{n}^{th} Heisenberg flows are pullbacks of the k^{th} $mcKP$. The natural big cell in this situation is determined by the cartesian diagram:

$$\begin{array}{ccccc} Gr_n & \longrightarrow & Gr_n/Heis_{\underline{n}}^+ & \longleftrightarrow & \Psi_{\underline{n}} \\ \downarrow m_{\underline{n}} & & \downarrow & & \downarrow \\ Gr_k & \longrightarrow & Gr_k/(C[[z]])^k & \longleftrightarrow & \Psi_k \end{array}$$

6.2 Krichever maps

The data

A basic Krichever datum (for the KP hierarchy) consists of a quintuple

$$(C, p, z, L, \eta)$$

where:

- C is a (compact, non-singular) algebraic curve
- $p \in C$
- z is a local (analytic or formal) coordinate at p
- $L \in PicC$
- $\eta : L \otimes \hat{\mathcal{O}}_p \xrightarrow{\cong} \hat{\mathcal{O}}_p \approx \mathbb{C}[[z]]$ is a (formal) trivialization of L near p .

If we fix C, p and z , we think of the Krichever datum as giving a point of

$$\mathcal{U}_C(1, \infty p) := \varinjlim_{\ell} \mathcal{U}_C(1, \ell p).$$

The Krichever map

$$\{\text{Krichever data}\} \rightarrow Gr$$

sends the above datum to the subspace

$$W := \eta(H^0(C, L(\infty p))) = \bigcup_k \eta(H^0(C, L(kp))) \subset \mathbb{C}((z)).$$

This subspace is comparable to $\mathbb{C}[z^{-1}]$, since it follows from Riemann-Roch that the dimension of $H^0(L(kp))$ differs from k by a bounded (and eventually constant) quantity.

The flows

Let's work with a coordinate z which is analytic, i.e. it actually converges on some disc. A line bundle L on C can be trivialized (analytically) on the Stein manifold $C \setminus p$. We can think of (L, η) as being obtained from $\mathcal{O}_{C \setminus p}$ by glueing it to $\hat{\mathcal{O}}_p$ via a 1-cocycle, or transition function, which should consist of an invertible function g on a punctured neighborhood of p in C . Conversely, we claim there is a map:

$$\exp : K \longrightarrow \mathcal{U}_C(1, \infty p),$$

$$f \longmapsto (L, \eta).$$

For $f \in \mathbb{C}(z)$, this is defined by the above analytic gluing, using $g := \exp f$, which is indeed analytic on a punctured neighborhood. For $f \in \mathbb{C}[[z]] \approx \hat{\mathcal{O}}_p$, on the other hand, we take $(L, \eta) := (\mathcal{O}, \exp f)$. These two versions agree on the intersection, $f \in \mathbb{C}[z]_{(0)}$, so the map is uniquely defined as claimed. (The bundles we get this way all have degree 0, but we can also obtain maps

$$\exp_{g_0} : K \longrightarrow \mathcal{U}_C(1, d, \infty p)$$

to the moduli space of level- ∞p line bundles of degree d , simply by fixing a meromorphic function g_0 on a neighborhood of p which has order d at p , and replacing the previous g by $g_0 \exp f$. We will continue to suppress the degree d in our notation.)

Any $a \in K$ gives an additive flow on K , which at $f \in K$ is

$$t \longmapsto f + ta.$$

Under the composed map

$$K \xrightarrow{\exp_{g_0}} \mathcal{U}_C(1, \infty p) \xrightarrow{\text{Krichever}} Gr,$$

this is mapped to the double KP flow KP_a on Gr . For $a \in \mathbb{C}[[z]]$ this flow does not affect the isomorphism class of L , and simply multiplies η by $\exp(ta)$. On the other hand, the i^{th} KP flow, given by $a = z^{-i}$, changes both L and η if $i > 0$. The projection to $\text{Pic } C$ is a linear flow, whose direction is the i^{th} derivative at p , with respect to the coordinate z , of the Abel-Jacobi map $C \rightarrow \text{Pic } C$. Dividing out the trivial flows corresponds to suppressing η , so we obtain, for each C, p, z and degree $d \in \mathbb{Z}$, a finite-dimensional orbit of the KP flows in $Gr/\mathbb{C}[[z]]^*$, isomorphic to $\text{Pic}^d C$.

Multi-Krichever data

Several natural generalizations of the Krichever map to the multi-component KP can be found in [AB, LM1] and elsewhere. Here are some of the possibilities. We can consider “multi-Krichever” data

$$(C, D, z_i, L, \eta)$$

involving a curve C with a divisor D consisting of k distinct points $p_i (1 \leq i \leq k)$, a coordinate z_i at each p_i , a line bundle L , and a formal trivialization η_i at each p_i . Fixing C, p_i and z_i , we have a multi-Krichever map

$$\{\text{multi-Krichever data}\} \approx \mathcal{U}_C(1, \infty D) \longrightarrow Gr_k$$

sending

$$(L, \eta_i) \mapsto W := (\eta_1, \dots, \eta_k)(H^0(C, L(\infty D))) \subset \mathbb{C}((z))^{\oplus k}.$$

The k -component KP flow on the right hand side given by $a = (a_1, \dots, a_k) \in K^k$ restricts to the flow on the multi-Krichever data which multiplies the transition function at p_i (for an analytic trivialization of L on $C \setminus D$) by $\exp a_i$.

We can also consider “vector-Krichever” data (C, p, z, E, η) where the line bundle L is replaced by a rank n vector bundle E , and

$$\eta : E \otimes \hat{\mathcal{O}}_p \xrightarrow{\cong} (\hat{\mathcal{O}}_p)^n \approx (\mathbb{C}[[z]])^n$$

is now a (formal) trivialization of E near p . Not too surprisingly, the vector-Krichever map

$$\{\text{vector-Krichever data}\} \rightarrow Gr_n$$

sends the above datum to the subspace

$$W := \eta(H^0(C, E(\infty p))) \subset (\mathbb{C}((z)))^n.$$

In the next subsection we will see that the interesting interaction of these two types of higher Krichever maps occurs not by extending further (to objects such as (C, D, z_i, E, η_i)), but by restricting to those vector data on one curve which match some multi-data on another.

KdV-type subhierarchies

Among the Krichever data one can restrict attention to those quintuples where z^{-n} (for some fixed n) happens to extend to a regular function on $C \setminus p$, i.e. gives a morphism

$$f = z^{-n} : C \rightarrow \mathbb{P}^1$$

of degree n , such that the fiber $f^{-1}(\infty)$ is $n \cdot p_0$. The Krichever map sends such data to the n^{th} KdV hierarchy, the distinguished subvariety of Gr (invariant under the (double) KP flows) given by

$$\text{KdV}_n := \{W \in Gr \mid z^{-n}W \subset W\}.$$

The corresponding subspace of Ψ is

$$\{\mathcal{L} \mid \mathcal{L}^n = \mathcal{L}_+^n \text{ is a differential operator}\}.$$

Fixing a partition $\underline{n} = (n_1, \dots, n_k)$ of n , we can similarly consider the covering data of type \underline{n} , consisting of the multi-Krichever data (C, p_i, z_i, L, η_i) plus a map $f : C \rightarrow \Sigma$ of degree n to a curve Σ with local coordinate z at a point $\infty \in \Sigma$, such that

$$f^{-1}(p) = \Sigma n_k p_i, \quad f^{-1}(z) = z_i^{n_i} \text{ at } p_i.$$

Such a covering datum clearly gives a multi-Krichever datum on C , but it also determines a vector-Krichever datum (E, η) on Σ : The standard m -sheeted branched cover

$$\begin{array}{ccc} f_m : \mathbb{C} & \rightarrow & \mathbb{C} \\ \tilde{z} & \mapsto & z = \tilde{z}^m \end{array}$$

of the z -line determines an isomorphism

$$s_m : (f_m)_* \mathcal{O} \xrightarrow{\cong} \mathcal{O}^{\oplus m}$$

given by

$$\sum_{i=0}^{m-1} a_i(\tilde{z}^m) \tilde{z}^i \mapsto (a_0(z), \dots, a_{m-1}(z)).$$

To the covering datum above we can then associate the rank- n vector bundle $E := f_* L$ on Σ , together with the trivialization at p obtained by composing

$$\oplus_i f_{*, p_i}(\eta_i) : (f_* L)_p \xrightarrow{\cong} \oplus_i f_{*, p_i}(\mathcal{O}_{p_i})$$

with the isomorphisms

$$f_{*,p_i}(\mathcal{O}_{p_i}) \xrightarrow{\sim} \mathcal{O}_p^{\oplus n_i}$$

which are conjugates of the standard isomorphisms s_{n_i} by the chosen local coordinates z, z_i . Finally, we note that there are obvious geometric flows on these covering data: L and η_i flow as before, while everything else stays put. The compatibility of the two types of higher Krichever data is expressed by the commutativity of the diagram:

$$\begin{array}{ccc}
\{\underline{n} - \text{covering data}\} & \approx & \{f : C \rightarrow \Sigma; p_i, z_i, L, \eta_i; z \mid \dots\} \\
\downarrow & & \downarrow \\
\{\text{vector Krichever data on } \Sigma\} & & \{\text{multi Krichever data on } C\} \\
\parallel & & \parallel \\
\cup_{\Sigma, p, z} \mathcal{U}_{\Sigma}(n, \infty p) & & \cup_{C, D, z_i} \mathcal{U}_C(1, \infty D) \\
\downarrow & & \downarrow \\
Gr_n & \xrightarrow{m_{\underline{n}}} & Gr_k \\
\downarrow & \longrightarrow & \downarrow \\
Gr_n / Heis_{\underline{n}}^+ & & Gr_k / (\mathbb{C}[[z]]^{\times})^k.
\end{array} \tag{17}$$

The mcKP flows on the bottom right pull back to the Heisenberg flows on the bottom left, and to the geometric flows on the \underline{n} -covering data.

6.3 Compatibility of hierarchies

Fix a smooth algebraic curve Σ of arbitrary genus and a point P in it. The moduli space $\text{Higgs}_D := \text{Higgs}_{\Sigma}^{sm}(n, d, \omega(D))$ (see definition 4.9) can be partitioned into type loci. We consider the Zariski dense subset consisting of the union of finitely many type loci $\text{Higgs}_D^{\underline{n}}$ indexed by partitions \underline{n} of n . A Higgs pair in $\text{Higgs}_D^{\underline{n}}$ has a spectral curve $C \rightarrow \Sigma$ whose ramification type over $P \in \Sigma$ is $\underline{n} = (n_1, \dots, n_k)$.

Fix a formal local parameter z on the base curve Σ at P . A Higgs pair in $\text{Higgs}_D^{\underline{n}}$ (or rather its spectral pair (C, L) , see proposition 4.5) can be completed to an $\text{Heis}_{\underline{n}}^+$ -orbit of an \underline{n} -covering data $(C, P_i, z_i, L, \eta_i) \rightarrow (\Sigma, P, z, E, \eta)$ in finitely many ways. These extra choices form a natural finite Galois cover $\widetilde{\text{Higgs}}_D^{\underline{n}}$ of each type locus $\text{Higgs}_D^{\underline{n}}$. We obtain Krichever maps (see diagram 17) from the Galois cover $\widetilde{\text{Higgs}}_D^{\underline{n}}$ to the quotients $Gr_n / \text{Heis}_{\underline{n}}^+$ and $Gr_k / (\mathbb{C}[[z]]^{\times})^k$. Both the mcKP and Heisenberg flows pull back to the same geometric flow on the Galois cover. It is natural to ask:

Question 6.1 (The compatibility question) *Is the Heisenberg flow Poisson with respect to the natural Poisson structure on Higgs_D ?*

The Compatibility Theorem 6.5 and its extension 6.13 provide an affirmative answer. We factor the moment map of the Heisenberg action through natural finite Galois covers of the ramification type loci in the space of characteristic polynomials (equations (20) and (21)).

The compatibility naturally follows from the construction of the Poisson structure via level structures. Recall the birational realization of the moduli space $\text{Higgs}_{\Sigma}^{sm}(n, d, \omega(lP +$

D) as a quotient of the cotangent bundle $T^*\mathcal{U}_{lP+D}$ of the moduli space $\mathcal{U}_{lP+D} := \mathcal{U}_\Sigma(n, d, lP + D)$ of vector bundles with level structures (Chapter 5). This realization is a finite dimensional approximation of the limiting realization of the moduli space

$$\text{Higgs}_{\infty P+D} := \varinjlim_{l \rightarrow \infty} \text{Higgs}_{lP+D}$$

as a quotient of (a subset of) the cotangent bundle $T^*\mathcal{U}_{\infty P+D}$. The ramification type loci $\text{Higgs}_{\underline{n}D}$, their Galois covers $\widetilde{\text{Higgs}}_{\underline{n}D}$ and the infinitesimal $\text{Heis}_{\underline{n}}$ -action on $\widetilde{\text{Higgs}}_{\underline{n}D}$ become special cases of those appearing in the construction of section 2.5.2. The compatibility theorem follows from corollary 2.22 accompanied by the concrete identification of the moment maps in our particular example.

The rest of this section is organized as follows. In section 6.3.1 we emphasize the ubiquity of the setup of relative Krichever maps. They can be constructed for any family $\mathcal{J} \rightarrow B$ of Jacobians of branched covers of a fixed base curve Σ . The analogue of the compatibility question 6.1 makes sense whenever the family $\mathcal{J} \rightarrow B$ is an integrable system (see for example question 6.2).

Starting with section 6.3.2 we concentrate on the moduli spaces of Higgs pairs. Sections 6.3.2 and 6.3.3 consider the case of smooth spectral curves. Especially well behaved is the case where the point $P \in \Sigma$ of the \underline{n} -covering data is in the support of the polar divisor D . In this case the symplectic leaves foliation of the moduli space of Higgs pairs is a refinement of the type loci partition (lemma 6.3).

Section 6.3.4 is a generalization to singular cases. As an example, we consider in section 6.3.5 the Elliptic Solitons studied by Treibich and Verdier. We conclude with an outline of the proof of the compatibility theorem in section 6.3.6.

Note: Type- $(1, 1, \dots, 1)$ relative Krichever maps were independently considered by Y. Li and M. Mulase in a recent preprint [LM2].

6.3.1 Galois covers and relative Krichever maps

Let $B_D := \bigoplus_{i=1}^n H^0(\Sigma, (\omega_\Sigma(D))^{\otimes i})$ be the space of characteristic polynomials. For simplicity, we restrict ourselves to the Zariski open subset Bsm_D of reduced and irreducible n -sheeted spectral curves $\pi : C \rightarrow \Sigma$ in $T_\Sigma^*(D)$ whose fiber over $P \in \Sigma$ consists of *smooth* points of C . Denote by Higgsm_D the corresponding open subset of Higgs_D . The ramification type stratification of Higgsm_D is induced by that of Bsm_D

$$Bsm_D = \cup_{\underline{n}} Bsm_{\underline{n}D}.$$

Given a Higgs pair (E, φ) in $\text{Higgsm}_{\underline{n}D}$ corresponding to a torsion free sheaf L on a spectral cover $C \rightarrow \Sigma$ we can complete it to an \underline{n} -covering data

$$(C, P_i, z_i, L, \eta_i) \rightarrow (\Sigma, P, z, E, \eta)$$

by choosing i) a formal local parameter z at P , ii) an n_i -th root z_i of π^*z at each point P_i of C over P and iii) formal trivializations η_i of the sheaf L at P_i . The $\text{Heis}_{\underline{n}}^+$ -orbit of an \underline{n} -covering data consists precisely of all possible choices of the η_i 's. Thus, for fixed P and z only a finite choice is needed in order to obtain the points of the quotients of

the Grassmannians Gr_n/Heis_n^+ and $Gr_k/(\mathbb{C}[[z]]^\times)^k$ (see diagram 17). These choices are independent of the sheaf L . The choices are parametrized by the Galois cover $\widetilde{Bsm}_D^n \rightarrow Bsm_D^n$ consisting of pairs (C, λ) of a spectral curve C in Bsm^n and the discrete data λ which amounts to:

- i) (Parabolic data) An ordering (P_1, P_2, \dots, P_k) of the points (eigenvalues) in the fiber over P compatible with the fixed order of the ramification indices (n_1, \dots, n_k) (say, $n_1 \leq n_2 \leq \dots \leq n_k$).
- ii) A choice of an n_i -th root z_i of π^*z at P_i .

Denote by $\widetilde{\text{Higgs}}_D^n$ the corresponding Galois cover of Higgs_D^n . We get a canonical relative Krichever map

$$\kappa_n : \widetilde{\text{Higgs}}_D^n \rightarrow Gr_n/\text{Heis}_n^+ \quad (18)$$

from the Galois cover to the quotient Grassmannian.

The Galois group of $\widetilde{Bsm}_D^n \rightarrow Bsm_D^n$ is the Weyl group $W_n := N(\text{Heis}_n^+)/\text{Heis}_n^+$ of the maximal torus of the level infinity group G_∞^+ . For example, $W_{(1, \dots, 1)}$ is the symmetric group S_n , while $W_{(n)}$ is the cyclic group of order n . The discrete data $\lambda = [(P_1, P_2, \dots, P_k), (z_1, \dots, z_k)]$ of a point (C, λ) in \widetilde{Bsm}_D^n is equivalent to a commutative $\mathbb{C}[[z]]$ -algebras isomorphism

$$\lambda : \mathbf{heis}_n^+ \xrightarrow{\cong} \bigoplus_{i=1}^k \widehat{\mathcal{O}_{C, (P_i)}} \quad (19)$$

of the torus algebra with the formal completion of the structure sheaf \mathcal{O}_C at the fiber over P . The inverse λ^{-1} sends z_i to the generator of the i -th block of the torus \mathbf{heis}_n^+ given by (16). The finite Weyl group W_n acts on \mathbf{heis}_n^+ , hence on λ , introducing the W_n -torsor structure on \widetilde{Bsm}_D^n . (See also lemma 6.17 part 2 for a group theoretic interpretation.)

We note that a Galois cover $\widetilde{B} \rightarrow B$ as above can be defined for any family $\mathcal{J} \rightarrow B$ of Jacobians of a family $\mathcal{C} \rightarrow B$ of branched covers with a fixed ramification type \underline{n} of a fixed triple (Σ, P, z) . We obtain a relative Krichever map

$$\kappa_n : \widetilde{\mathcal{J}} \rightarrow Gr_n/\text{Heis}_n^+$$

as above. The Heisenberg flow pulls back to an infinitesimal action, i.e., a Lie algebra homomorphism

$$d\rho : \mathbf{heis}_n \rightarrow V(\widetilde{\mathcal{J}})$$

into a commutative algebra of vertical tangent vector fields. When \mathcal{J} (and hence $\widetilde{\mathcal{J}}$) is an integrable system we are led to ask the compatibility question 6.1: *is the action Poisson? i.e., can $d\rho$ be lifted to a Lie algebra homomorphism*

$$\mu_{\mathbf{heis}_n}^* : \mathbf{heis}_n \rightarrow \Gamma(\mathcal{O}_{\widetilde{B}}) \hookrightarrow [\Gamma(\mathcal{O}_{\widetilde{\mathcal{J}}}), \{, \}]?$$

A priori, the vector fields $d\rho(a)$, $a \in \mathbf{heis}_n$ may not even be *locally* Hamiltonian.

Inherently nonlinear examples arise from the Mukai-Tyurin integrable system of a family of Jacobians of a linear system $B := \mathbb{P}H^0(S, \mathcal{L})$ of curves on a symplectic or Poisson surface S (see chapter 8). Consider for example the:

Question 6.2 *Let $\pi : S \rightarrow \mathbb{P}^1$ be an elliptic K3 surface and \mathcal{L} a very ample line bundle on S . Fix $P \in \mathbb{P}^1$ and a local parameter z and consider the Galois cover $\tilde{B}^{(1,1,\dots,1)}$ of the generic ramification type locus. Is the Heisenberg action Poisson on $\tilde{\mathcal{J}} \rightarrow \tilde{B}^{(1,1,\dots,1)}$ (globally over $\tilde{B}^{(1,1,\dots,1)}$)?*

The compatibility question has an intrinsic algebro-geometric formulation: The j -th KP flow of the P_i -component is the vector field whose direction along the fiber over $(b, \lambda) \in \tilde{B}^{(1,1,\dots,1)}$ is the j -th derivative of the Abel-Jacobi map at P_i . Using the methods of chapter 8 it is easy to see that the Heisenberg action is symplectic. As we move the point $P(0) := P$ in \mathbb{P}^1 and its (Lagrangian) fiber in S we obtain an analytic (or formal) family of Lagrangian sections $\mathcal{AJ}(P_i(z))$ of $\tilde{\mathcal{J}} \rightarrow \tilde{B}^{(1,1,\dots,1)}$ (see corollary 8.13). Translations by the sections $\mathcal{AJ}(P_i(z)) - \mathcal{AJ}(P_i(0))$ is a family of symplectomorphisms of $\tilde{\mathcal{J}}$. Thus, the vector field corresponding to its j -th derivative with respect to the local parameter z is locally Hamiltonian. It seems unlikely however that the Heisenberg flow integrates to a global Poisson action for a general system as in question 6.2. It is the *exactness* of the symplectic structure in a neighborhood of the fiber over P in $T^*\Sigma$ which lifts the infinitesimal symplectic Heisenberg action to a Poisson action in the Hitchin's system case (see equation (21)).

6.3.2 Compatibility of stratifications

Prior to stating the compatibility theorem 6.5 we need to examine the Poisson nature of the Galois covers $\widetilde{\text{Higgs}}_D^{\underline{n}}$. $\text{Higgs}_D^{(1,1,\dots,1)}$ is a Zariski *open Poisson* subvariety of Higgs_D . Hence, the unramified Galois cover $\widetilde{\text{Higgs}}_D^{(1,1,\dots,1)}$ is endowed with the canonical pullback Poisson structure.

Though non generic, the other type strata are as important. The cyclic ramification type (n) , for example, corresponds to the single component KP-hierarchy (see [SW]). When the point P of the \underline{n} -covering data is in the support of the divisor D , we obtain a *strict compatibility* between the P -ramification type stratification of Higgs_D and its symplectic leaves foliation. All Galois covers $\widetilde{\text{Higgs}}_D^{\underline{n}}$, $P \in D$ are thus endowed with the canonical pullback Poisson structure:

Lemma 6.3 (*conditional compatibility of stratifications*) *When the point $P \in \Sigma$ is in the support of D , the symplectic leaves foliation is a refinement of the ramification type stratification $\text{Higgs}_D = \cup_{\underline{n}} \text{Higgs}_D^{\underline{n}}$.*

Proof: We need to show that the ramification type \underline{n} of the spectral cover $\pi : C \rightarrow \Sigma$ over $P \in D$ is fixed throughout the symplectic leaf $\text{Higgs}_S \subset \text{Higgs}_D$ of a Higgs pair (E_0, φ_0) . The symplectic leaves of Higgs_D are determined by coadjoint orbits of the level D algebra \mathfrak{g}_D . The coadjoint orbit S is determined by the residue of the Higgs field, namely, the infinitesimal data $(E|_D, \varphi|_D)$ encoded in the value of φ at D (see [Ma1] Remark 8.9 and Proposition 7.17)). Thus, the Jordan type of the Higgs field φ at P is fixed throughout Higgs_S . In general, (allowing singularities over P), the Jordan type depends both on the ramification type of $C \rightarrow \Sigma$ and on the sheaf L on C corresponding to the Higgs pair. For Higgs pairs (E, φ) in Higgs_D , the spectral curve C is smooth over P , hence, its ramification type coincides with the Jordan type of φ at P .

□

6.3.3 The compatibility theorem, the smooth case

We proceed to introduce the moment map of the infinitesimal Poisson action

$$d\rho : \mathbf{heis}_{\underline{n}} \rightarrow V(\widetilde{\text{Higgsm}}_{\underline{n}}^D).$$

Throughout the end of this subsection we will assume the

Condition 6.4 *Ramification types \underline{n} other than $(1, 1, \dots, 1)$ are considered only if P is in the support of D .*

This condition will be relaxed later by conditions 6.11 or 6.12.

Let $b \in \text{Bsm}_D^{\underline{n}}$ be the polynomial of the spectral curve $\pi_b : C_b \rightarrow \Sigma$. Recall that spectral curves are endowed with a tautological meromorphic 1-form $y_b \in H^0(C_b, \pi_b^* \omega_{\Sigma}(D))$ with poles over $D \subset \Sigma$ (see section 4.2). Let $\phi_{P_i}^j$ be the function on $\widetilde{\text{Bsm}}_D^{\underline{n}}$ given at a pair $(b, \lambda) \in \widetilde{\text{Bsm}}_D^{\underline{n}}$ by

$$\phi_{P_i}^j(b, \lambda) := \text{Res}_{P_i}((z_i)^{-j} \cdot y_b). \quad (20)$$

The Lie algebra homomorphism $\mu_{\mathbf{heis}_{\underline{n}}}^*$ sends the inverse (in $\mathbf{heis}_{\underline{n}}$) of the generator of the i -th block of the torus $\mathbf{heis}_{\underline{n}}^+$ given in (16) to the function $\phi_{P_i}^1 \circ \widetilde{\text{char}}$ on $\widetilde{\text{Higgsm}}_D^{\underline{n}}$. In other words, $\mu_{\mathbf{heis}_{\underline{n}}}^* : \mathbf{heis}_{\underline{n}} \rightarrow [\Gamma(\mathcal{O}_{\widetilde{\text{Higgsm}}_D^{\underline{n}}}), \{, \}]$ factors as a composition $\phi \circ \widetilde{\text{char}}^*$ through $\Gamma(\mathcal{O}_{\widetilde{\text{Bsm}}_D^{\underline{n}}})$. If we regard λ also as an isomorphism from $\mathbf{heis}_{\underline{n}}$ to $\bigoplus_{i=1}^k \widehat{\mathcal{O}}_{C, (P_i)}$ (eq. (19)), then $\phi : \mathbf{heis}_{\underline{n}} \rightarrow \Gamma(\mathcal{O}_{\widetilde{\text{Bsm}}_D^{\underline{n}}})$ is given by

$$(\phi(a))(b, \lambda) = \sum_{\{P_i\}} \text{Res}_{P_i}(\lambda(a) \cdot y_b), \quad a \in \mathbf{heis}_{\underline{n}}. \quad (21)$$

Theorem 6.5 (The Compatibility Theorem, smooth case) *(Assuming condition 6.4) The relative Krichever map*

$$\kappa_{\underline{n}} : \widetilde{\text{Higgsm}}_D^{\underline{n}} \rightarrow Gr_n / \text{Heis}_{\underline{n}}^+$$

intertwines the Heisenberg flow on $Gr_n / \text{Heis}_{\underline{n}}^+$ (and the mcKP flow on $Gr_k / (\mathbb{C}[[z]])^{\times k}$) with an infinitesimal Poisson action of the maximal torus $\mathbf{heis}_{\underline{n}}$ on $\widetilde{\text{Higgsm}}_D^{\underline{n}}$. The latter is induced by the Lie algebra homomorphism

$$\mu_{\mathbf{heis}_{\underline{n}}}^* = \phi \circ \widetilde{\text{char}}^* : \mathbf{heis}_{\underline{n}} \xrightarrow{\phi} \Gamma(\mathcal{O}_{\widetilde{\text{Bsm}}_D^{\underline{n}}}) \xrightarrow{\widetilde{\text{char}}^*} [\Gamma(\mathcal{O}_{\widetilde{\text{Higgsm}}_D^{\underline{n}}}), \{, \}]$$

which factors through the homomorphism ϕ given by (21).

Remarks 6.6 1. The subalgebra $\mathbf{heis}_{\underline{n}}^+$ acts trivially on $Gr_n / \text{Heis}_{\underline{n}}^+$ hence also on $\widetilde{\text{Higgsm}}_D^{\underline{n}}$. This corresponds to the fact that the functions $\phi_{P_i}^j$, $j \leq 0$ are Casimir. Indeed, if P is not contained in D , the 1-form y_b is holomorphic at the fiber over P and $\phi_{P_i}^j$ is identically zero for $j \leq 0$. If P is in D then the finite set of non-zero $\phi_{P_i}^j$, indexed by finitely many non-positive integers j , are among the Casimirs that induce the highest rank symplectic leaves foliation (see [Ma1] Proposition 8.8).

2. The multi-Krichever map $\kappa_{\underline{n}}$ depends on auxiliary parameters P and z . In other words, it lives naturally on an infinite dimensional space $\cup_{P,z} \widetilde{\text{Higgs}}_{D,P,z}^{\underline{n}}$ in which P and z are allowed to vary. This is not as bad as it might seem, since the j -th flow on $\widetilde{\text{Higgs}}_D^{\underline{n}}$ really depends only on our finite dimensional choices of P and the j -th order germ of z there. Similarly, our Hamiltonians $\phi_{P_i}^j$ depend at most on the $(j + n \deg D)$ -th germ of z , the shift arising, as in part 1 of this remark, from the possible poles above P of the tautological 1-form on the spectral curve. So we may think of $\cup_{P,z} \widetilde{\text{Higgs}}_{D,P,z}^{\underline{n}}$ as the inverse limit of a family of finite dimensional moduli spaces, indexed by the level. Each KP flow or Hamiltonian is defined for sufficiently high level.

6.3.4 The compatibility theorem, singular cases

The condition that the fiber over P of the embedded spectral curve be smooth is too restrictive. The embedded spectral data $(\bar{C} \subset T_{\Sigma}^*(D), \bar{L})$ of a Higgs pair (E, φ) may have singularities over P which are canonically resolvable. The point is that the rank 1 torsion free sheaf \bar{L} on \bar{C} determines a partial normalization $\nu : C \rightarrow \bar{C}$ and a unique rank 1 torsion free sheaf L on C such that i) \bar{L} is isomorphic to the direct image $\nu_* L$, and ii) L is locally free at the fiber over P . We are interested in those Higgs pairs for which the fiber of C over P is smooth. Such data may also be completed in finitely many ways to an \underline{n} -covering data as in section 6.3.1.

Definition 6.7 The singularities over P of a spectral pair $(\bar{C} \subset T_{\Sigma}^*(D), \bar{L})$ are said to be *resolved by the spectral sheaf \bar{L}* if i) \bar{C} is irreducible and reduced. ii) The sheaf \bar{L} is the direct image $\nu_* L$ of a rank 1 torsion free sheaf L on the normalization $\nu : C \rightarrow \bar{C}$ of the fiber of \bar{C} over P .

Fixing a symplectic leaf Higgs_S we may consider the type sub-loci in the locus of Higgs pairs whose spectral curve has at worst singularities over P which are resolved by the spectral sheaf. The topology of these type loci is quite complicated. As a result, the Galois covers of these type loci may not have a symplectic structure. Nevertheless, the construction of section 2.5.2, as used in section 6.3.6, provides *canonical embeddings* of the Galois covers of these type loci in (finite dimensional) symplectic varieties. These embeddings realize the Heisenberg flow as a Hamiltonian flow.

Control over the topology is regained below by restraining the singularities. In the smooth case (section 6.3.3) it is the smoothness which assures that the generic ramification type locus in a symplectic leaf Higgs_S is open (rather than only Zariski dense). The point is that degenerations from a ramification type \underline{n} through other types back to type \underline{n} must end with a singular fiber over P (and are thus excluded). If $P \in D$, there are symplectic leaves Higgs_S , $S \subset \mathfrak{g}_D^*$ of Higgs_D for which the singularities over P are encoded in the infinitesimal data associated to the coadjoint orbit S and shared by the generic Higgs pair in Higgs_S . Often, this is an indication that the Poisson surface $T_{\Sigma}^*(D)$ is not the best to work with. Moreover, a birational transformation $T_{\Sigma}^*(D) \rightarrow X_S$, centered at points of the fiber over P and encoded in S , can simultaneously resolve the singularities (over P) of the spectral curve of the generic point in Higgs_S (see example 6.9). In such a case, smoothness of the proper transform of the spectral curve in X_S at points of the

fiber over P is an *open* condition and the corresponding locus in Higgs_S with generic ramification type is *symplectic*. When the multiplicity of P in D is greater than 1 the correspondence between the coadjoint orbits $S \subset \mathfrak{g}_D^*$ and their surfaces X_S can be quite complicated. Instead of working the correspondence out, we will use the following notion of S -smoothness to assure (see condition 6.11) that the generic ramification type locus in a symplectic leaf Higgs_S is open (rather than only Zariski dense).

Definition 6.8 Let S be a coadjoint orbit of \mathfrak{g}_D . An irreducible and reduced spectral curve $\pi : \bar{C} \rightarrow \Sigma$ is *S -smooth over D* if i) a line bundle L on the resolution of the singularities $\nu : C \rightarrow \bar{C}$ of the fibers over D results in a Higgs pair $(E, \varphi) := \pi \circ \nu_*(L, \otimes \nu^*(y))$ in Higgs_S . and ii) the arithmetic genus of the normalization C above is equal to half the dimension of the symplectic leaf Higgs_S .

If \bar{C} is an irreducible and reduced spectral curve then, by the construction of [Sim1], the fiber of the characteristic polynomial map in $\text{Higgs}_\Sigma^{sm}(n, d, \omega(D))$ is its compactified Jacobian, the latter being the moduli space of all rank 1 torsion free sheaves on \bar{C} with a fixed Euler characteristic. The compactified Jacobian is known to be irreducible for irreducible and reduced curve on a surface (i.e., with planar singularities, [AIK]). Moreover, the symplectic leaf Higgs_S intersects the compactified Jacobian of \bar{C} in a union of strata determined by partial normalizations. If (E, φ) in Higgs_S corresponds to (\bar{C}, \bar{L}) and $\nu : (C, L) \rightarrow (\bar{C}, \bar{L})$ is a partial normalization where L is a locally free sheaf on C , then any twist F of L by a locally free sheaf in $\text{Pic}^0(C)$ (the component of \mathcal{O}_C) will result in a Higgs pair $(E', \varphi') := (\pi \circ \nu_*(F), \pi_*(\otimes y))$ in Higgs_S . The point is that the residue of (E', φ') (with respect to any level- D structure) will be in the same coadjoint orbit as that of (E, φ) . S -smoothness of \bar{C} over D is thus equivalent to the geometric condition:

The fiber of the characteristic polynomial map over \bar{C} intersects Higgs_S in a Lagrangian subvariety isomorphic to the compactified Jacobian of the resolution C of the singularities of \bar{C} over D .

The following example will be used in section 6.3.5 to describe a symplectic leaf which parametrizes Elliptic solitons.

Example 6.9 Let $D = P$, $S \subset \mathfrak{g}_D^* \cong \mathfrak{gl}(\mathbf{n})^* \cong \mathfrak{gl}(\mathbf{n})$ be the coadjoint orbit containing the diagonal matrix

$$A = \begin{pmatrix} -1 & 0 & \dots & 0 \\ 0 & -1 & & \\ & & \cdot & \vdots \\ \vdots & & & \cdot \\ & & & -1 & 0 \\ 0 & \dots & 0 & n-1 \end{pmatrix}$$

S is the coadjoint orbit of lowest dimension with characteristic polynomial $(x+1)^{n-1}(x-(n-1))$. Its dimension $2n-2$ is $(n-2)(n-1)$ less than the generic rank of the Poisson structure of $\mathfrak{gl}(\mathbf{n})^*$. If non-empty, each component of Higgs_S is a smooth symplectic variety of dimension $\dim \text{Higgs}_P - (n-1) - (n-2)(n-1) = [n^2(2g-1)+1] - (n-1)^2$ (see theorem 4.8 and [Ma1] proposition 7.17). The spectral curves $\bar{C} \subset T_\Sigma^*(P)$ which are S -smooth over P will have two points in the fiber over P , one smooth at the eigenvalue with

residue $n-1$ and one (singular if $n \geq 3$) at the eigenvalue with residue -1 . Assume $n \geq 3$. The resolution C of the singularity of a typical (though not all) such \bar{C} will be unramified over P with $n-1$ points collapsed to one in \bar{C} . The sheaf \bar{L} (of a Higgs pair in Higgs_S with spectral curve \bar{C}) will be a pushforward of a torsion free sheaf L on C . In contrast, line bundles on that \bar{C} will result in Higgs pairs in another symplectic leaf $\text{Higgs}_{S_{reg}}$ corresponding to the *regular* coadjoint orbit S_{reg} in $\mathfrak{gl}(\mathbf{n})^* \cong \mathfrak{gl}(\mathbf{n})$ with characteristic polynomial $(x+1)^{n-1}(x-(n-1))$. These Higgs pairs will *not* be S_{reg} -smooth. S_{reg} -smoothness coincides with usual smoothness of the embedded spectral curve which is necessarily ramified with ramification index $n-1$ at the point with residue -1 over P .

The S -smooth spectral curves will be smooth on the blowup \widehat{X}_S of $T_\Sigma^*(P)$ at residue -1 over P . If we blow down in \widehat{X}_S the proper transform of the fiber of $T_\Sigma^*(P)$ we get a surface X_S with a marked point x_{n-1} over P . An S -smooth spectral curve \bar{C} will correspond to a curve on X_S through x_{n-1} . It will be smooth at x_{n-1} if in addition it is of ramification type $(1, 1, \dots, 1)$ over P . Consider the compactification $\mathbb{P}(T_\Sigma^*(P) \oplus \mathcal{O}_\Sigma)$ of $T_\Sigma^*(P)$. Blowing it up and down as above we get a ruled surface \bar{X}_S over Σ which is isomorphic to the projectivization $\mathbb{P}W$ of the unique nontrivial extension

$$0 \rightarrow \omega_\Sigma \rightarrow W \rightarrow \mathcal{O}_\Sigma \rightarrow 0.$$

In particular, the surface \bar{X}_S is *independent of the point P* . The point is that blowing up and down the ruled surface $\mathbb{P}V := \mathbb{P}(T_\Sigma^*(P) \oplus \mathcal{O}_\Sigma)$ at residue -1 over P results with the ruled surface of a rank 2 vector bundle W whose sheaf of sections is a subsheaf of $V := T_\Sigma^*(P) \oplus \mathcal{O}_\Sigma$. This subsheaf consists of all sections which restrict at P to the subspace of the fiber $V|_P$ spanned by $(-1, 1)$ (i.e., W is a Hecke transform of V , see [Ty2]). Clearly, ω_Σ is a subsheaf of W and the quotient W/ω_Σ is isomorphic to \mathcal{O}_Σ . The resulting extension is non-trivial because $H^0(\Sigma, W)$ and $H^0(\Sigma, \omega_\Sigma)$ are equal as subspaces of $H^0(\Sigma, V)$.

Remark 6.10 S -smoothness over D is stronger than having singularities over D which are resolved by the spectral sheaf. They differ when the singularity appears at an infinitesimal germ of too high an order to be detected by S . E.g., take $n = 3$ in example 6.9 and consider a pair (\bar{C}, \bar{L}) with a tacnode at residue -1 over P (two branches meet with a common tangent). The arithmetic genus of the normalization $\nu : C \rightarrow \bar{C}$ of the fiber over P will drop by 2 while the pushforward $\bar{L} := \nu_*(L)$ of a line bundle L on C will belong to a symplectic leaf Higgs_S whose rank is 2 less than the maximal rank (rather than 4). Hence (\bar{C}, \bar{L}) is S -singular.

We denote by $\text{Higgs}_{S/D}$ the subset of $\text{Higgs}_S \subset \text{Higgs}_D$ parametrizing Higgs pairs whose spectral curve is S -smooth over D .

Unfortunately, the compatibility of stratifications (lemma 6.3) does not extend to the S -smooth case. To overcome this inconvenience we may either assume condition 6.11 or condition 6.12.

Condition 6.11 *Consider the ramification locus $\text{Higgs}_{S/D}^n$ in a component of $\text{Higgs}_{S/D}$ only if it is the generic ramification type in this component.*

Note that $\text{Higgs}_{S/D}$ parametrizes only Higgs pairs whose spectral curve is S -smooth over D . S -smoothness over D assures that if the type \underline{n} is a generic ramification type in a component of $\text{Higgs}_{S/D}$, then the corresponding component of $\text{Higgs}_{S/D}^{\underline{n}}$ is an *open* subset of $\text{Higgs}_{S/D}$ (i.e., it excludes degenerations of type \underline{n} Higgs pairs through other types back to type \underline{n}). In particular, these components of $\text{Higgs}_{S/D}^{\underline{n}}$ are *symplectic*. Alternatively, we may relax condition 6.11 even further at the expense of losing the symplectic nature of the loci and having to resort to convention 2.20:

Condition 6.12 *Consider only Higgs pairs with a spectral curve whose singularities over P are resolved by its spectral sheaf. Adopt convention 2.20.*

Theorem 6.13 *The compatibility theorem 6.5 holds for: i) The $W_{\underline{n}}$ -Galois covers of the type loci in $\text{Higgs}_{S/D}$ satisfying the genericity condition 6.11 (instead of condition 6.4). ii) The $W_{\underline{n}}$ -Galois covers of the locus in $\text{Higgs}_D^{\underline{n}}$ consisting of Higgs pairs satisfying condition 6.12. In ii) however we adopt convention 2.20.*

6.3.5 Elliptic Solitons

In this subsection we illustrate the possibilities in the singular case with a specific example. Let Σ be a smooth elliptic curve. A Σ -periodic *Elliptic KP soliton* is a finite dimensional solution to the KP hierarchy, in which the orbit of the first KP equation is isomorphic to Σ . Its Krichever data $(C, \tilde{P}, \frac{\partial}{\partial z}, L)$ consists of a reduced and irreducible curve C , a smooth point \tilde{P} , a nonvanishing tangent vector at \tilde{P} and a rank 1 torsion free sheaf L on C of Euler characteristic 0 (we suppress the non essential formal trivialization η and consider only the first order germ of z which is equivalent to choosing a nonzero tangent vector at P). We will denote the global vector field extending $\frac{\partial}{\partial z}$ also by $\frac{\partial}{\partial z}$. The periodicity implies that the image of the Abel Jacobi map $AJ : C \hookrightarrow J_C, Q \mapsto Q - \tilde{P}$ is tangent at 0 to a subtorus isomorphic to Σ . Composing the Abel Jacobi map with projection to Σ we get a *tangential morphism* $\pi : (C, \tilde{P}) \rightarrow (\Sigma, P)$. Its degree n is called the *order* of the Elliptic soliton. In general, a tangential morphism $\pi : (C, \tilde{P}) \rightarrow (\Sigma, P)$ is a morphism with the property that $AJ(C)$ is tangent at $AJ(\tilde{P})$ to $\pi^* J_{\Sigma}^0$. Notice that composing a tangential morphism $\pi : (C, \tilde{P}) \rightarrow (\Sigma, P)$ with a normalization $\nu : \tilde{C} \rightarrow C$ results in a tangential morphism. A tangential morphism is called *minimal* if it does not factor through another tangential morphism.

The KP elliptic solitons enjoyed a careful and detailed study by A. Treibich and J.-L. Verdier in a series of beautiful papers (e.g., [TV1, TV2]). Their results fit nicely with our picture:

Theorem 6.14 *The variety of Krichever data of Elliptic KP solitons of order n with a fixed pointed elliptic curve and a tangent vector $(\Sigma, P, \frac{\partial}{\partial z})$ is canonically birational to the divisor of traceless Higgs pairs in the symplectic leaf Higgs_S of $\text{Higgs}_{\Sigma}^{sm}(n, 0, \omega_{\Sigma}(P))$ corresponding to the coadjoint orbit S of example 6.9. The KP flows are well defined on $\text{Higgs}_{S/P} \subset \text{Higgs}_S$ as the Hamiltonian vector fields of the functions $\phi_{\tilde{P}}^j$ given in (20).*

Note that if non-empty (which is the case) Higgs_S is $2n$ -dimensional (see example 6.9). The correspondence between tangential covers and spectral covers is a corollary of the

following characterization of tangential covers due to I. M. Krichever and A. Treibich. For simplicity we consider only the case in which the tangency point \tilde{P} is not a ramification point of π .

Theorem 6.15 [TV2] *Assume that $\pi : C \rightarrow \Sigma$ is unramified at \tilde{P} . Then π is tangential if and only if there exists a section $y \in H^0(C, \pi^*[\omega_\Sigma(P)])$ satisfying:*

- a) *Near a point of $\pi^{-1}(P) - \tilde{P}$ (away from the tangency point \tilde{P}), $y - \pi^*(dz/z)$ is a holomorphic multiple of $\pi^*(dz)$, where z is a local parameter at P . (If $\pi : C \rightarrow \Sigma$ is unramified over P , this is equivalent to saying that the residues $\text{Res}_{P_i}(y)$ are the same at all P_i other than \tilde{P} in the fiber over $P \in \Sigma$).*
- b) *The residue $\text{Res}_{\tilde{P}}(y)$ at \tilde{P} does not vanish if $n \geq 2$.*

It follows by the residue theorem that there is a unique such section which has residue $n - 1$ at \tilde{P} and which is moreover traceless $\text{tr}(y) = 0 \in H^0(\Sigma, \omega_\Sigma(P))$. Let dz be a global holomorphic non zero 1-form on Σ . The function $k := y/\pi^*(dz)$ is called a *tangential function* in [TV2]. It was also proven that a tangential morphism of order n has arithmetic genus $\leq n$ and is minimal if and only if its arithmetic genus is n . ([TV1] Corollaire 3.10).

Sketch of proof of theorem 6.15: (for C smooth, $\pi : C \rightarrow \Sigma$ unramified over P .)

Step 1: (Cohomological identification of the differential of the Abel-Jacobi map)

The differential $dAJ : T_Q C \cong H^0(Q, \mathcal{O}_Q(Q)) \rightarrow H^1(C, \mathcal{O}_C)$ of the Abel-Jacobi map at $Q \in C$ is identified as the connecting homomorphism of the short exact sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_C(Q) \rightarrow \mathcal{O}_Q(Q) \rightarrow 0.$$

Similarly, the differential $d(AJ \circ \pi^{-1})$ of the composition $\Sigma \xrightarrow{\pi^{-1}} \text{Sym}^n C \xrightarrow{AJ} J_C^n$ is given by

$$T_P \Sigma \cong H^0(P, \mathcal{O}_P(P)) \xrightarrow{\pi^*} H^0(\pi^{-1}(P), \mathcal{O}_{\pi^{-1}(P)}(\pi^{-1}(P))) \xrightarrow{\cong} T_{[\pi^{-1}(P)]} \text{Sym}^n C \xrightarrow{dAJ} H^1(C, \mathcal{O}_C),$$

where the composition of the last two arrows is the connecting homomorphism of the short exact sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_C(\pi^{-1}(P)) \rightarrow \mathcal{O}_{\pi^{-1}(P)}(\pi^{-1}(P)) \rightarrow 0.$$

Step 2: (residues as coefficients in a linear dependency of tangent lines)

Clearly, the tangent line to $(AJ \circ \pi^{-1})(\Sigma)$ at the image of P is in the span of the tangent lines to $AJ(C)$ at the points P_i . If, in addition, π is tangential with tangency point $\tilde{P} \in C$, then the tangent lines to $AJ(C)$ at the points in the fiber over P are linearly dependent. If $\pi : C \rightarrow \Sigma$ is unramified over P we can write these observations in the form of two linear equations:

$$\sum dAJ_{P_i} \left(\frac{\partial}{\partial z_i} \right) = d(AJ \circ \pi^{-1})_P \left(\frac{\partial}{\partial z} \right), \quad (22)$$

and

$$\sum a_i dAJ_{P_i} \left(\frac{\partial}{\partial z_i} \right) = 0 \quad \text{linear dependency.} \quad (23)$$

Above, $\frac{\partial}{\partial z_i}$ is the lift of $\frac{\partial}{\partial z}$, i.e., $d\pi \left(\frac{\partial}{\partial z_i} \right) = \frac{\partial}{\partial z}$. We claim that the coefficients a_i in (23) are residues of a meromorphic 1-form y at the points of the fiber. More precisely we have:

Lemma 6.16 *Assume that $\pi : C \rightarrow \Sigma$ is unramified over P . There exists a section $y \in H^0(C, \pi^* \omega_\Sigma(P))$ with residues (a_1, a_2, \dots, a_n) at the fiber over P if and only if (a_1, a_2, \dots, a_n) satisfy equation (23).*

Proof: The global tangent vector field $\frac{\partial}{\partial z} \in H^0(\Sigma, T\Sigma)$ gives rise to the commutative diagram:

$$\begin{array}{ccccc}
 H^0(C, \pi^* \omega_\Sigma(P)) & \longrightarrow & H^0(\pi^{-1}(P), \pi^* \omega_\Sigma(P)|_{\pi^{-1}(P)}) & \longrightarrow & H^1(C, \pi^* \omega_\Sigma) \\
 \cong \downarrow \downarrow \frac{\partial}{\partial z} & & \cong \downarrow \downarrow \frac{\partial}{\partial z} & & \cong \downarrow \downarrow \frac{\partial}{\partial z} \\
 H^0(C, \mathcal{O}_C(\pi^{-1}(P))) & \rightarrow & H^0(\pi^{-1}(P), \mathcal{O}_{\pi^{-1}(P)}(\pi^{-1}(P))) & \xrightarrow{dAJ} & H^1(C, \mathcal{O}_C).
 \end{array} \tag{24}$$

The middle contraction $\downarrow \frac{\partial}{\partial z}$ maps residues $(a_1, a_2, \dots, a_n) \in H^0(\pi^{-1}(P), \pi^* \omega_\Sigma(P)|_{\pi^{-1}(P)})$ to $(a_1 \frac{\partial}{\partial z_1}, a_2 \frac{\partial}{\partial z_2}, \dots, a_n \frac{\partial}{\partial z_n})$. The lemma follows by the exactness of the horizontal sequences in the diagram.

□

Step 3: We conclude that $\pi : C \rightarrow \Sigma$ is tangential if and only if there is a 1-form y as in the theorem. If $\pi : C \rightarrow \Sigma$ is tangential then using lemma 6.16 we see that equation (22) gives rise to a 1-form y with residues $(-1, -1, \dots, n-1)$ as required. Conversely, given a 1-form y with residues $(-1, -1, \dots, n-1)$ lemma 6.16 and equations (23) and (22) imply the tangentiality. This completes the proof of theorem 6.15 in the generic case considered.

□

Theorem 6.14 would follow once the existence of either n -sheeted spectral covers S -smooth over P , or degree n tangential covers smooth and unramified over P is established for every choice of (Σ, P) . This was done in ([TV1] Theorem 3.11) by studying the linear system of transferred spectral curves on the surface \bar{X}_S of example 6.9 and applying Bertini's theorem to show that the generic transferred spectral curve is smooth in X_S . It follows that the blow up of the point with residue -1 over P resolves the generic n -sheeted spectral curve of Higgs pairs in Higgs_S to a smooth curve of genus n unramified over P .

Any S -smooth spectral curve \bar{C} in $T^*\Sigma(P)$ of a Higgs pair (E, φ) in Higgs_S admits a unique partial normalization C of arithmetic genus n by the spectral sheaf \bar{L} corresponding to (E, φ) . The tautological 1-form \bar{y} pulls back to a 1-form on C of the type which characterize tangential covers by theorem 6.15.

Conversely, a degree n tangential cover $\pi : (C, \tilde{P}) \rightarrow (\Sigma, P)$ of arithmetic genus n which is smooth and unramified over P is sent to the spectral curve \bar{C} in $T^*\Sigma(P)$ of the Higgs pair

$$(E, \varphi) := (\pi_*(L), [\otimes y : \pi_*(L) \rightarrow \pi_*(L) \otimes \omega_\Sigma(P)])$$

for some, hence every, choice of a line bundle L on C . Note that \bar{C} is reduced since it is irreducible and the branch through residue $n-1$ over P is reduced. The canonical morphism $\nu : C \rightarrow \bar{C}$ is the resolution by the spectral sheaf $\nu_* L$. Hence $(\bar{C}, \nu_* L)$ is

S -smooth. (The arithmetic genus of \bar{C} is $\frac{1}{2}(n^2 - n + 2)$, the common arithmetic genus to all n -sheeted spectral curves in $T^*\Sigma(P)$.)

Finally we note that, as the tangency point \tilde{P} over P is marked by having residue $n - 1$, it does not have monodromy and all the KP flows corresponding to it are well define on $\text{Higgs}_{S/P}$ as the Hamiltonian vector fields of the functions $\phi_{\tilde{P}}^j$ given in (20).

6.3.6 Outline of the proof of the compatibility theorem

For simplicity we assume that $D = lP$, $l \geq 0$. The general case is similar replacing $\mathcal{U}_\Sigma(n, d, \infty P)$ by $\mathcal{U}_\Sigma(n, d, \infty P + D)$. Let $G_\infty := GL(n, K)$ be the loop group and G_∞^+ the level infinity subgroup. (More canonically, $K \cong \mathbb{C}((z))$ should be thought of as the completion of the function field of Σ at P , and we may postpone the choice of a coordinate z until we need to choose generators for a maximal torus $\text{Heis}_{\underline{n}}$ of G_∞ .)

Denote by $M_{l,k}$, $k \geq l$ the pullback of $T^*\mathcal{U}_\Sigma(n, d, lP)$ to $\mathcal{U}_\Sigma(n, d, kP)$ via the rational forgetful morphism. $M := T^*\mathcal{U}_\Sigma(n, d, \infty P)$ is defined as the limit of finite dimensional approximations (see 2.5.1 for the terminology)

$$T^*\mathcal{U}_\Sigma(n, d, \infty P) := \lim_{l \rightarrow \infty} \lim_{\infty \leftarrow k} M_{l,k}.$$

Denote by M^s (resp. $M_{l,k}^s$) the subset of $T^*\mathcal{U}_\Sigma(n, d, \infty P)$ (resp. $M_{l,k}$) consisting of triples (E, φ, η) with a *stable* Higgs pair (E, φ) . We arrive at the setup of section 2.5.1 with the Poisson quotient $Q_\infty := M^s/G^+$ being the direct limit

$$\text{Higgs}_\infty := \lim_{l \rightarrow \infty} \text{Higgs}_\Sigma^{sm}(n, d, lp).$$

We emphasize that the stability condition is used here for the morphism

$$M_{l,k}^s \rightarrow \text{Higgs}_\Sigma^{sm}(n, d, lp)$$

between the two *existing* coarse moduli spaces to be well defined, and *not* to define the quotient.

The *infinitesimal* loop group action on $\mathcal{U}_{\infty P} := \mathcal{U}_\Sigma(n, d, \infty P)$ (the derivative of the action defined in section 6.2 on the level of Čech 1-cocycles) may be lifted to an infinitesimal action on its cotangent bundle. The point is that the infinitesimal action of $a \in \mathfrak{g}_\infty$, with poles of order $\leq l_0$, is well defined on the finite dimensional approximation $\mathcal{U}_\Sigma(n, d, lP)$ for $l \geq l_0$. Thus, it lifts to all cotangent bundles $M_{l,l}$, $l \geq l_0$. $M_{l,k}$ embeds naturally in $M_{k,k}$ as an invariant subvariety. This defines the action on the limit $T^*\mathcal{U}_\Sigma(n, d, \infty P)$.

As a lifted action it is automatically Poisson. Its moment map

$$\mu_\infty^* : \mathfrak{g}_\infty \rightarrow \Gamma[\mathcal{O}_{T^*\mathcal{U}_{\infty P}}, \{, \}]$$

(the limit of the moment maps for the finite dimensional approximations) is given by the same formula that we have already encountered for the level groups (see 14):

$$(\mu_\infty^*(a))(E, \varphi, \eta) = \text{Res}_P \text{trace}(a \cdot (\varphi)^\eta), \quad a \in \mathfrak{g}_\infty. \quad (25)$$

Choosing a maximal torus $\mathbf{heis}_{\underline{n}} \subset \mathfrak{g}_\infty$ of type \underline{n} we arrive at the setup of section 2.5.2. In particular, we obtain the type locus $\text{Higgs}_\Sigma^{\underline{n}}(n, d, lp)$ in $\text{Higgs}_\Sigma^{sm}(n, d, lp)$.

Lemma 6.17 1. *The algebro-geometric definition of the ramification type loci coincides with the group theoretic definition 2.18 when $\text{char}(E, \varphi)$ is an integral (irreducible and reduced) spectral curve.*

2. *A choice of generators for a maximal torus $\mathbf{heis}_{\underline{n}}$ as in (16) determines a canonical isomorphism between the group theoretic and the algebro-geometric $W_{\underline{n}}$ -Galois covers.*

Proof: 1) The stabilizer $\mathfrak{t} \subset \mathfrak{g}_{\infty}$ of $(\varphi)^{\eta} \in \mathfrak{g}_{\infty} \otimes_{\mathcal{O}_{(P)}} \omega_{\Sigma, (P)} \cong \mathfrak{g}_{\infty}^*$ with spectral curve $\pi : C = \text{char}(E, \varphi) \rightarrow \Sigma$ is precisely U^{η} where U is the stalk of $\text{Ker}[ad\varphi : \text{End}E \rightarrow \text{End}E \otimes \omega_{\Sigma}(lP)]$ at the formal punctured neighborhood of P . In addition, U is canonically isomorphic to the stalk of the structure sheaf at the formal punctured neighborhood of the fiber of C over P via the completion of the canonical embedding:

$$\pi_* \mathcal{O}_C \hookrightarrow \pi_* \text{End}L \hookrightarrow \text{End}E.$$

Hence, the level infinity structure η provides a *canonical* isomorphism

$$\lambda : \mathfrak{t} \xrightarrow{\cong} U \tag{26}$$

from the stabilizer algebra \mathfrak{t} to the structure sheaf at the formal punctured neighborhood of the fiber of C over P .

2) As the types coincide, we may choose the level infinity structure η so that the stabilizer \mathfrak{t} coincides with the fixed torus $\mathbf{heis}_{\underline{n}}$. We may further require that the isomorphism λ given by (26) coincides with the one in (19). This determines the $\text{Heis}_{\underline{n}}^+$ -orbit of η uniquely, i.e., a point in the group theoretic Galois cover. □

Theorems 6.5 and 6.13 would now follow from corollary 2.22 provided that we prove that the homomorphism ϕ of the theorems (given by 21) is indeed the factorization of the $\mathbf{heis}_{\underline{n}}$ -moment map through the characteristic polynomial map. (Note that the existence of this factorization follows from diagram (10)). In other words, we need to prove the identity

$$\sum_{\{P_i\}} \text{Res}_{P_i}(\lambda(a) \cdot y_b) = \text{Res}_P \text{trace}(a \cdot (\varphi)^{\eta}) \quad a \in \mathbf{heis}_{\underline{n}}, \quad b = \text{char}(E, \varphi) \tag{27}$$

as functions on the set of all Higgs pairs (E, φ, λ) in $\widetilde{\text{Higgs}}_{lP}^{\underline{n}}$ for which the spectral sheaf resolves the singularities of their spectral curve over P (see definition 6.7). Above, η is any level infinity structure in the $\text{Heis}_{\underline{n}}^+$ -orbit as in the proof of lemma 6.17 or, equivalently, $\lambda(a) = \eta^{-1} \circ a \circ \eta$ where we identify the structure sheaf of the formal punctured neighborhood with U of that lemma. If the embedded spectral curve \bar{C}_b is singular, the P_i are the points over P of its resolution $\nu : C_b \rightarrow \bar{C}_b$, and the tautological meromorphic 1-form y_b should be replaced by the pullback $\nu^*(y)$ of the tautological 1-form y on the surface $T_{\Sigma}^*(lP)$.

Conjugating the right hand side of (27) by η , we get

$$\sum_{\{P_i\}} \text{Res}_{P_i}(A \cdot \nu^*(y)) = \text{Res}_P \text{trace}((\pi \circ \nu)_*[A \cdot \nu^*(y)])$$

for A a (formal) meromorphic function at the fiber over P . Working formally, we can consider only the “parts” with first order pole $r_i d\log z_i$ of $A \cdot \nu^*(y)$ at P_i . The equality follows from the identity $d\log z_i = (\pi \circ \nu)^*[\frac{1}{n_i} d\log z]$ which imply (projection formula) that $(\pi \circ \nu)_*(\otimes d\log z_i)$ acts as $\frac{1}{n_i} e_{P_i} \otimes d\log z$ were e_{P_i} is the projection onto the eigenspace of the point P_i .

□

7 The Cubic Condition and Calabi-Yau threefolds

We pose in section 7.1 the general question: when does a family of polarized abelian varieties or complex tori support a completely integrable system? In section 7.2 we describe a general necessary infinitesimal symmetry condition on the periods of the family (the cubic condition of lemmas 7.1 and 7.2) and a sufficient local condition (lemmas 7.4 and 7.5).

In section 7.3 we use the Yukawa cubic to construct a symplectic structure (and an ACIHS) on the relative intermediate Jacobian over the moduli space of gauged Calabi-Yau threefolds (theorem 7.7). The symplectic structure extends to the bundle of Deligne cohomologies and we show that the image of the relative cycle map as well as bundles of sub-Hodge-structures are isotropic (corollary 7.11).

7.1 Families of Tori

Consider a Poisson manifold (X, ψ) together with a Lagrangian fibration

$$\pi : \mathcal{X} \longrightarrow B$$

over a base B , whose fibers

$$X_b := \pi^{-1}(b), \quad b \in B$$

are tori. (We say π is Lagrangian if each fiber X_b is a Lagrangian submanifold of some symplectic leaf in \mathcal{X} .) All these objects may be C^∞ , or may be equipped with a complex analytic or algebraic structure.

On B we have the tangent bundle \mathcal{T}_B as well as the vertical bundle \mathcal{V} , whose sections are vector fields along the fibers of π which are constant on each torus. The pullback $\pi^*\mathcal{V}$ is the relative tangent bundle $\mathcal{T}_{\mathcal{X}/B}$; in the analytic or algebraic situations, we can define \mathcal{V} as $\pi_*\mathcal{T}_{\mathcal{X}/B}$. The data π and ψ determine an injection

$$i : \mathcal{V}^* \hookrightarrow \mathcal{T}_B$$

or, equivalently, a surjection

$$i' : \mathcal{T}_B^* \rightarrow \mathcal{V}$$

sending a 1-form α on B to the vertical vector field

$$i'(\alpha) := \pi^*\alpha \rfloor \psi.$$

The image $i(\mathcal{V}^*) \subset \mathcal{T}_B$ is an integrable distribution on B . Its integral manifolds are the images in B of symplectic leaves in \mathcal{X} .

In this section we start with a family of tori $\pi : \mathcal{X} \rightarrow B$ and ask whether there is a *Lagrangian structure* for π , i.e. a Poisson structure on \mathcal{X} making the map π Lagrangian. More precisely, we fix $\pi : \mathcal{X} \rightarrow B$ and an injection $i : \mathcal{V}^* \hookrightarrow \mathcal{T}_B$ with integrable image, and ask for existence of a Lagrangian structure ψ on \mathcal{X} inducing the given i .

In the C^∞ category there are no local obstructions to existence of a Lagrangian structure: the fibration π is locally trivial, so one can always find action-angle coordinates near each fiber, and use them to define ψ . In the analytic or algebraic categories, on the other hand, the fibers X_b (complex tori, or abelian varieties) have invariants, given

essentially by their *period matrix* $p(X_b)$, so the fibration may not be analytically locally trivial. We will see that there is an obstruction to existence of a Lagrangian structure for $\pi : \mathcal{X} \rightarrow B$, which we formulate as a symmetry condition on the derivatives of the period map p . These derivatives can be considered as a linear system of quadrics, and the condition is, roughly, that they be the polars of some cubic (= section of $\text{Sym}^3\mathcal{V}$).

Let X be a g -dimensional complex torus, and $\gamma_1, \dots, \gamma_{2g}$ a basis of the integral homology $H_1(X, \mathbb{Z})$. There is a unique basis $\alpha_1, \dots, \alpha_g$ for the holomorphic differentials $H^0(X, \Omega_X^1)$ satisfying

$$\int_{\gamma_{g+i}} \alpha_j = \delta_{ij}, \quad 1 \leq i, j \leq g,$$

so we define the period matrix $P = p(X, \gamma)$ by

$$p_{ij} := \int_{\gamma_i} \alpha_j, \quad 1 \leq i, j \leq g.$$

Riemann's first and second bilinear relations say that X is a principally polarized abelian variety (PPAV) if and only if P is in *Siegel's half space*:

$$\mathbb{H}_g := \{\text{symmetric } g \times g \text{ complex matrices whose imaginary part is positive definite}\}.$$

In terms of a dual basis $\gamma_1^*, \dots, \gamma_{2g}^*$ of $H^1(X, \mathbb{Z})$, the integral class $\omega := \sum_{i=1}^g \gamma_i^* \wedge \gamma_{g+i}^* \in H^2(X, \mathbb{Z})$ is a Kähler class if and only if P is in Siegel's half space. In this case we call ω a *principal polarization*.

Given a family $\pi : \mathcal{X} \rightarrow B$ of PPAVs together with a continuously varying family of symplectic bases $\gamma_1, \dots, \gamma_{2g}$ for the fiber homologies, we then get a period map

$$p : B \longrightarrow \mathbb{H}_g.$$

If we change the basis γ by a symplectic transformation

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(2g, \mathbb{Z}),$$

the period matrix P goes to $(AP + B)(CP + D)^{-1}$. So given a family π without the choice of γ , we get a multi-valued map of B to \mathbb{H}_g , or a map

$$p : B \longrightarrow \mathcal{A}_g$$

to the moduli space of PPAV. The latter is a quasi projective variety, which can be described analytically as the quotient

$$\mathcal{A}_g = \mathbb{H}_g / \Gamma$$

of \mathbb{H}_g by the action of the modular group

$$\Gamma := Sp(2g, \mathbb{Z}) / (\pm 1).$$

A PPAV X determines a point $[X]$ (or carelessly, X) of \mathcal{A}_g . This point is non-singular if X has no automorphisms other than ± 1 , and then the tangent space $T_{[X]}\mathcal{A}_g$ can be identified with $\text{Sym}^2 V_X$, where V_X is the tangent space (at $0 \in X$) to X . This can be seen

by identifying $T_{[X]}\mathcal{A}_g$ with $T_{[X]}\mathbb{H}_g$ and recalling that \mathbb{H}_g is an open subset of Sym^2V_X . More algebraically, this follows from elementary deformation theory: all first-order deformations of X are given by

$$H^1(X, \mathcal{T}_X) \approx H^1(X, V_X \otimes_{\mathbb{C}} \mathcal{O}_X) \approx V_X \otimes H^1(X, \mathcal{O}_X) \approx \otimes^2 V_X,$$

and in there the deformations as abelian variety, i.e., the deformations preserving the polarization bilinear form on $H_1(X, \mathbb{Z})$, are given by the symmetric tensors Sym^2V_X .

7.2 The Cubic Condition

Our condition for an analytic or algebraic family $\pi : \mathcal{X} \rightarrow B$ of PPAVs, given by a period map $p : B \rightarrow \mathcal{A}_g$, to have a Lagrangian structure ψ inducing a given $i : \mathcal{V}^* \hookrightarrow \mathcal{T}_B$, can now be stated as follows. The differential of p is a map of bundles:

$$dp : \mathcal{T}_B \longrightarrow \text{Sym}^2\mathcal{V},$$

so the composite

$$dp \circ i : \mathcal{V}^* \longrightarrow \text{Sym}^2\mathcal{V}$$

can be considered as a section of $\mathcal{V} \otimes \text{Sym}^2\mathcal{V}$, and the condition is that it should come from the subbundle $\text{Sym}^3\mathcal{V}$. In other words, there should exist a cubic $c \in H^0(B, \text{Sym}^3\mathcal{V})$ whose polar quadrics give the directional derivatives of the period map: if the tangent vector $\partial/\partial b \in T_b B$ equals $i(\beta)$ for some $\beta \in \mathcal{V}^*$, then:

$$\frac{\partial p}{\partial b} = \beta \rfloor c.$$

We give two versions of this cubic condition. In the first, we check the existence of a two vector ψ , not necessarily satisfying the Jacobi identity, for which π is Lagrangian, and which induces a given injection $i : \mathcal{V}^* \hookrightarrow \mathcal{T}_B$. (Note that neither the definition of the map i induced by the two-vector ψ , nor the notion of π being Lagrangian, require ψ to be Poisson.)

Lemma 7.1 (*Weak cubic condition, Poisson form*). *A family $\pi : \mathcal{X} \rightarrow B$ of polarized abelian varieties has a two vector ψ satisfying*

- a) $\pi : \mathcal{X} \rightarrow B$ is Lagrangian
- b) ψ induces a given $i : \mathcal{V}^* \hookrightarrow \mathcal{T}_B$

if and only if

$$dp \circ i \in \text{Hom}(\mathcal{V}^*, \text{Sym}^2\mathcal{V})$$

comes from a cubic

$$c \in H^0(B, \text{Sym}^3\mathcal{V}).$$

Moreover, in this case there is a unique such 2-vector ψ which satisfies also

- c) *The zero section $z : B \rightarrow \mathcal{X}$ is Lagrangian, i.e., $(T_{\mathcal{X}/B}^*)|_{z(B)}$ is ψ -isotropic (here we identify the conormal bundle of the zero section with $(T_{\mathcal{X}/B}^*)|_{z(B)}$.)*

Proof: (Note: we refer below to the vertical bundle $\mathcal{T}_{\mathcal{X}/B}$ by its, somewhat indirect, realization as the pullback $\pi^*\mathcal{V}$.) The short exact sequence of sheaves on \mathcal{X} :

$$0 \rightarrow \pi^*\mathcal{V} \rightarrow \mathcal{T}_{\mathcal{X}} \rightarrow \pi^*\mathcal{T}_B \rightarrow 0$$

determines a subsheaf \mathcal{F} of $\overset{2}{\wedge} \mathcal{T}_{\mathcal{X}}$ which fits in the exact sequences:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathcal{F} & \rightarrow & \overset{2}{\wedge} \mathcal{T}_{\mathcal{X}} & \rightarrow & \pi^* \overset{2}{\wedge} \mathcal{T}_B \rightarrow 0 \\ 0 & \rightarrow & \pi^* \overset{2}{\wedge} \mathcal{V} & \rightarrow & \mathcal{F} & \rightarrow & \pi^*(\mathcal{V} \otimes \mathcal{T}_B) \rightarrow 0. \end{array}$$

The map π is Hamiltonian with respect to the two-vector $\psi \in H^0(B_{\mathcal{X}}, \overset{2}{\wedge} \mathcal{T}_{\mathcal{X}})$ if and only if ψ goes to 0 in $\overset{2}{\wedge} \mathcal{T}_B$, i.e., if and only if it comes from $H^0(\mathcal{F})$. The question is therefore whether $i \in H^0(B, \mathcal{V} \otimes \mathcal{T}_B) \subset H^0(\mathcal{X}, \pi^*(\mathcal{V} \otimes \mathcal{T}_B))$ is in the image of $H^0(\mathcal{X}, \mathcal{F})$. Locally in B , this happens if and only if i goes to 0 under the coboundary map

$$\begin{array}{ccc} \pi_* \pi^*(\mathcal{V} \otimes \mathcal{T}_B) & \longrightarrow & R^1 \pi_* \pi^* \overset{2}{\wedge} \mathcal{V} \\ \parallel & & \parallel \\ \mathcal{V} \otimes \mathcal{T}_B & \longrightarrow & \overset{2}{\wedge} \mathcal{V} \otimes \mathcal{V}. \end{array}$$

This latter map factors through the period map

$$1 \otimes dp : \mathcal{V} \otimes \mathcal{T}_B \longrightarrow \mathcal{V} \otimes \text{Sym}^2 \mathcal{V}$$

and a Koszul map

$$\mathcal{V} \otimes \text{Sym}^2 \mathcal{V} \longrightarrow \overset{2}{\wedge} \mathcal{V} \otimes \mathcal{V}.$$

Now exactness of the Koszul sequence

$$0 \rightarrow \text{Sym}^3 \mathcal{V} \rightarrow \mathcal{V} \otimes \text{Sym}^2 \mathcal{V} \rightarrow \overset{2}{\wedge} \mathcal{V} \otimes \mathcal{V}$$

shows that the desired ψ exists if and only if

$$dp \circ i = (1 \otimes dp)(i) \in \mathcal{V} \otimes \text{Sym}^2 \mathcal{V}$$

is in the subspace $\text{Sym}^3 \mathcal{V}$. (The Hamiltonian map π will automatically be Lagrangian, since i is injective.)

We conclude that, locally on B , i lifts to a 2-vector ψ satisfying conditions a), b), if and only if $dp \circ i$ is a cubic. If ψ_1, ψ_2 are two such lifts then $\psi_1 - \psi_2 \in H^0(\mathcal{X}, \overset{2}{\wedge} \pi^* \mathcal{V})$. Moreover, $\psi_1 - \psi_2$ is determined by its restriction to the zero section because $\overset{2}{\wedge} \pi^* \mathcal{V}$ restricts to a trivial bundle on each fiber. The zero section induces a splitting $\mathcal{T}_{\mathcal{X}|_{z(B)}} \simeq \pi^* \mathcal{T}_B \oplus (\pi^* \mathcal{V})|_{z(B)}$ and hence a well defined pullback $z^*(\psi) \in H^0(B, \overset{2}{\wedge} \mathcal{V})$ (locally on B). The normalizations $\psi - \pi^*(z^*(\psi))$ patch to a unique global section satisfying a), b) c).

□

The symplectic version of this lemma is:

Lemma 7.2 : (Weak cubic condition, quasi-symplectic form). A family $\pi : \mathcal{X} \rightarrow B$ of principally polarized abelian varieties has a 2-form σ satisfying

- a) $\pi : \mathcal{X} \rightarrow B$ has isotropic fibers,
 - b) σ induces a given (injective) homomorphism $j : \mathcal{T}_B \hookrightarrow \mathcal{V}^*$,
- if and only if

$$(1 \otimes j^*) \circ dp \in \text{Hom}(\mathcal{T}_B, \mathcal{T}_B^* \otimes \mathcal{V}) \cong \mathcal{T}_B^* \otimes \mathcal{T}_B^* \otimes \mathcal{V} \text{ is in } \text{Sym}^2 \mathcal{T}_B^* \otimes \mathcal{V}.$$

Moreover, in this case, there exists a unique 2-form σ satisfying a), b), and the additional condition

- c) the zero section is isotropic ($z^* \sigma = 0$).

Remark 7.3 Riemann's first bilinear condition implies further that $(1 \otimes j^*) \circ dp$ maps to $\text{Sym}^3 \mathcal{T}_B^*$, i.e., $(\text{Sym}^2 j^*) \circ dp \in \text{Hom}(\mathcal{T}_B, \text{Sym}^2 \mathcal{T}_B^*)$ comes from a cubic $c \in H^0(B, \text{Sym}^3 \mathcal{T}_B^*)$.

The cubic condition for an embedding $j : \mathcal{T}_B \hookrightarrow \mathcal{V}^*$ does not guarantee that the induced 2-form σ on \mathcal{X} is closed. In that sense, the cubic condition is a necessary condition for j to induce a symplectic structure while the following condition is necessary and sufficient (but, in general, harder to verify).

Closedness Criterion for a Symplectic Form: Given a family $\pi : \mathcal{X} \rightarrow B$ of polarized abelian varieties and a surjective $j' : \mathcal{V} \rightarrow \mathcal{T}_B^*$, there exists a closed 2-form σ on \mathcal{X} satisfying conditions a), b), c) of Lemma 7.2 if and only if $j'(\mathcal{H}_1(\mathcal{X}/B, \mathbb{Z})) \subset \mathcal{T}_B^*$ is a Lagrangian lattice in T^*B , i.e., if locally on B it consists of closed 1-forms. Moreover, the 2-form σ is uniquely determined by j' .

Proof: $j'(\mathcal{H}_1(\mathcal{X}/B, \mathbb{Z}))$ is Lagrangian \iff the canonical symplectic structure $\tilde{\sigma}$ on T^*B is translation invariant under $j'(\mathcal{H}_1(\mathcal{X}/B, \mathbb{Z})) \iff (j')^*(\tilde{\sigma})$ descends to the unique 2-form σ on $\mathcal{X} = \mathcal{V}/\mathcal{H}_1(\mathcal{X}/B, \mathbb{Z})$ satisfying conditions a), b), c) of Lemma 7.2.

□

It is instructive to relate the cubic condition to the above criterion. This is done in lemma 7.4 in a down to earth manner and is reformulated in lemma 7.5 as a coordinate free criterion.

Lemma 7.4 (“Strong Cubic Condition”) Let V be a g -dimensional vector space, $\{e_1, \dots, e_g\}$ a basis, $B \subset V^*$ an open subset, $p : B \rightarrow \mathbb{H}_g \hookrightarrow \text{Sym}^2 V$ a holomorphic map (\mathbb{H}_g is embedded in $\text{Sym}^2 V$ via the basis $\{e_j\}$), $\pi : \mathcal{X} \rightarrow B$ the corresponding family of principally polarized abelian varieties. Then the following are equivalent:

- (i) There exists a symplectic structure σ on \mathcal{X} such that $\pi : (\mathcal{X}, \sigma) \rightarrow B$ is a Lagrangian fibration and σ induces the identity isomorphism

$$\text{id} \in \text{Hom}(\mathcal{T}_{\mathcal{X}/B}, \pi^* \mathcal{T}_B^*) \simeq \text{Hom}(\pi^* \mathcal{V}, \pi^* \mathcal{V}).$$

- (ii) $p : B \rightarrow \text{Sym}^2 V$ is, locally in B , the Hessian of a function on B ,
- (iii) $dp \in \text{Hom}(\mathcal{T}_B, \text{Sym}^2 \mathcal{V}) \simeq (\mathcal{V} \otimes \text{Sym}^2 \mathcal{V})$ is a section of $\text{Sym}^3 \mathcal{V}$.

Proof: Let $\{e_j^*\}$ be the dual basis of V .

(i) \Leftrightarrow (iii): By the closedness criterion above, there exists σ as in (i) if and only if the subsheaf of lattices $\mathcal{H}^1(\mathcal{X}/B, \mathbb{Z}) \subset T^*B$ is Lagrangian, i.e., if and only if its basis

$$\{e_1, \dots, e_g, (p \rfloor e_1^*), \dots, (p \rfloor e_g^*)\}$$

consists of closed 1-forms. The e_i 's are automatically closed. If we regard the differential dp as a section of $\mathcal{T}_B^* \otimes \text{Sym}^2 \mathcal{V}$, then the two-form $d(p \rfloor e_j^*)$ is equal to the anti-symmetric part of the contraction $dp \rfloor e_j^* \in \mathcal{T}_B^* \otimes \mathcal{V} \cong \mathcal{V} \otimes \mathcal{V}$. Hence, closedness of $(p \rfloor e_j^*)$, $1 \leq j \leq g$, is equivalent to the symmetry of $dp \in \mathcal{V} \otimes \text{Sym}^2 \mathcal{V}$ also with respect to the first two factors, i.e., to dp being a section of $\text{Sym}^3 \mathcal{V}$.

(ii) \Rightarrow (iii): Clear.

(iii) \Rightarrow (ii): Follows from the Poincare lemma.

□

The additional information contained in the ‘‘Strong Cubic Condition’’ and lacking in the ‘‘Weak Cubic Condition’’ is that a Lagrangian sublattice (with respect to the polarization) $\mathcal{L} \subset \mathcal{H}_1(\mathcal{X}/B, \mathbb{Z})$ is mapped via $j' : \mathcal{V} \xrightarrow{\sim} \mathcal{T}_B^*$ to a sublattice $j'(\mathcal{L}) \subset T^*B$ Lagrangian with respect to the holomorphic symplectic structure on T^*B . (In the above lemma, $\mathcal{L} = \text{Sp}\{e_1, \dots, e_g\}$). The coordinate free reformulation of lemma 7.4 is:

Lemma 7.5 (*‘‘Strong cubic condition’’*) *Let $j' : \mathcal{V} \xrightarrow{\sim} \mathcal{T}_B^*$ be an isomorphism of the vertical bundle $\mathcal{V} = R_{\pi_*}^0(T_{\mathcal{X}/B})$ of the family $\pi : \mathcal{X} \rightarrow B$ of polarized abelian varieties with the cotangent bundle of the base. Assume only that j' maps a sublattice $\mathcal{L} \subset \mathcal{H}_1(\mathcal{X}/B, \mathbb{Z})$ Lagrangian with respect to the polarization to a sublattice $j'(\mathcal{L}) \subset T^*B$ Lagrangian with respect to the holomorphic symplectic structure on T^*B . Then there exists a symplectic structure σ on \mathcal{X} s.t. $\pi : \mathcal{X} \rightarrow B$ is a Lagrangian fibration and inducing j' if and only if j' satisfies the weak cubic condition, i.e.*

$$dp \circ i \in H^0(B, \text{Sym}^3 \mathcal{V}) \quad \text{where } i = (j')^{*-1}.$$

Remark 7.6 In most cases however, $j'(\mathcal{L})$ being Lagrangian implies $j'(\mathcal{H}_1(\mathcal{X}/B, \mathbb{Z}))$ being Lagrangian via the global monodromy action and without reference to the weak cubic condition.

Finally we remark that the above discussion applies verbatim to the case of polarized complex tori (not necessarily algebraic) since only the first Riemann bilinear condition was used.

7.3 An Integrable System for Calabi-Yau Threefolds

The Hodge group $H^{p,q}$ of an n -dimensional compact Kähler manifold X is defined as the space of harmonic forms on X of type (p, q) , i.e. involving p holomorphic and q antiholomorphic differentials. Equivalently, $H^{p,q}$ is isomorphic to the q -th cohomology

$H^q(X, \Omega^p)$ of the sheaf of holomorphic p -forms on X . The *Hodge theorem* gives a natural decomposition of the complex cohomology,

$$H^k(X, \mathbb{C}) \approx \bigoplus_{p+q=k} H^{p,q} \approx \bigoplus_{p+q=k} H^q(\Omega^p).$$

The *Hodge number* $h^{p,q}$ is the complex dimension of $H^{p,q}$. The *Hodge filtration* of $H^k(X, \mathbb{C})$ is defined by

$$F^i H^k(X, \mathbb{C}) := \bigoplus_{\substack{p+q=k \\ p \geq i}} H^{p,q}.$$

The k -th *intermediate Jacobian* of X [CG] is:

$$\begin{aligned} J^k(X) &:= H^{2k-1}(X, \mathbb{C}) / (F^k H^{2k-1}(X, \mathbb{C}) + H^{2k-1}(X, \mathbb{Z})) \\ &\approx (F^{n-k+1} H^{2n-2k+1}(X, \mathbb{C}))^* / H_{2n-2k+1}(X, \mathbb{Z}). \end{aligned}$$

Elementary properties of the Hodge filtration imply that this is a complex torus, but generally not an abelian variety unless $k = 1$ or $k = n$: it satisfies Riemann's first bilinear condition (which expresses the skew symmetry of the cup product on H^{2k-1}), but not the second, since the sign of the product (on primitive pieces) will vary with the parity of p . The extreme cases correspond to the connected component of the Picard ($k = 1$) and Albanese ($k = n$) varieties.

The Hodge decomposition does not depend holomorphically on parameters, since both holomorphic and antiholomorphic differentials are involved. The advantage of the Hodge filtration is that it does vary holomorphically and even algebraically when X is algebraic. The F^p can be defined algebraically, as the hypercohomology of the complex

$$0 \rightarrow \Omega^p \rightarrow \Omega^{p+1} \rightarrow \dots \rightarrow \Omega^n \rightarrow 0. \quad (28)$$

In particular, the intermediate Jacobian $J^k(X)$ varies holomorphically with X . This means that a smooth analytic family $\mathcal{X} \rightarrow B$ of compact Kähler manifolds gives rise to analytic vector bundles $F^i \mathcal{H}^k(\mathcal{X}/B)$ and to smooth analytic families $\mathcal{J}^k(\mathcal{X}/B) \rightarrow B$ of intermediate Jacobians of the fibers.

The bundle $\mathcal{H}^k(\mathcal{X}/B)$ is the complexification of a bundle $\mathcal{H}^k(\mathcal{X}/B, \mathbb{Z})$ of discrete groups. In particular, it has a natural local trivialization. In other words, it admits a natural flat connection, called the *Gauss-Manin* connection. The holomorphic subbundles $F^i \mathcal{H}^k(\mathcal{X}/B)$ are in general not invariant with respect to this connection, since the Hodge decomposition and filtration do change from point to point. *Griffiths' transversality* says that when a holomorphic section of $F^i \mathcal{H}^k$ is differentiated, it can move at most one step:

$$\nabla(F^i \mathcal{H}^k) \subset F^{i-1} \mathcal{H}^k \otimes \Omega_B^1.$$

An n -dimensional compact Kähler manifold X is called *Calabi-Yau* if it has trivial canonical bundle,

$$\omega_X = \Omega_X^n \approx \mathcal{O}_X,$$

and satisfies

$$h^{p,0} = 0 \text{ for } 0 < p < n.$$

A *gauged Calabi-Yau* is a pair (X, s) consisting of a Calabi-Yau manifold X together with a non-zero volume form

$$s : \mathcal{O}_X \xrightarrow{\approx} \omega_X.$$

A theorem of Bogomolov, Tian and Todorov [Bo, Ti, To] says that X has a smooth (local analytic) universal deformation space M_X . We say that a family $\chi : \mathcal{X} \rightarrow \mathcal{M}$ of Calabi-Yaus $X_t, t \in \mathcal{M}$, is *complete* if the local classifying map $\mathcal{M} \supset U_t \rightarrow M_{X_t}$ is an isomorphism for some neighborhood of every point $t \in \mathcal{M}$. It follows that \mathcal{M} is smooth and that the tangent space at t to \mathcal{M} is naturally isomorphic to $H^1(X, \mathcal{T}_X)$. Typically, such families might consist of all Calabi-Yaus in some open subset of moduli, together with some “level” structure.

The choice of gauge s gives an isomorphism

$$\lrcorner s : \mathcal{T}_X \longrightarrow \Omega_X^{n-1},$$

hence an isomorphism

$$T_X \mathcal{M} \approx H^{n-1,1}(X).$$

Starting with a complete family $\chi : \mathcal{X} \rightarrow \mathcal{M}$, we can construct

- The bundle $\mathcal{J}^k \rightarrow \mathcal{M}$ of intermediate Jacobians of the Calabi-Yau fibers.
- The space $\tilde{\mathcal{M}}$ of gauged Calabi-Yaus, a \mathbb{C}^* -bundle over \mathcal{M} obtained by removing the 0-section from the line bundle $\chi_*(\omega_{\mathcal{X}/\mathcal{M}})$.
- The fiber product

$$\tilde{\mathcal{J}}^k := \mathcal{J}^k \times_{\mathcal{M}} \tilde{\mathcal{M}},$$

which is an analytic family of complex tori $\pi : \tilde{\mathcal{J}}^k \rightarrow \tilde{\mathcal{M}}$.

Theorem 7.7 *Let $\mathcal{X} \rightarrow \mathcal{M}$ be a complete family of Calabi-Yau manifolds of odd dimension $n = 2k - 1 \geq 3$. Then there exists a canonical closed holomorphic 2-form σ on the relative k -th intermediate Jacobian $\pi : \tilde{\mathcal{J}} \rightarrow \tilde{\mathcal{M}}$ with respect to which π has maximal isotropic fibers. When $n = 3$, the 2-form σ is a symplectic structure and $\pi : \tilde{\mathcal{J}} \rightarrow \tilde{\mathcal{M}}$ is an analytically completely integrable Hamiltonian system.*

Proof.

Step I. There is a canonical isomorphism

$$T_{(X,s)} \tilde{\mathcal{M}} \approx F^{n-1} H^n(X, \mathbb{C}).$$

Indeed, the natural map $p : \tilde{\mathcal{M}} \rightarrow \mathcal{M}$ gives a short exact sequence

$$0 \rightarrow T_{(X,s)}(\tilde{\mathcal{M}}/\mathcal{M}) \rightarrow T_{(X,s)} \tilde{\mathcal{M}} \rightarrow T_X \mathcal{M} \rightarrow 0,$$

in which the subspace can be naturally identified with $H^0(\omega_X) = H^0(\Omega_X^n)$, and the quotient with $H^1(\mathcal{T}_X)$, which goes isomorphically to $H^1(\Omega_X^{n-1})$ by $\lrcorner s$. What we are claiming is that this sequence can be naturally identified with the one defining $F^{n-1} H^n$:

$$0 \rightarrow H^0(\omega_X) \rightarrow F^{n-1} \rightarrow H^1(\Omega_X^{n-1}) \rightarrow 0,$$

i.e., that the extension data match, globally over $\tilde{\mathcal{M}}$. To see this we need a natural map $T_{(X,x)} \tilde{\mathcal{M}} \rightarrow F^{n-1} H^n$ inducing the identity on the sub and quotient spaces.

Over $\tilde{\mathcal{M}}$ there is a tautological section s of $F^n\mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}},\mathbb{C})$. The Gauss-Manin connection defines an embedding

$$\nabla_{(\cdot)}s : T_{(X,s)}\tilde{\mathcal{M}} \longrightarrow H^n(X,\mathbb{C}).$$

Griffiths' transversality implies that the image is in $F^{n-1}H^n(X,\mathbb{C})$. Clearly $\nabla_{(\cdot)}s$ has the required properties.

We will need also a description of the isomorphism in terms of Dolbeault cohomology. We think of a 1-parameter family $(X_t, s_t) \in \tilde{\mathcal{M}}$, depending on the parameter t , as living on a fixed topological model X on which there are families $\bar{\partial}_t$ of complex structures (given by their $\bar{\partial}$ -operator) and s_t of C^∞ n -forms, such that s_t is of type $(n, 0)$ with respect to $\bar{\partial}_t$, all t . Since the s_t are now on a fixed underlying X , we can differentiate with respect to t :

$$s_t = s_0 + ta \quad (\text{mod } t^2).$$

Griffiths transversality now says that a is in $F^{n-1}H^n(X_0)$. It clearly depends only on the tangent vector to $\tilde{\mathcal{M}}$ along (X_t, s_t) at $t = 0$, so we get a map $T_{(X,s)}\tilde{\mathcal{M}} \longrightarrow F^{n-1}H^n$ with the desired properties.

Step II. Let \mathcal{V} be the vertical bundle on $\tilde{\mathcal{M}}$ coming from $\pi : \tilde{\mathcal{J}} \rightarrow \tilde{\mathcal{M}}$. It is isomorphic to

$$F^k\mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}})^*$$

(recall $n = 2k - 1$). Combining with Step I, we get a natural injection

$$j : T_{\tilde{\mathcal{M}}} \hookrightarrow \mathcal{V}^*,$$

which above a given (X, s) is the inclusion of $F^{n-1}H^n(X)$ into $F^kH^n(X)$. Its transpose

$$j' : \mathcal{V} \rightarrow T^*\tilde{\mathcal{M}}$$

determines a closed 2-form $\sigma := (j')^*\tilde{\sigma}$ on \mathcal{V} , where $\tilde{\sigma}$ is the standard symplectic form on $T^*\tilde{\mathcal{M}}$ (see example 2.1). By construction, the fibers of \mathcal{V} over $\tilde{\mathcal{M}}$ are maximal isotropic with respect to this form.

Step III. We need to verify that $\tilde{\sigma}$ descends to

$$\mathcal{J}^k(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}) = \mathcal{V}/\mathcal{H}_n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}},\mathbb{Z}).$$

Equivalently, a locally constant integral cycle

$$\gamma \in \Gamma(B, \mathcal{H}_n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}},\mathbb{Z})),$$

defined over some open subset B of $\tilde{\mathcal{M}}$, gives a section of \mathcal{V} on B ; hence through j' , a 1-form ξ on B , and we need this 1-form to be closed. Explicitly, if a is a section of $T_{\tilde{\mathcal{M}}}$ over B , we have

$$a \rfloor s \in \Gamma(B, F^{n-1}\mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}})) \subset \Gamma(B, \mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}))$$

and ξ is defined by:

$$\langle \xi, a \rangle := \int_\gamma (a \rfloor s).$$

Consider the function

$$g : B \longrightarrow \mathbb{C}$$

$$g(X, s) := \int_{\gamma} s.$$

If we set

$$a = \left. \frac{\partial}{\partial t} \right|_{t=0} (X_t, s_t)$$

as in Step I, we get:

$$\langle dg, a \rangle = \left. \frac{\partial}{\partial t} \right|_{t=0} g(X_t, s_t) = \left. \frac{\partial}{\partial t} \right|_{t=0} \int_{\gamma} s_t = \int_{\gamma} (a \rfloor s) = \langle \xi, a \rangle,$$

so $\xi = dg$ is closed. □

Remark 7.8 (1) The most interesting case is clearly $n = 3$, when $\tilde{\mathcal{J}}$ has an honest symplectic structure. The cubic field on $\tilde{\mathcal{M}}$ corresponding to this structure by lemma 7.1 made its first appearance in [BG] and is essentially the *Yukawa coupling*, popular among physicists and mirror-symmetry enthusiasts. At $(X, x) \in \tilde{\mathcal{M}}$ there is a natural cubic form on $H^1(\mathcal{T}_x)$:

$$c : \otimes^3 H^1(\mathcal{T}_X) \rightarrow H^3(\wedge^3 \mathcal{T}_X) = H^3(\omega_X^{-1}) \xrightarrow{\cdot s^2} H^3(\omega_X) \xrightarrow{\int} \mathbb{C},$$

which pulls back to the required cubic on $\mathcal{T}_{(X,s)}\tilde{\mathcal{M}}$. Hodge theoretically, this cubic can be interpreted as the third iterate of the infinitesimal variation of the periods, or the Hodge structure, of X c.f. [IVHS] and [BG]. By Griffiths transversality, each tangent direction on \mathcal{M} , $\theta \in H^1(\mathcal{T}_X)$, determines a linear map

$$\theta_i : H^{i,3-i} \longrightarrow H^{i-1,4-i} \quad i = 3, 2, 1,$$

and clearly the composition

$$\theta_1 \circ \theta_2 \circ \theta_3 : H^{3,0} \longrightarrow H^{0,3}$$

becomes $c(\theta)$ when we use s to identify $H^{3,0}$ and its dual $H^{0,3}$ with \mathbb{C} .

- (2) For $n = 2k - 1 \geq 5$, we get a closed 2-form on $\tilde{\mathcal{J}}$ which is in general not of maximal rank. The corresponding cubic is identically 0. Hodge theoretically, the ‘‘cubic’’ multiplies the gauge $s \in H^0(\omega_X)$ by two elements of $H^1(\mathcal{T}_X)$ (landing in $H^{n-2,2}$) and then with an element of $F^k H^n$. When $k > 2$ there are too many dz' s, so the product vanishes.
- (3) The symplectic form σ which we constructed on $\tilde{\mathcal{J}}$ is actually exact. Recall that the natural symplectic form $\tilde{\sigma}$ on $T^*\tilde{\mathcal{M}}$ is exact: $\tilde{\sigma} = d\tilde{\alpha}$, where $\tilde{\alpha}$ is the action 1-form. We obtained σ by pulling $\tilde{\sigma}$ back to $(j')^*\tilde{\sigma}$ on \mathcal{V} , and observing that the latter is invariant under translation by locally constant integral cycles γ , hence descends to

$\tilde{\mathcal{J}}$. Now a first guess for the anti-differential of σ would be the 1-form $(j')^*\tilde{\alpha}$; but this is *not* invariant under translation: if the cycle γ corresponds, as in Step III of the proof, to a 1-form ξ on $\tilde{\mathcal{M}}$, then the translation by γ changes $(j')^*\tilde{\alpha}$ by $\pi^*\xi$, where $\pi : \mathcal{V} \rightarrow \tilde{\mathcal{M}}$ is the projection. To fix this discrepancy, we consider the tautological function $f \in \Gamma(\mathcal{O}_{\mathcal{V}})$ whose value at a point $(X, s, v) \in \mathcal{V}$ (where $(X, s) \in \tilde{\mathcal{M}}$ and $v \in F^k H^n(X)^*$) is given by

$$f(X, s, v) = v(s). \quad (29)$$

This f is linear on the fibers of π , so df is constant on these fibers, and therefore translation by γ changes df by π^* of a 1-form on the base $\tilde{\mathcal{M}}$. This 1-form is clearly ξ , so we conclude that

$$(j')^*\tilde{\alpha} - df \quad (30)$$

is a global 1-form on \mathcal{V} which is invariant under translation by each γ , hence descends to a 1-form α on $\tilde{\mathcal{J}}$. It satisfies $d\alpha = \sigma$, as claimed.

- (4) Another way to see the exactness of σ on $\tilde{\mathcal{J}}$ is to note that it comes from a *quasi-contact structure* κ on \mathcal{J} . By a quasi-contact structure we mean a line subbundle κ of $T^*\mathcal{J}$. It determines a tautological 1-form on the \mathbb{C}^* -bundle $\tilde{\mathcal{J}}$ obtain from κ by omitting its zero section. Hence, it determines also an exact 2-form σ on $\tilde{\mathcal{J}}$. We refer to the pair $(\tilde{\mathcal{J}}, \sigma)$ as the *quasi-symplectification* of (\mathcal{J}, κ) . Conversely, according to [AG], page 78, a 2-form σ on a manifold $\tilde{\mathcal{J}}$ with a \mathbb{C}^* -action ρ is the quasi-symplectification of a line subbundle of the cotangent bundle of the quotient \mathcal{J} if and only if σ is homogeneous of degree 1 with respect to ρ (and the contraction of σ with the vector field generating ρ is nowhere vanishing).

In our case, there are two independent \mathbb{C}^* -actions on the total space of $T^*\tilde{\mathcal{M}} \simeq [F^{n-1}\mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}, \mathbb{C})]^*$: the \mathbb{C}^* -action on $\tilde{\mathcal{M}}$ lifts to an action $\bar{\rho}'$ on $T^*\tilde{\mathcal{M}}$, and there is also the action $\bar{\rho}''$ which commutes with the projection to $\tilde{\mathcal{M}}$ and is linear on the fibers. The symplectic form $\tilde{\sigma}$ is homogeneous of weight 0 with respect to $\bar{\rho}'$ and of weight 1 with respect to $\bar{\rho}''$, hence of weight 1 with respect to $\bar{\rho} := \bar{\rho}' \cdot \bar{\rho}''$. Hence, $\tilde{\sigma}$ is the symplectification of a contact structure on $T^*\tilde{\mathcal{M}}/\bar{\rho} \simeq [F^{n-1}\mathcal{H}^n(\mathcal{X}/\mathcal{M}, \mathbb{C})]^*$ (suppressing the gauge).

Denote a point in $[F^{n-1}\mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}, \mathbb{C})]^*$ by (X, s, ξ) . The actions, for $t \in \mathbb{C}^*$, are given by:

$$\begin{aligned} \bar{\rho}' & : (X, s, \xi) \longmapsto (X, ts, t^{-1}\xi) \\ \bar{\rho}'' & : (X, s, \xi) \longmapsto (X, s, t\xi) \\ \bar{\rho} & : (X, s, \xi) \longmapsto (X, ts, \xi). \end{aligned}$$

The function f on \mathcal{V} , given by (29), is the pullback $(j')^*(\bar{f})$ of the function \bar{f} on $T^*\tilde{\mathcal{M}}$ given by

$$\bar{f}(X, s, \xi) = \xi(s).$$

The symplectic structure $\tilde{\sigma}$ on $T^*\tilde{\mathcal{M}}$ takes the vector fields generating the actions $\bar{\rho}', \bar{\rho}''$, and $\bar{\rho}$ to the 1-forms $-d\bar{f}, \tilde{\alpha}$ and $\tilde{\alpha} - d\bar{f}$, respectively. The 1-form $\tilde{\alpha} - d\bar{f}$, which is homogeneous of degree 1 with respect to $\bar{\rho}$, is the 1-form canonically associated to the contact structure on $T^*\tilde{\mathcal{M}}/\bar{\rho} \simeq [F^{n-1}\mathcal{H}^n(\mathcal{X}/\mathcal{M}, \mathbb{C})]^*$ (namely, the contraction of $\tilde{\sigma}$ with the vector field of $\bar{\rho}$.)

Similarly, we have three action ρ', ρ'' and $\rho = \rho' \cdot \rho''$ on the total space of $\mathcal{V} \simeq [F^k \mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}, \mathbb{C})]^*$. The surjective homomorphism $j' : \mathcal{V} \rightarrow T^* \tilde{\mathcal{M}}$ is $(\rho', \bar{\rho}')$, $(\rho'', \bar{\rho}'')$, and $(\rho, \bar{\rho})$ -equivariant. The 1-form $\tilde{\alpha} - d\tilde{f}$ pulls back to the 1-form $(j')^*(\tilde{\alpha}) - d\tilde{f}$ given by (30). Clearly, the action ρ commutes with translations by $\mathcal{H}_n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}, \mathbb{Z})$. Since the 2-form $(j')^* \tilde{\sigma}$ is also $\mathcal{H}_n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}, \mathbb{Z})$ -equivariant, $(j')^*(\alpha) - d\tilde{f}$ descends to a 1-form α on $\tilde{\mathcal{J}}$. Clearly, $d\alpha = \sigma$ and α is homogeneous of degree 1 with respect to the \mathbb{C}^* -action on $\tilde{\mathcal{J}}$. Hence α comes from a quasi-contact structure κ on \mathcal{J} .

The Abel-Jacobi map of a curve to its Jacobian has an analogue for intermediate Jacobians. Let Z be a codimensional- k cycle in X , i.e. a formal linear combination $Z = \sum m_i Z_i$, with integer coefficients, of codimension k subvarieties $Z_i \subset X$. If Z is homologous to 0, we can associate to it a point $\mu(Z) \in \mathcal{J}^k(X)$, as follows. Choose a real $(2n - 2k + 1)$ -chain Γ in X whose boundary is Z , and let $\mu(Z)$ be the image in

$$\mathcal{J}^k(X) \simeq (F^{n-k+1} H^{2n-2k+1}(X, \mathbb{C}))^* / H_{2n-2k+1}(X, \mathbb{Z})$$

of the linear functional

$$\int_{\Gamma} \in (F^{n-k+1} H^{2n-2k+1}(X, \mathbb{C}))^*$$

sending a cohomology class represented by a harmonic form α to $\int_{\Gamma} \alpha$. Changing the choice of Γ changes \int_{Γ} by an integral class, so $\mu(Z)$ depends only on Z . This construction depends continuously on its parameters: given a family $\pi : \mathcal{X} \rightarrow B$ and a family $\mathcal{Z} \rightarrow B$ of codimension- k cycles in the fibers which are homologous to 0 in the fibers, we get the *normal function*, or Abel-Jacobi map

$$\mu : B \longrightarrow \mathcal{J}^k(\mathcal{X}/B)$$

to the family of intermediate Jacobians of the fibers.

Abstractly, a normal function $\nu : B \rightarrow \mathcal{J}^k(\mathcal{X}/B)$ is a section satisfying the infinitesimal condition:

Any lift

$$\tilde{\nu} : B \longrightarrow \mathcal{H}^n(\mathcal{X}/B, \mathbb{C})$$

of

$$\nu : B \rightarrow \mathcal{J}^k(\mathcal{X}/B) \simeq \mathcal{H}^n(\mathcal{X}/B, \mathbb{C}) / [F^k \mathcal{H}^n(\mathcal{X}/B, \mathbb{C}) + \mathcal{H}^n(\mathcal{X}/B, \mathbb{Z})]$$

satisfies

$$\nabla \tilde{\nu} \in F^{k-1} \mathcal{H}^n(\mathcal{X}/B, \mathbb{C}) \otimes \Omega_B^1 \tag{31}$$

or equivalently

$$(\nabla \tilde{\nu}, s) = 0 \text{ for any section } s \text{ of } F^{k+1} \mathcal{H}^n(\mathcal{X}/B, \mathbb{C})$$

where $\nabla \tilde{\nu}$ is the Gauss-Manin derivative of $\tilde{\nu}$.

This condition is independent of the choice of the lift $\tilde{\nu}$ by Griffiths' transversality. It is satisfied by the Abel-Jacobi image of a relative codimension k -cycle (see [Gr]). More generally, we can consider maps

$$\begin{array}{ccc} B & \xrightarrow{\mu} & \mathcal{J}^k \\ & \searrow q & \swarrow \\ & \mathcal{M} & \end{array}$$

The pullback $\mathcal{J}^k(\mathcal{X}/B) \rightarrow B$ of the relative intermediate Jacobian to B has a canonical section $\nu : B \rightarrow \mathcal{J}^k(\mathcal{X}/B)$. We will refer to the subvariety $\mu(B)$ as a *multivalued normal function* if $\nu : B \rightarrow \mathcal{J}^k(\mathcal{X}/B)$ is a normal function.

Theorem 7.9 *Let $\tilde{\mathcal{X}} \rightarrow \tilde{\mathcal{M}}$ be a complete family of gauged Calabi-Yau manifolds of dimension $n = 2k - 1 \geq 3$, $\tilde{\mathcal{J}} \rightarrow \tilde{\mathcal{M}}$ the relative intermediate Jacobian, $B \xrightarrow{q} \tilde{\mathcal{M}}$ a base of a family $\mathcal{Z} \rightarrow B$ of codimension- k cycles homologous to 0 in the fibers of $q^*\tilde{\mathcal{X}} \rightarrow B$. Then i) the Abel-Jacobi image in $\tilde{\mathcal{J}}$ of B is isotropic with respect to the quasi-symplectic form σ of theorem 7.7. ii) Moreover, the Abel-Jacobi image is also integral with respect to the 1-form α given by (30).*

Proof. i) We follow Step III of the proof of theorem 7.7. We thus think, locally in B , of X as being a fixed C^∞ manifold with variable complex structure $\bar{\partial}_b$, n -form s_b , and cycle Z_b , subject to the obvious compatibility. We choose a family Γ_b , $b \in B$ of n -chains whose boundary is Z_b , and consider the 1-form ξ on B given at $b \in B$ by \int_{Γ_b} ; we need to show that ξ is closed. (The new feature here is that instead of the cycles $\gamma_b \in H_n(X_b, \mathbb{Z})$ we have chains, or relative cycles $\Gamma_b \in H_n(X_b, |Z_b|, \mathbb{Z})$, where $|Z_b|$ is the support of Z_b , which varies with b .)

As before, we consider the function

$$g : B \longrightarrow \mathbb{C}$$

$$g(X, s, Z, \Gamma) := \int_{\Gamma} s,$$

and we claim $\xi = dg$. This time, in the integral $\int_{\Gamma_b} s_b$, both the integrand and the chain depend on b . So if we take a normal vector v to the supports $|Z_b|$ along Γ_b , we obtain two terms:

$$\frac{\partial}{\partial b} \int_{\Gamma_b} s_b = \int_{\Gamma_b} \frac{\partial s}{\partial b} + \int_{\partial \Gamma_b} (v \lrcorner s_b).$$

In the second term, however, s_b is of type $(n, 0)$ with respect to the complex structure $\bar{\partial}_b$, so the contraction $v \lrcorner s_b$ is of type $(n - 1, 0)$ regardless of the type of v . Since $\partial \Gamma_b = Z_b$ is of the type $(k - 1, k - 1)$, the second term vanishes identically, so we have $dg = \xi$ as desired.

ii) Integration $\int_{\Gamma}(\cdot)$ defines a section of $\mathcal{V} \simeq [F^k \mathcal{H}^n]^*$. The function g on B is the pullback via $\int_{\Gamma}(\cdot)$ of the function f on \mathcal{V} given by the formula (29). Similarly, integration $\int_{\Gamma}(\cdot)$ defines the section ξ of $T^*\tilde{\mathcal{M}} \simeq [F^{n-1} \mathcal{H}^n]^*$. The pullback of the tautological 1-form $\tilde{\alpha}$ by ξ is ξ itself. The equation $\xi - dg = 0$ translates to the statement that the 1-form $(j')^* \tilde{\alpha} - df$ vanishes on the section $\int_{\Gamma}(\cdot)$ of \mathcal{V} (see formula (30)). In particular, its descent α vanishes on the Abel-Jacobi image of $\mathcal{Z} \rightarrow B$.

□

Again, the most interesting case is $n = 3$. When B dominates the moduli space $\tilde{\mathcal{M}}$, i.e. for a multivalued choice of cycles on the general gauged Calabi-Yau of a given type, the normal function produces a Lagrangian subvariety of the symplectic $\tilde{\mathcal{J}}$, generically transversal to the fibers of the completely integrable system.

Remark 7.10 (1) The result of Theorem 7.9 holds for every multi-valued normal function $\mu : B \rightarrow \tilde{\mathcal{J}}^k(\tilde{\mathcal{X}}/\tilde{\mathcal{M}})$, not only for those coming from cycles. Given a vector field $\frac{\partial}{\partial b}$ on B , a lift $\tilde{\nu} : B \rightarrow \mathcal{H}^n$, and any section s of $F^{k+1}\mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}, \mathbb{C})$, the infinitesimal condition for normal functions (31) becomes

$$0 = \left(\nabla_{\frac{\partial}{\partial b}} \tilde{\nu}, s \right) = \frac{\partial}{\partial b}(\tilde{\nu}, s) - \left(\tilde{\nu}, \nabla_{\frac{\partial}{\partial b}} s \right). \quad (32)$$

When s is the tautological gauge, $\frac{\partial}{\partial b}(\tilde{\nu}, s)$ is the pullback of df by the projection of $\tilde{\nu}$ to $\mathcal{V} \cong \mathcal{H}^n/F^k\mathcal{H}^n$ (where f is defined by the equation (29)). Similarly, $\left(\tilde{\nu}, \nabla_{\frac{\partial}{\partial b}} s \right)$ is the contraction $\xi \rfloor \frac{\partial}{\partial b}$ of the pullback ξ of the tautological 1-form $\tilde{\alpha}$ on $T^*\tilde{\mathcal{M}}$ by the composition

$$\tilde{\mu} : B \rightarrow \mathcal{H}^n(\tilde{\mathcal{X}}/\tilde{\mathcal{M}}) \rightarrow \mathcal{H}^n/F^2 \simeq [F^{n-1}\mathcal{H}^n]^* \simeq T^*\tilde{\mathcal{M}}.$$

Thus, the infinitesimal condition for a normal function (32) implies that the image $\mu(B) \subset \tilde{\mathcal{J}}^k(\tilde{\mathcal{X}}/\tilde{\mathcal{M}})$ is integral with respect to the 1-form α (defined in (30)).

In the case of CY 3-folds ($n = 3, k = 2$) we see that the Legendre subvarieties of $\mathcal{J}^2 \rightarrow \mathcal{M}$ (i.e., the κ -integral subvarieties of maximal dimension $h^{2,1}$ where κ is the contact structure of Remark 7.8(4)) are precisely the multivalued normal functions.

- (2) Both the infinitesimal condition for a normal function (31) and the (quasi) contact structure κ on the relative Jacobian $\mathcal{J}^k \rightarrow \mathcal{M}$ (see Remark 7.8(4)) are special cases of a more general filtration of Pfaffian exterior differential systems on the relative intermediate Jacobian $\mathcal{J}^k \rightarrow \mathcal{M}$ of any family $\mathcal{X} \rightarrow \mathcal{M}$ of $n = 2k - 1$ dimensional projective algebraic varieties.

The tangent bundle $T\mathcal{J}^k$ has a canonical decreasing filtration (defined by (34) below)

$$T\mathcal{J}^k = F^0T\mathcal{J}^k \supset F^1T\mathcal{J}^k \supset \dots \supset F^{k-1}T\mathcal{J}^k \supset 0.$$

The quotient $T\mathcal{J}^k/F^iT\mathcal{J}^k$ is canonically isomorphic to the pullback of the Hodge bundle $\mathcal{H}^n/F^i\mathcal{H}^n$. The $F^{k-1}T\mathcal{J}^k$ integral subvarieties are precisely the multi-valued normal functions.

When \mathcal{J}^k is the relative intermediate Jacobian of a family of CY n -folds, the subbundle $F^1T\mathcal{J}^k$ is a hyperplane distribution on \mathcal{J}^k which defines the (quasi) contact structure κ of Remark 7.8(4).

When $n = 3, k = 2$, the filtration is a two step filtration

$$T\mathcal{J}^2 \supset F^1T\mathcal{J}^2 \supset 0$$

and the $F^1T\mathcal{J}^2$ -integral subvarieties are precisely the normal functions.

The filtration $F^iT\mathcal{J}^k$, $0 \leq i \leq k - 1$ is defined at a point $(b, y) \in \mathcal{J}^k$ over $b \in \mathcal{M}$ as follows: Choose a section $\tilde{\nu} : \mathcal{M} \rightarrow \mathcal{J}^k$ through (b, y) with the property that any lift $\tilde{\nu} : \mathcal{M} \rightarrow \mathcal{X}^n(\mathcal{H}/\mathcal{M}, \mathbb{C})$ of ν satisfies the horizontality condition

$$\nabla \tilde{\nu} \in F^i\mathcal{H}^n(\mathcal{X}/\mathcal{M}, \mathbb{C}) \otimes \Omega_{\mathcal{M}}^1 \quad (33)$$

The section ν defines a splitting

$$T_{(b,y)}\mathcal{J}^k = T_b\mathcal{M} \oplus [H^n(X_b, \mathbb{C})/F^k H^n(X_b, \mathbb{C})]$$

and the i -th piece of the filtration is defined by

$$F^i T_{(b,y)}\mathcal{J}^k := T_b\mathcal{M} \oplus [F^i H^n(X_b, \mathbb{C})/F^k H^n(X_b, \mathbb{C})]. \quad (34)$$

The horizontality condition (33) implies that the subspace $F^i T_{(b,y)}\mathcal{J}^k$ is independent of the choice of the section ν through (b, y) . Moreover, the subbundle $F^i T_{(b,y)}\mathcal{J}^k$ is invariant under translations by its integral sections, namely, by sections $\nu : \mathcal{M} \rightarrow \mathcal{J}^k$ satisfying the i -th horizontality condition.

We noted above that when $k = 1$ the intermediate Jacobian $\mathcal{J}^k(X)$ becomes the connected component $\text{Pic}^0(X)$ of the Picard variety. The generalization of the Picard variety itself is the *Deligne cohomology* group $D^k(X)$, cf. [EZ]. This fits in an exact sequence

$$0 \rightarrow \mathcal{J}^k(X) \rightarrow D^k(X) \xrightarrow{p} H^{k,k}(X, \mathbb{Z}) \rightarrow 0, \quad (35)$$

where the quotient is the group of Hodge (k, k) -classes,

$$H^{k,k}(X, \mathbb{Z}) := H^{k,k}(X, \mathbb{C}) \cap H^{2k}(X, \mathbb{Z}).$$

Any codimension- k cycle Z in X has an Abel-Jacobi image, or cycle class $\mu(Z)$ in $D^k(X)$. Its image $p(\mu(Z))$ is the cycle class of Z in ordinary cohomology.

Formally, $D^k(X)$ is defined as the hypercohomology \mathbb{H}^{2k} of the following complex of sheaves on X starting in degree 0.

$$0 \rightarrow \mathbb{Z} \rightarrow \mathcal{O}_X \rightarrow \Omega_X^1 \rightarrow \cdots \rightarrow \Omega_X^{k-1} \rightarrow 0.$$

The forgetful map to \mathbb{Z} is a map of complexes, with kernel the complex

$$0 \rightarrow \mathcal{O}_X \rightarrow \Omega_X^1 \rightarrow \cdots \rightarrow \Omega_X^{k-1} \rightarrow 0.$$

The resulting long exact sequence of hyper cohomologies gives (35).

Let $H_{alg}^{k,k}$ be the subgroup of $H^{k,k}(X, \mathbb{Z})$ of classes of algebraic cycles. (The Hodge conjecture asserts that $H_{alg}^{k,k}$ is of finite index in $H^{k,k}(X, \mathbb{Z})$.) The inverse image

$$D_{alg}^k(X) := p^{-1}(H_{alg}^{k,k})$$

has an elementary description: it is the quotient of

$$\mathcal{J}^k(X) \times \{\text{codimension-}k \text{ algebraic cycles}\}$$

by the subgroup of codimension- k cycles homologous to 0, embedded naturally in the second component and mapped to the first by Abel-Jacobi.

As X varies in a family, the rank of $H^{k,k}(X, \mathbb{Z})$ can jump up (at those X for which the variable vector subspace $H^{k,k}(X, \mathbb{C})$ happens to be in special position with respect to the “fixed” lattice $H^{2k}(X, \mathbb{Z})$). To obtain a well-behaved family of Deligne cohomology groups, we require that

$$H^{k,k}(X, \mathbb{C}) = H^{2k}(X, \mathbb{C}).$$

For example, this holds for $k = 1$ or $k = n - 1$ if $h^{2,0} = 0$. In this case we also have $H_{alg}^{k,k} = H^{k,k}(X, \mathbb{Z})$ and hence $D_{alg}^k(X) = D^k(X)$, by the Lefschetz theorem on $(1, 1)$ -classes [GH].

Corollary 7.11 *Let $\mathcal{X} \rightarrow \mathcal{M}$ be a complete family of 3-dimensional Calabi-Yau manifolds, $\tilde{\mathcal{X}} \rightarrow \tilde{\mathcal{M}}$ the corresponding gauged family. Let $\mathcal{D} \rightarrow \mathcal{M}$, $\tilde{\mathcal{D}} \rightarrow \tilde{\mathcal{M}}$, be their families of (second) Deligne cohomology groups, \mathcal{J} , $\tilde{\mathcal{J}}$ their relative intermediate Jacobians. Then there is a natural contact structure κ on \mathcal{D} with symplectification $\sigma = d\alpha$ on $\tilde{\mathcal{D}}$ with the following properties:*

- (a) σ , α , and κ restrict to the previously constructed structures on $\tilde{\mathcal{J}}$ and \mathcal{J} .
- (b) The fibration $\tilde{\mathcal{D}} \rightarrow \tilde{\mathcal{M}}$ is Lagrangian.
- (c) The multivalued normal functions of \mathcal{D} (resp. $\tilde{\mathcal{D}}$) are precisely the κ -integral (resp. α -integral) subvarieties. In particular, all multi-valued normal functions in $\tilde{\mathcal{D}}$ are isotropic.

Proof: The contact structure κ on \mathcal{J} defines one on $\mathcal{J} \times \{\text{cycles}\}$, which descends to \mathcal{D} since the equivalence relation is κ -integral by remark 7.10.

□

The mirror conjecture of conformal field theory predicts that to a family $\mathcal{X} \rightarrow \mathcal{M}$ of Calabi-Yau three folds, with some extra data, corresponds a “mirror” family $\mathcal{X}' \rightarrow \mathcal{M}'$, cf. [Mor] for the details. A first property of the conjectural symmetry is that for $X \in \mathcal{M}$, $X' \in \mathcal{M}'$,

$$h^{2,1}(X) = h^{1,1}(X'), h^{1,1} = h^{2,1}(X').$$

The conjecture goes much deeper, predicting a relation between the Yukawa cubic of \mathcal{M} and the numbers of rational curves of various homology classes in a typical $X' \in \mathcal{M}'$. This has been used spectacularly in [CdOGP] and subsequent works, to predict those numbers on a non-singular quintic hypersurface in \mathbb{P}^4 and in a number of other families.

We wonder whether the conjecture could be reformulated and understood as a type of Fourier transform between the integrable systems on the universal Deligne cohomologies $\tilde{\mathcal{D}}$ and $\tilde{\mathcal{D}}'$ of the mirror families $\tilde{\mathcal{M}}$ and $\tilde{\mathcal{M}}'$. Note that the dimensions $h^{2,1}$ and $h^{1,1}$ which are supposed to be interchanged by the mirror, can be read off the continuous and discrete parts of the fibers of $\pi : \tilde{\mathcal{D}} \rightarrow \tilde{\mathcal{M}}$, respectively. One may try to imagine the mirror as a transform, taking these Lagrangian fibers over $\tilde{\mathcal{M}}$ (which encode the Yukawa cubic, as in Section 7.2) to Lagrangian sections over $\tilde{\mathcal{M}}'$, which should somehow encode the numbers of curves in X' via their Abel-Jacobi images.

8 The Lagrangian Hilbert scheme and its relative Picard

8.1 Introduction

The Lagrangian Hilbert scheme of a symplectic variety X parametrizes Lagrangian subvarieties of X . Its relative Picard parametrizes pairs (Z, L) consisting of a line bundle L on a Lagrangian subvariety Z . We use the cubic condition of chapter 7 to construct an integrable system structure on components of the relative Picard bundle over the Lagrangian Hilbert scheme.

We interpret the generalized Hitchin integrable system, supported by the moduli space of Higgs pairs over an algebraic curve (see Ch V), as a special case of this construction. Other examples discussed include:

- a) Higgs pairs over higher dimensional base varieties (example 8.5.1), and
- b) Fano varieties of lines on hyperplane sections of a cubic fourfold (example 8.5.2).

Understanding the *global* geometry of such an integrable system requires a compactification and a study of its boundary. Our compactifications of the relative Picard are moduli spaces of sheaves and we study the symplectic structure at (smooth, stable) points of the boundary.

Let X be a smooth projective symplectic algebraic variety, σ an everywhere non degenerate algebraic 2-form on X . A smooth projective Lagrangian subvariety Z_0 of X determines a component \bar{B} of the Hilbert scheme parametrizing deformations of Z_0 in X . The component \bar{B} consists entirely of Lagrangian subschemes. Its dense open subset B , parametrizing smooth deformations of Z_0 , is a smooth quasi-projective variety [Ra, V].

Choose a very ample line bundle $\mathcal{O}_X(1)$ on X and a Hilbert polynomial p . The relative Picard $h : \mathcal{M}^p \rightarrow B$, parametrizing line bundles with Hilbert polynomial p which are supported on Lagrangian subvarieties of X , is a quasi-projective variety (see [Sim1]). If the Chern class $c_1(L_0) \in H^2(Z_0, \mathbb{Z})$ of a line bundle on Z_0 deforms as a $(1, 1)$ -class over the whole of B , then L_0 belongs to a component \mathcal{M} of \mathcal{M}^p which *dominates* the Hilbert scheme B . (By Griffiths' and Deligne's Theorem of the Fixed Part, [Sch] Corollary 7.23, this is the case for example, if $c_1(L_0)$ belongs to the image of $H^2(X, \mathbb{Q})$). Such components \mathcal{M} are integrable systems, in other words:

Theorem 8.1 *There exists a canonical symplectic structure $\sigma_{\mathcal{M}}$ on the relative Picard bundle $\mathcal{M} \xrightarrow{h} B$ over the open subset B of the Hilbert scheme of smooth projective Lagrangian subvarieties of X . The support map $h : \mathcal{M} \rightarrow B$ is a Lagrangian fibration.*

The relative Picard over the Hilbert scheme of curves on a $K3$ or abelian surface is an example [Mu1]. In example 8.5.2, X is a symplectic fourfold.

Remark 8.2 Theorem 8.1 holds in a more general setting where X is a smooth projective algebraic variety, σ is a meromorphic, generically non degenerate closed 2-form on X . We let D_0 denote its degeneracy divisor, D_∞ its polar divisor, and set $D = D_0 \cup D_\infty$. Let Z_0 be a smooth projective Lagrangian subvariety of X which does not intersect D . Denote by B

the open subset of a component of the Hilbert scheme parametrizing smooth deformations of Z_0 which stay in $X - D$. Then B is smooth and Theorem 1 holds. A special case is when X has a generically non-degenerate Poisson structure ψ . In this case D_∞ , the polar divisor of the inverse symplectic structure, is just the degeneracy locus of ψ , while D_0 is empty. The case where the subvariety Z_0 does intersect the degeneracy locus D_∞ of the Poisson structure is also of interest. It is discussed below under the category of Poisson integrable systems.

The moduli space of 1-form valued Higgs pairs is related to the case where $X = \mathbb{P}(\Omega_Y^1 \oplus \mathcal{O}_Y)$ is the compactification of the cotangent bundle of a smooth projective algebraic variety Y , and $D = \mathbb{P}\Omega_Y^1$ is the divisor at infinity (see example 8.5.1).

The relative Picard bundle \mathcal{M} is in fact also a Zariski open subset of a component of the moduli space of stable coherent sheaves on X (see [Sim1] for the construction of the moduli space). Viewed in this way, Theorem 8.1 extends a result of Mukai [Mu1] for sheaves on a $K3$ or abelian surface.

Theorem [Mu1]: *Any component of the moduli space of simple sheaves on X is smooth and has a canonical symplectic structure.*

Kobayashi [Ko] generalized the above theorem to the case of simple vector bundles on a (higher dimensional) compact complex symplectic manifold (X, σ) :

The smooth part of the moduli space has a canonical symplectic structure.

In view of Theorem 8.1 and Kobayashi's result one might be tempted to speculate that every component of the moduli space of (simple) sheaves on a symplectic algebraic variety has a symplectic structure. This is *false*. In fact, some components are odd dimensional (see example 8.19).

Returning to our symplectic relative Picard \mathcal{M} , it is natural to ask whether its *compactification* is symplectic. More precisely:

- (i) *Does the symplectic structure extend to the smooth locus of the closure of the relative Picard \mathcal{M} in the moduli space of stable (Lagrangian) sheaves?*
- (ii) *Which of these components $\bar{\mathcal{M}}$ admits a smooth projective birational model which is symplectic?*

A partial answer to (i) is provided in Theorem 8.18. We provide a cohomological identification of the symplectic structure which extends as a 2-form $\sigma_{\bar{\mathcal{M}}}$ over the smooth locus of $\bar{\mathcal{M}}$. We do not know at the moment if the 2-form $\sigma_{\bar{\mathcal{M}}}$ is *non-degenerate* at every smooth point of $\bar{\mathcal{M}}$. The cohomological identification of $\sigma_{\bar{\mathcal{M}}}$ involves a surprisingly rich *polarized Hodge-like structure* on the algebra $\text{Ext}_X^*(L, L)$ of extensions of a Lagrangian line bundle L by itself as an \mathcal{O}_X -module.

Much of the above generalizes to Poisson integrable systems. Tyurin showed in [Ty1] that Mukai's theorem generalizes to Poisson surfaces:

The smooth part of any component of the moduli space of simple sheaves on a Poisson surface has a canonical Poisson structure.

When the sheaves are supported as line bundles on curves in the surface, we get an integrable system. More precisely:

Theorem 8.3 *Let (X, ψ) be a Poisson surface, D_∞ the degeneracy divisor of ψ . Let B be the Zariski open subset of a component of the Hilbert scheme of X parametrizing smooth irreducible curves on X which are not contained in D_∞ . Then*

- i) B is smooth,
- ii) the relative Picard bundle $h : \mathcal{M} \rightarrow B$ has a canonical Poisson structure $\psi_{\mathcal{M}}$,
- iii) The bundle map $h : \mathcal{M} \rightarrow B$ is a Lagrangian fibration and
- iv) The symplectic leaf foliation of \mathcal{M} is induced by the canonical morphism $B \rightarrow \text{Hilb}_{D_\infty}$ sending a curve Z to the subscheme $Z \cap D_\infty$ of D_∞ .

The generalization to higher dimensional Poisson varieties is treated here under rather restrictive conditions on the component of the Lagrangian Hilbert scheme (see condition 8.9). These restrictions will be relaxed in [Ma2].

The rest of this chapter is organized as follows: In section 8.2 we review the deformation theory of Lagrangian subvarieties. The construction of the symplectic structure is carried out in section 8.3 where we prove Theorems 8.1 and 8.3. In section 8.4 we outline the extension of the symplectic structure to the smooth locus of the moduli space of Lagrangian sheaves (Theorem 8.18). We discuss the examples of Higgs pairs and of Fano varieties of lines on cubics in section 8.5.

8.2 Lagrangian Hilbert Schemes

Let X be a smooth n -dimensional projective algebraic variety, $Z \subset X$ a codimension q subvariety and $\mathcal{O}_X(1)$ a very ample line bundle. The Hilbert polynomial p of Z is defined to be

$$p(n) := \chi(\mathcal{O}_Z(n)) := \sum (-1)^i \dim H^i(Z, \mathcal{O}_Z(n)).$$

Grothendieck proved in [Gro] that there is a projective scheme Hilb_X^p parametrizing all algebraic subschemes of X with Hilbert polynomial p and having all the expected functoriality and naturality properties.

The Zariski tangent space $T_{[Z]}\text{Hilb}_X^p$ at the point $[Z]$ parametrizing a subvariety Z is canonically identified with the space of sections $H^0(Z, N_{X/Z})$ of the normal bundle (normal sheaf if Z is singular).

The scheme Hilb_X^p may, in general, involve pathologies. In particular, it may be non-reduced. A general criterion for the smoothness of the Hilbert scheme at a point $[Z]$ parameterizing a locally complete intersection subscheme Z is provided by:

Definition 8.4 *The semi-regularity map $\pi : H^1(Z, N_{X/Z}) \rightarrow H^{q+1}(X, \Omega_X^{q-1})$ is the dual of the natural homomorphism*

$$\pi^* : H^{n-q-1}(X, \Omega_X^{n-q+1}) \rightarrow H^{n-q-1}(Z, \omega_Z \otimes N_{Z/X}^*).$$

Here $\omega_Z \simeq \bigwedge^q N_{Z/X} \otimes \omega_X$ is the dualizing sheaf of Z and the homomorphism π^* is induced by the sheaf homomorphism

$$\Omega_X^{n-q+1} \simeq \omega_X \otimes \bigwedge^{q-1} T_X \rightarrow \omega_X \otimes \bigwedge^{q-1} N_{Z/X} \cong \omega_Z \otimes N_{Z/X}^*. \quad (36)$$

Theorem (Severi-Kodaira-Spencer-Bloch [Ka]) *If the semi-regularity map π is injective, then the Hilbert scheme is smooth at $[Z]$.*

Together with a result of Ran it implies:

Corollary 8.5 *Let (X, ψ) be a Poisson surface with a degeneracy divisor D_∞ (possibly empty). Let $Z \subset X$ be a smooth irreducible curve which is not contained in D_∞ . Then the Hilbert scheme Hilb_X^p is smooth at $[Z]$.*

Proof: The Poisson structure induces an injective homomorphism $\phi : N_{Z/X}^* \hookrightarrow T_Z$. If Z intersects D_∞ non-trivially then $N_{Z/X} \simeq \omega_Z(Z \cap D_\infty)$ and hence $H^1(Z, N_{Z/X}) = (0)$ and the semi-regularity map is trivially injective.

Note that in our case $n = 2$, $q = 1$ and the dual of the semi-regularity map

$$\pi^* : H^0(X, \omega_X) \longrightarrow H^0(Z, \omega_Z \otimes N_{Z/X}^*)$$

is induced by the sheaf homomorphism

$$\omega_X \rightarrow \omega_{X|_Z} \rightarrow \omega_Z \otimes N_{Z/X}^*$$

given by (36). If $D_\infty = \emptyset$ (X is symplectic) then ω_X , $\omega_{X|_Z}$ and $\omega_Z \otimes N_{Z/X}^*$ are all trivial line bundles and hence both π^* and the semi-regularity map are isomorphisms. If $D_\infty \cap Z = \emptyset$ but $D_\infty \neq \emptyset$ then π fails to be injective but the result nevertheless holds by a theorem of Ran which we recall below (Theorem 8.7).

□

The condition that the curve Z is not contained in D_∞ is necessary as can be seen by the following counterexample due to Severi and Zappa:

Example 8.6 ([Mum1] Section 22) Let C be an elliptic curve, E a nontrivial extension $0 \rightarrow E_1 \rightarrow E \rightarrow E_2 \rightarrow 0$, $E_i \simeq \mathcal{O}_C$ and $\pi : X = \mathbb{P}(E) \rightarrow C$ the corresponding ruled surface over C . Denote by Z the section $s : C \rightarrow X$ given by the line subbundle $E_1 \subset E$. Let $\mathcal{O}_X(-1)$ be the tautological subbundle of π^*E . Then $\mathcal{O}_X(1)$ is isomorphic to the line bundle $\mathcal{O}_X(Z)$ and the canonical bundle ω_X is isomorphic to $\pi^*(\omega_C) \otimes \mathcal{O}_X(-2) \simeq \mathcal{O}_X(-2)$. $H^0(X, \wedge^2 T_X)$ is thus isomorphic to $H^0(C, \text{Sym}^2 E^*)$ which is one dimensional. It follows that X has a unique Poisson structure ψ up to a scalar factor. The divisor $D_\infty = 2Z$ is the degeneracy divisor of ψ .

Clearly, $N_{Z/X} \simeq \pi^*T_C \simeq T_Z$ and hence $H^0(Z, N_{Z/X})$ is one dimensional. On the other hand, Z has no deformations in X (its self intersection is 0 and a deformation Z' of Z will contradict the nontriviality of the extension $0 \rightarrow E_1 \rightarrow E \rightarrow E_2 \rightarrow 0$).

□

A curve Z on a symplectic surface X is automatically Lagrangian. In the higher dimensional case we replace the curve Z by a Lagrangian subvariety. Lagrangian subvarieties of symplectic varieties have two pleasant properties:

- i) *The condition of being Lagrangian is both open and closed,*
- ii) *Their deformations are unobstructed.*

More precisely, we have:

Theorem 8.7 (*Voisin [V], Ran [Ra]*) *Let X be a smooth projective algebraic variety, σ a generically non degenerate meromorphic closed 2-form, D_∞ its polar divisor, D_0 its degeneracy divisor. Assume that $Z_0 \subset X - D_\infty - D_0$ is a smooth projective Lagrangian subvariety. Then*

- (i) *The subset of the Hilbert scheme Hilb_X^p parametrizing deformations of Z_0 in $X - D_\infty$ consists entirely of Lagrangian subvarieties.*
- (ii) *The Hilbert scheme is smooth at $[Z_0]$.*

Sketch of Proof: (i) The Lagrangian condition is closed. Thus, it suffices to prove that the open subset of smooth deformations of Z_0 is Lagrangian. If $Z \subset X - D_\infty$ then $\sigma|_Z$ is a closed holomorphic 2-form and the cohomology class $[\sigma|_Z]$ in $H^{2,0}(Z)$ vanishes if and only if $\sigma|_Z$ is identically zero. Since σ induces a *flat* section of the Hodge bundle of relative cohomology with \mathbb{C} -coefficients, then $[\sigma|_Z] = 0$ is an open and closed condition.

(ii). The symplectic structure σ induces a canonical isomorphism $N_{Z/X} \simeq \Omega_Z^1$ for any Lagrangian projective smooth subvariety $Z \subset X - D_\infty - D_0$. Ran proved a criterion for unobstructedness of deformations: the T^1 -lifting property (see [Ra, Ka]). Let $S_n = \text{Spec}(\mathbb{C}[t]/t^{n+1})$. Any flat $(n+1)$ -st order infinitesimal embedded deformation $Z_{n+1} \rightarrow S_{n+1}$ of $Z_0 = Z$ restricts canonically to an n -th order deformation $Z_n \rightarrow S_n$. In our context, the T^1 -lifting property amounts to the following criterion:

Given any $(n+1)$ -st order flat embedded deformation $Z_{n+1} \rightarrow S_{n+1}$, every extension

(a) *of $Z_n \rightarrow S_n$ to a flat embedded deformation $\tilde{Z}_n \rightarrow S_n \times_{\mathbb{C}} S_1$*

lifts to an extension

(b) *of $Z_{n+1} \rightarrow S_{n+1}$ to $\tilde{Z}_{n+1} \rightarrow S_{n+1} \times_{\mathbb{C}} S_1$.*

Extensions in (a) and (b) are classified by $T^1(Z_i/S_i) \cong H^0(Z_i, \mathcal{N}_{\varphi_i/S_i})$ where $\varphi_i : Z_i \rightarrow S_i \times X$ is the canonical morphism and $\mathcal{N}_{\varphi_i/S_i}$ is the relative normal sheaf. Recall that the De Rham cohomology and its Hodge filtration can be computed using the algebraic De Rham complex (28). Consequently, the discussion of part (i) applies in the infinitesimal setting to show that T_{Z_i/S_i} is *Lagrangian* as a subbundle of the pullback $(\varphi^i)^*T_X$ with respect to the non-degenerate 2-form $(\varphi^i)^*(\sigma)$ on $(\varphi^i)^*T_X$. The relative normal sheaf is the quotient

$$0 \rightarrow T_{Z_i/S_i} \rightarrow (\varphi^i)^*T_X \rightarrow \mathcal{N}_{\varphi_i/S_i} \rightarrow 0.$$

Hence the symplectic structure induces an isomorphism $\mathcal{N}_{\varphi_i/S_i} \simeq \Omega_{Z_i/S_i}^1$. By a theorem of Deligne, $H^0(\Omega_{Z_i/S_i}^1)$, and hence also $H^0(\mathcal{N}_{\varphi_i/S_i})$, is a free \mathcal{O}_{S_i} -module [De1]. Thus, $H^0(\mathcal{N}_{\varphi_{n+1}/S_{n+1}}) \rightarrow H^0(\mathcal{N}_{\varphi_n/S_n})$ is surjective and the T^1 -lifting property holds.

□

Note that the naive analogue of the above theorem fails for Poisson varieties. In general, deformations of Lagrangian subvarieties need not stay Lagrangian. Consider for example (\mathbb{P}^{2n}, ψ) where the Poisson structure ψ is the extension of the standard (non degenerate) symplectic structure on $\mathcal{A}^{2n} \subset \mathbb{P}^{2n}$. The Lagrangian Grassmannian has positive codimension in $Gr(n+1, 2n+1)$.

8.3 The construction of the symplectic structure

The construction of the symplectic structure on the relative Picard bundle is carried out in three steps:

In Step I we reduce it to the construction of the symplectic structure on the relative Pic^0 -bundle.

In Step II we verify the cubic condition and thus construct the 2-form (or the 2-tensor in the Poisson case).

In Step III we prove the closedness of the 2-form.

Step I: Reduction to the Pic^0 -Bundle Case:

The construction of a 2-form on the relative Picard bundle $\mathcal{M} \xrightarrow{h} B$ reduces to constructing it on its zero component $\mathcal{M}^0 \xrightarrow{h} B$, namely the Pic^0 -bundle, by the following:

Proposition 8.8 *Any closed 2-form $\sigma_{\mathcal{M}^0}$ on \mathcal{M}^0 , with respect to which the zero section of \mathcal{M}^0 is Lagrangian, extends to a closed 2-form $\sigma_{\mathcal{M}}$ on the whole Picard bundle $h : \mathcal{M} \rightarrow B$. The extension $\sigma_{\mathcal{M}}$ depends canonically on $\sigma_{\mathcal{M}^0}$ and the polarization $\mathcal{O}_X(1)$ of X .*

Proof: The point is that Picard bundles are rationally split. For any polarized projective variety $(Z, \mathcal{O}_Z(1))$, we have the Lefschetz map

$$Lef : Pic Z \longrightarrow Alb Z$$

$$[D] \mapsto [D \cap [\mathcal{O}_Z(1)]^{n-1}]$$

inducing an isogeny

$$Lef^0 : Pic^0 Z \longrightarrow Alb^0 Z.$$

We can set

$$L_Z := \{s \in Pic Z \mid \exists \ell, m, \ell \neq 0, \text{ such that } \ell \cdot Lef(s) = m \cdot Lef(\mathcal{O}(1))\}.$$

This is an extension of $H_Z^{1,1}(Z)$ by the torsion subgroup L_Z^{tor} of $Pic(Z)$. In a family $\mathcal{Z} \rightarrow B$, these groups form a subsheaf \mathcal{L} of $\mathcal{M} := Pic(\mathcal{Z}/B)$, intersecting $\mathcal{M}^0 := Pic^0(\mathcal{Z}/B)$ in its torsion subsheaf \mathcal{L}^0 . In our situation, the 2-form $\sigma_{\mathcal{M}^0}$ is \mathcal{L}^0 -invariant, so it extends uniquely to an \mathcal{L} -invariant closed 2-form $\sigma_{\mathcal{M}}$ on \mathcal{M} .

□

Step II: Verification of the Cubic Condition:

In this step we construct the 2-form (or 2-vector) on the relative Picard bundle. In the next step we will prove that it is closed (respectively, a Poisson structure).

Let (X, ψ) be a smooth projective variety, ψ a generically non-degenerate holomorphic Poisson structure. Denote by D_∞ the degeneracy divisor of ψ . We will assume throughout this step that $Z \subset X$ is a smooth subvariety, $Z \cap (X - D_\infty)$ is non empty and Lagrangian, and

Condition 8.9 i) $[Z]$ is a smooth point of the Hilbert scheme, and

ii) all deformations of Z in X are Lagrangian.

As we saw in the previous section, conditions i) and ii) hold in case ψ is everywhere non-degenerate ((X, ψ^{-1}) is a symplectic projective algebraic variety), and also in case X is a surface. Such $[Z]$ vary in a smooth Zariski open subset B of the Hilbert scheme and we denote by $h : \mathcal{M} \rightarrow B$ the relative *Pic*⁰-bundle. Condition 8.9 can be relaxed considerably (see [Ma2]).

Let

$$\phi : N_{Z/X}^* \hookrightarrow T_Z \quad (37)$$

be the injective homomorphism induced by the Poisson structure ψ . Its dual $\phi^* : T_Z^* \rightarrow N_{Z/X}$ induces an injective homomorphism.

$$i : H^0(Z, T_Z^*) \hookrightarrow H^0(Z, \mathcal{N}_{Z/X}). \quad (38)$$

The vertical tangent bundle $V := h_* \mathcal{T}_{\mathcal{M}/B}$ is isomorphic to the Hodge bundle $\mathcal{H}^{0,1}(Z/B)$. The polarization induces an isomorphism $V^* \simeq \mathcal{H}^{1,0}$. We get a global injective homomorphism $i : V^* \hookrightarrow T_B$.

Proposition 8.10 *The homomorphism i is induced by a canonical 2-vector $\psi_{\mathcal{M}} \in H^0(\mathcal{M}, \overset{2}{\wedge} T_{\mathcal{M}})$ with respect to which $h : \mathcal{M} \rightarrow B$ is a Lagrangian fibration. (We do not assert yet that $\psi_{\mathcal{M}}$ is a Poisson structure).*

Proof: It suffices to show that i satisfies the (weak) cubic condition, namely, that $dp \circ i$ comes from a cubic. The derivative of the period map

$$dp : H^0(Z, N_{Z/X}) \longrightarrow \text{Hom}(H^{1,0}(Z), H^{0,1}(Z)) \simeq [H^{1,0}(Z)^*]^{\otimes 2} \quad (39)$$

is identified by the composition

$$H^0(Z, N_{Z/X}) \xrightarrow{\text{K-S}} H^1(Z, T_Z) \xrightarrow{\text{VHS}} \text{Sym}^2 H^{1,0}(Z)^*$$

where K-S is the Kodaira-Spencer map given by cup product with the extension class of $T_{X|_Z}$:

$$\tau \in \text{Ext}^1(N_{Z/X}, T_Z) \simeq H^1(Z, N_{Z/X}^* \otimes T_Z), \quad (40)$$

and the variation of Hodge structure map VHS is given by cup product and contraction

$$H^1(Z, T_Z) \otimes H^0(Z, T_Z^*) \longrightarrow H^1(Z, \mathcal{O}_Z).$$

The composition $(\text{K-S}) \circ i : H^0(Z, T_Z^*) \rightarrow H^1(Z, T_Z)$ is then given by cup product with the class $(\phi \otimes \text{id})(\tau) \in H^1(Z, T_Z \otimes T_Z)$. We will show that $(\phi \otimes \text{id})(\tau)$ is symmetric, that is, an element of $H^1(Z, \text{Sym}^2 T_Z)$. This would imply that $d\phi$, regarded as a section of $H^0(Z, N_{Z/X})^* \otimes \text{Sym}^2 H^{1,0}(Z)^* \cong H^{1,0}(Z)^* \otimes \text{Sym}^2 H^{1,0}(Z)^*$, is symmetric also with respect to the first two factors. The cubic condition will follow.

Lemma 8.11 below implies that $(\phi \otimes \text{id})(\tau)$ is in $H^1(Z, \text{Sym}^2 T_Z)$ if and only if ϕ is induced by a section ψ in $H^0(Z, \overset{2}{\wedge} T_{X|Z})$ with respect to which Z is Lagrangian (i.e., $N_{Z/X}^*$ is isotropic). This is indeed the way ϕ was defined.

□

Lemma 8.11 *Let T be an extension*

$$0 \rightarrow Z \rightarrow T \rightarrow N \rightarrow 0 \quad (41)$$

of a vector bundle N by a vector bundle Z . Then the following are equivalent for any homomorphism $\phi : N^ \rightarrow Z$.*

i) The homomorphism ϕ is induced by a section $\psi \in H^0(\overset{2}{\wedge} T)$ with respect to which N^ is isotropic.*

ii) The homomorphism $\phi_ := H^1(\phi \otimes 1) : H^1(N^* \otimes Z) \rightarrow H^1(Z \otimes Z)$ maps the extension class $\tau \in H^1(N^* \otimes Z)$ of T to a symmetric class $\phi_*(\tau) \in H^1(\text{Sym}^2 Z) \subset H^1(Z \otimes Z)$.*

Proof: We argue as in the proof of the cubic condition (lemma 7.1). The extension (41) induces an extension

$$0 \rightarrow \overset{2}{\wedge} Z \rightarrow F \rightarrow Z \otimes N \rightarrow 0,$$

where F is the subsheaf of $\overset{2}{\wedge} T$ of sections with respect to which N^* is isotropic. The homomorphism ϕ , regarded as a section of $Z \otimes N$, lifts to a section ψ of F if and only if it is in the kernel of the connecting homomorphism

$$\delta : H^0(Z \otimes N) \rightarrow H^1(\overset{2}{\wedge} Z).$$

The latter is given by a) pairing with the extension class τ

$$(\cdot)_*\tau : H^0(Z \otimes N) \rightarrow H^1(Z \otimes Z),$$

followed by b) wedge product

$$H^1(Z \otimes Z) \xrightarrow{\wedge} H^1(\overset{2}{\wedge} Z).$$

Thus, $\delta(\phi)$ vanishes if and only if $\phi_*(\tau)$ is in the kernel of \wedge , i.e., in $H^1(\text{Sym}^2 Z)$.

□

The identification of the cubic is particularly simple in the case of a curve Z on a surface X . In that case Serre's duality identifies VHS with the dual of the multiplication map

$$\mathrm{Sym}^2 H^0(Z, \omega_Z) \xrightarrow{\mathrm{VHS}^*} H^0(Z, \omega_Z^{\otimes 2}).$$

The cubic $c \in \mathrm{Sym}^3 H^0(Z, \omega_Z)^*$ is given by composing the multiplication

$$\mathrm{Sym}^3 H^0(Z, \omega_Z) \longrightarrow H^0(Z, \omega_Z^{\otimes 3})$$

with the linear functional

$$(\phi \otimes \mathrm{id})(\tau) \in H^1(Z, T_Z^{\otimes 2}) \simeq H^0(Z, \omega_Z^{\otimes 3})^*$$

corresponding to the extension class τ .

In higher dimension (say n), the cubic depends on the choice of a polarization $\alpha \in H^{1,1}(X)$:

$$\mathrm{Sym}^3 H^0(Z, \Omega_Z^1) \rightarrow H^0(Z, \mathrm{Sym}^3 \Omega_Z^1) \xrightarrow{\phi_*(\tau)} H^1(Z, \Omega_Z^1) \xrightarrow{\alpha|_Z^{n-1}} H^{n,n}(Z) \cong \mathbb{C}.$$

The choice of α is implicitly made in the proof of proposition 8.10 when we identify $H^{0,1}(Z)$ with $H^{1,0}(Z)^*$ via the Lefschetz isomorphism (see (39)).

Step III: Closedness:

In this step we prove that the canonical 2-vector $\psi_{\mathcal{M}}$ constructed in the previous step is a Poisson structure. We first prove it in the symplectic case and later indicate the modifications needed for the Poisson case (assuming condition 8.9 of the previous step). This completes the proof of Theorems 8.1 and 8.3 stated in the introduction to this chapter.

Symplectic Case:

We assume, for simplicity of exposition, that (X, σ) is a smooth projective symplectic algebraic variety. The arguments apply verbatim to the more general setup involving a smooth projective algebraic variety X , a closed generically non-degenerate meromorphic 2-form σ on X with degeneracy divisor D_0 and polar divisor D_∞ , and Lagrangian smooth projective subvarieties which do not intersect $D_0 \cup D_\infty$.

We then have a non-degenerate 2-tensor $\psi_{\mathcal{M}}$ on $h : \mathcal{M} \rightarrow B$ and hence a 2-form $\sigma_{\mathcal{M}}$. The closedness of $\sigma_{\mathcal{M}}$ follows from that of σ_X as we now show. A polarization of X induces a relative polarization on the universal Lagrangian subvariety

$$\begin{array}{ccc} \mathcal{Z} & \longrightarrow & B \times X \\ \pi \downarrow & \swarrow p_B & \searrow p_X \\ B & & X. \end{array}$$

The relative polarization induces an isogeny

$$\begin{array}{ccc} \mathcal{M} & \longrightarrow & \mathcal{A} \\ & \searrow & \swarrow \\ & B & \end{array}$$

between the relative Pic^0 -bundle and the relative Albanese $h : \mathcal{A} \rightarrow B$. Hence, a 2-form $\sigma_{\mathcal{A}}$ on \mathcal{A} . Clearly, closedness of $\sigma_{\mathcal{M}}$ is equivalent to that of $\sigma_{\mathcal{A}}$. Since the question is local, we may assume that we have a section $\xi : B \rightarrow \mathcal{Z}$. We then get for each positive integer t a relative Albanese map

$$\begin{array}{ccc} \mathcal{Z}^t & \xrightarrow{a_t} & \mathcal{A} \\ & \searrow \pi & \swarrow h \\ & & B \end{array}$$

from the fiber product over B of t copies of the universal Lagrangian subvariety $\mathcal{Z} \rightarrow B$. For a fixed subvariety Z_b and points $(z_1, \dots, z_t) \in Z_b^t$, a_t is given by integration

$$\sum_{i=1}^t \int_{\xi(b)}^{z_i} (\cdot) \pmod{H_1(Z_b, \mathbb{Z})} \in H^{1,0}(Z)^* / H_1(Z_b, \mathbb{Z}).$$

We may assume, by choosing t large enough, that a_t is surjective. Thus, closedness of $\sigma_{\mathcal{A}}$ is equivalent to closedness of $a_t^*(\sigma_{\mathcal{A}})$. The closedness of $a_t^*(\sigma_{\mathcal{A}})$ now follows from that of σ_X by lemma 8.12.

Lemma 8.12 *Let $\ell : \mathcal{Z}^t \rightarrow X^t$ be the natural morphism; σ_{X^t} the product symplectic structure on X^t . Then,*

$$a_t^*(\sigma_{\mathcal{A}}) = \ell^*(\sigma_{X^t}) - \pi^*(\xi^t)^* \ell^*(\sigma_{X^t}). \quad (42)$$

Proof: The fibers of $\pi : \mathcal{Z}^t \rightarrow B$ are isotropic with respect to the 2-forms on both sides of the equation (42). Hence, these 2-forms induce (by contraction) homomorphisms

$$f_{\mathcal{A}}, f_X : T_{Z_b^t} \longrightarrow N_{Z_b^t/Z^t}^*.$$

The section $\xi^t(B) \subset \mathcal{Z}^t$ is also isotropic with respect to the 2-forms on both sides of equation (42). Thus, equality in (42) will follow from equality of the induced homomorphisms $f_{\mathcal{A}}, f_X$. Proving the equality $f_{\mathcal{A}} = f_X$ is a straightforward, though lengthy, unwinding of cohomological identifications.

The relative normal bundle is identified as the pullback of the tangent bundle of the Hilbert scheme

$$N_{Z_b^t/Z^t} \simeq \mathcal{O}_{Z_b^t} \otimes (T_b B) \simeq \mathcal{O}_{Z_b^t} \otimes H^0(Z_b, N_{Z_b/X}).$$

We will show that the duals of both $f_{\mathcal{A}}$ and f_X

$$f_{\mathcal{A}}^*, f_X^* : \mathcal{O}_{Z_b^t} \otimes H^0(Z_b, N_{Z_b/X}) \rightarrow T_{Z_b^t}^*$$

are identified as the composition of

i) the diagonal homomorphism

$$\mathcal{O}_{Z_b^t} \otimes H^0(Z_b, N_{Z_b/X}) \xrightarrow{\Delta} \mathcal{O}_{Z_b^t} \otimes [H^0(Z_b, N_{Z_b/X})]^t \quad \text{followed by}$$

ii) the evaluation map

$$e_t : \mathcal{O}_{Z_b^t} \otimes [H^0(Z_b, N_{Z_b/X})]^t \simeq \mathcal{O}_{Z_b^t} \otimes H^0(Z_b^t, N_{Z_b^t/X^t}) \longrightarrow N_{Z_b^t/X^t} \quad \text{followed by}$$

iii) contraction with the 2-form σ_{X^t}

$$(\phi^{-1*})^t : N_{Z_b^t/X^t} \xrightarrow{\sim} T_{Z_b^t}^*$$

(ϕ is given by contraction with the Poisson structure (37)).

Identification of $f_{\mathcal{A}}$: (for simplicity assume $t=1$).

The 2-form $\sigma_{\mathcal{A}}$ is characterized as the unique 2-form with respect to which the three conditions of lemma 7.1 hold, i.e., i) $\mathcal{A} \rightarrow B$ is a Lagrangian fibration, ii) the zero section is Lagrangian, and iii) $\sigma_{\mathcal{A}}$ induces the homomorphism

$$H^0(\phi^{-1*}) = i^{-1} : H^0(Z, N_{Z/X}) \xrightarrow{\sim} H^0(Z, T_Z^*).$$

Thus, $a^*(\sigma_{\mathcal{A}})$ induces

$$f_{\mathcal{A}}^* = \left(\mathcal{O}_{Z_b} \otimes H^0(Z_b, N_{Z_b/X}) \xrightarrow{(i^{-1})} \mathcal{O}_{Z_b} \otimes H^0(Z_b, T_{Z_b}^*) \xrightarrow{da^*} T_{Z_b}^* \right)$$

and the codifferential da^* of the Albanese map is the evaluation map.

Identification of f_X : ($t = 1$)

Both 2-forms $\ell^*(\sigma_X)$ and $\ell^*(\sigma_X) - \pi^*\xi^*\ell^*(\sigma_X)$ induce the same homomorphism $f_X^* : \mathcal{O}_{Z_b} \otimes T_b B \rightarrow T_{Z_b}^*$. This homomorphism is the composition $\phi^{-1*} \circ \overline{d\ell}$, where $\overline{d\ell}$ is the homomorphism $N_{Z_b/Z} \rightarrow N_{Z_b/X}$ induced by the differential of $\ell : Z \rightarrow X$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & T_{Z_b} & \longrightarrow & (T_Z)|_{Z_b} & \longrightarrow & H^0(Z_b, N_{Z_b/X}) \otimes \mathcal{O}_{Z_b} \longrightarrow 0 \\ & & \downarrow = & & \downarrow d\ell & & \downarrow \overline{d\ell} \\ 0 & \longrightarrow & T_{Z_b} & \longrightarrow & (\ell^*TX)|_{Z_b} & \longrightarrow & N_{Z_b/X} \longrightarrow 0. \end{array}$$

Clearly $\overline{d\ell}$ is given by evaluation. This completes the proof of lemma 8.12. □

As a simple corollary of lemma 8.12 we have:

Corollary 8.13 *There exists a canonical symplectic structure $\sigma_{\mathcal{A}^t}$ on the relative Albanese of degree $t \in \mathbb{Z}$, depending canonically on the symplectic structure σ_X (independent of the polarization $\mathcal{O}_X(1)$!) and satisfying, for $t \geq 1$,*

$$a_t^*(\sigma_{\mathcal{A}^t}) = \ell^*(\sigma_{X^t}).$$

(the pullback to the fiber product $\mathcal{Z}^t := \times_B^t \mathcal{Z}$ via the Albanese map coincides with the pullback of the symplectic structure σ_{X^t} on X^t).

Proof: The $t = 0$ case is proven. We sketch the proof of the $t \geq 1$ case. The $t \leq -1$ case is similar. Let ξ be a local section of $\mathcal{Z}^t \rightarrow B$. Translation by the section $-a_t(\xi)$ of \mathcal{A}^{-t} defines a local isomorphism

$$\tau_{\xi} : \mathcal{A}^t \longrightarrow \mathcal{A}^0.$$

Let

$$\sigma_{\mathcal{A}^t} := \tau_{\xi}^*(\sigma_{\mathcal{A}^0}) + h^*\xi^*\ell^*(\sigma_{X^t}).$$

We claim that $\sigma_{\mathcal{A}^t}$ is independent of ξ . This amounts to the identity

$$\tau_{(\xi_1 - \xi_2)}^*(\sigma_{\mathcal{A}^0}) = \sigma_{\mathcal{A}^0} - h^*[a_0(\xi_1 - \xi_2)]^*\sigma_{\mathcal{A}^0}$$

for any two sections ξ_1, ξ_2 of $\mathcal{Z}^t \rightarrow B$.

□

Poisson Case: (assuming condition 8.9)

Showing that the 2-vector $\psi_{\mathcal{M}}$ constructed in step II is a Poisson structure, amounts to showing that

Lemma 8.14 $\psi_{\mathcal{M}}(T_{\mathcal{M}}^*) \subset T_{\mathcal{M}}$ is an involutive distribution,

and

Lemma 8.15 the induced 2-form on each symplectic leaf is closed.

Sketch of Proof of Lemma 8.14: Since $h : \mathcal{M} \rightarrow B$ is a Lagrangian fibration with respect to $\psi_{\mathcal{M}}$ (by proposition 8.10), the distribution is the pullback of the distribution on the base B . The latter is induced by the image of the injective homomorphism $i : V^* \hookrightarrow T_B$ identified by (38)

$$i = H^0(\phi^*) : H^0(Z, T_Z^*) \hookrightarrow H^0(Z, N_{Z/X}).$$

Recall (37) that ϕ , in turn, is induced by the Poisson structure ψ_X on X . The involutivity now follows from that of $\psi_X(T_X^*) \subset T_X$ by a deformation theoretic argument. The details are omitted.

□

In case X is a surface, the degeneracy divisor D_{∞} of ψ_X is a curve and $iH^0(Z, T_Z^*) \subset H^0(Z, N_{Z/X})$ is the subspace of all infinitesimal deformations of Z which fix the divisor $Z \cap D_{\infty}$. Thus, the distribution $i(V^*)$ on the Hilbert scheme B (as in the proof of lemma 8.14) corresponds to the foliation by level sets of the algebraic morphism

$$\begin{aligned} R : B &\longrightarrow \text{Hilb}(D_{\infty}) \\ Z &\longmapsto Z \cap D_{\infty}. \end{aligned}$$

The higher dimensional case is analogous. The degeneracy divisor D_{∞} has an algebraic rank stratification

$$D_{\infty} = \bigcup_{r=0}^{n-1} D_{\infty}[2r] \quad (\dim X = 2n).$$

Each rank stratum is foliated, local analytically, by symplectic leaves. The subspace

$$i : H^0(Z, T_Z^*) \subset H^0(Z, N_{Z/X})$$

is characterized as the subspace of all infinitesimal deformations of Z which deform the subscheme $Z \cap D_\infty[2r]$ fixing the image $f(Z \cap D_\infty[2r])$ with respect to any Casimir function f on $D_\infty[2r]$. As an illustration, consider the case where X is the logarithmic cotangent bundle $T_M^*(\log(D))$ and Z is a 1-form with logarithmic poles along a divisor D with normal crossing. In this case the residues induce the symplectic leaves foliation.

Sketch of Proof of Lemma 8.15: The proof is essentially the same as in the symplectic case. We consider an open (analytic) subset B_1 of a leaf in B , the universal Lagrangian

$$\begin{array}{ccc} \mathcal{Z}_1 & \xrightarrow{\ell} & X & & \mathcal{A} \\ \text{subvariety} & \downarrow & & , \text{ and the relative Albanese} & \downarrow & \\ & B_1 & & & B_1 & \end{array} .$$

One has to choose the section $\xi : B_1 \rightarrow \mathcal{Z}_1$ outside $\ell^{-1}(D_\infty)$ and notice that the identity (42) implies that the pullback $\ell^*(\sigma_X^t)$ of the *meromorphic* closed 2-form σ_X^t (inverse of the generically non-degenerate Poisson structure on the product of t copies of X) is a *holomorphic* 2-form on \mathcal{Z}_1^t (because $a_t^*(\sigma_{\mathcal{A}})$ is) and that $a_t^*(\sigma_{\mathcal{A}})$ is *closed* (because $\ell^*(\sigma_{X^t})$ is).

□

8.4 Partial compactifications: a symplectic structure on the moduli space of Lagrangian sheaves

We describe briefly in this section the extension of the symplectic structure on the relative Picard \mathcal{M} to an algebraic 2-form on the smooth locus of a partial compactification. For details see [Ma2]. For simplicity, we assume that (X, σ) is a smooth $2n$ -dimensional projective symplectic variety. We note that with obvious modifications, the extension of the 2-form will hold in the setup $(X, \sigma, D_0, D_\infty)$ as in remark 8.2 allowing σ to degenerate and have poles away from the support of the sheaves.

When X is a symplectic surface, some of these extensions give rise to smooth projective symplectic compactifications [Mu1]. These projective symplectic compactifications appear also in the higher dimensional case:

Example 8.16 A somewhat trivial reincarnation of a relative Picard of a linear system on a K3 surface S as a birational model of a relative Picard of a Lagrangian Hilbert scheme over a higher dimensional symplectic variety X is realized as follows. Let X be the Beauville variety $S^{[n]}$ which is the resolution of the n -th symmetric product of S provided by the Hilbert scheme of zero cycles of length n [B2]. The symmetric powers $C^{[n]}$ of smooth curves on S are smooth Lagrangian subvarieties of $S^{[n]}$. Components of the relative Picard over the smooth locus in the linear system $|C|$ are isomorphic to Zariski open subsets of components of the relative Picard over the Lagrangian Hilbert scheme of $S^{[n]}$.

This leads us to speculate that genuinely new examples of smooth symplectic *projective* varieties will arise as birational models of moduli spaces of Lagrangian line bundles. (see section 8.5.2 for new *quasiprojective* examples).

We worked so far with a component $\mathcal{M} \rightarrow B$ of the relative Picard of the universal smooth Lagrangian subvariety $\mathcal{Z} \rightarrow B$ which dominates the corresponding component \bar{B} of the Lagrangian Hilbert scheme (i.e., if L is supported on Z , $c_1(L) \in H_{\mathbb{Z}}^{1,1}(Z)$ remains

of type $(1, 1)$ over B). Let $p(n) := \chi(L \otimes_{\mathcal{O}_X} \mathcal{O}_X(n))$ be the Hilbert polynomial of a Lagrangian line bundle L parametrized by \mathcal{M} . A construction of C. Simpson enables us to compactify \mathcal{M} as an open subset of a component \mathcal{M}^{ss} of the moduli space of equivalence classes of coherent semistable sheaves on X with Hilbert polynomial p [Sim1]. Denote by \mathcal{M}^s the open subset of \mathcal{M}^{ss} parametrizing isomorphism classes of stable sheaves, $\mathcal{M}^{s,sm}$ the smooth locus of \mathcal{M}^s . Then $\mathcal{M} \subseteq \mathcal{M}^{s,sm} \subseteq \mathcal{M}^s \subseteq \mathcal{M}^{ss}$. In addition, the moduli space \mathcal{M}^s embeds as a Zariski open subset of the moduli space of simple sheaves [AK2]. The Zariski tangent space $T_{[L]}\mathcal{M}^s$ at a stable sheaf L is thus canonically isomorphic to the Zariski tangent space of the moduli space of simple sheaves. The latter is identified as the group $\text{Ext}_{\mathcal{O}_X}^1(L, L)$ of extensions $0 \rightarrow L \rightarrow E \rightarrow L \rightarrow 0$ of L by L as an \mathcal{O}_X -module. When X is a $K3$ or abelian surface, Mukai's symplectic structure is given by the pairing

$$\text{Ext}_{\mathcal{O}_X}^1(L, L) \otimes \text{Ext}_{\mathcal{O}_X}^1(L, L) \xrightarrow{\text{Yoneda}} \text{Ext}_{\mathcal{O}_X}^2(L, L) \xrightarrow{\text{S.D.}} \text{Hom}_X(L, L \otimes \omega_X)^* \xrightarrow{id \otimes \sigma} \mathbb{C}$$

(Composition of the Yoneda pairing, Serre Duality, and evaluation at

$$id \otimes \sigma \in \text{Hom}_X(L, L \otimes \omega_X)).$$

The generalization of Mukai's pairing requires the construction of a homomorphism, depending linearly on the Poisson structure ψ ,

$$y : H^{1,1}(X) \rightarrow \text{Ext}_{\mathcal{O}_X}^2(L, L). \quad (43)$$

It sends the Kahler class $\alpha := c_1(\mathcal{O}_X(1)) \in H^{1,1}(X)$ to a 2-extension class $y(\alpha) \in \text{Ext}_{\mathcal{O}_X}^2(L, L)$. Once this is achieved, the 2-form $\sigma_{\mathcal{M}}$ will become:

$$\begin{aligned} & \text{Ext}_{\mathcal{O}_X}^1(L, L) \otimes \text{Ext}_{\mathcal{O}_X}^1(L, L) \xrightarrow{\text{Yoneda}} \text{Ext}_{\mathcal{O}_X}^2(L, L) \xrightarrow{y(\alpha)^{n-1}} \text{Ext}_{\mathcal{O}_X}^{2n}(L, L) \xrightarrow{\text{S.D.}} \\ & \text{Hom}_X(L, L \otimes \omega_X)^* \xrightarrow{id \otimes \sigma^n} \mathbb{C}. \end{aligned} \quad (44)$$

Remark 8.17 When L is a line bundle on a smooth Lagrangian subvariety Z the construction involves a surprisingly rich polarized Hodge-like structure on the algebra

$$\text{Ext}_{\mathcal{O}_X}^*(L, L) := \bigoplus_{k=0}^{2n} \text{Ext}_{\mathcal{O}_X}^k(L, L).$$

Since $\text{Ext}_{\mathcal{O}_X}^k(L, \cdot)$ is the right derived functor of the composition $\Gamma \circ \mathcal{H}om_{\mathcal{O}_X}(L, \cdot)$ of the Sheaf Hom and the global sections functors, there is a spectral sequence converging to $\text{Ext}_{\mathcal{O}_X}^k(L, L)$ with

$$E_2^{p,q} = H^p(Z, \mathcal{E}xt_{\mathcal{O}_X}^q(L, L))$$

(see [HS]). The sheaf of q -extensions $\mathcal{E}xt_{\mathcal{O}_X}^q(L, L)$ is canonically isomorphic to $\bigwedge^q N_{Z/X}$ and thus, via the symplectic structure, to Ω_Z^q . We obtain a canonical isomorphism $E_2^{p,q} \simeq H^{q,p}(Z)$ with the Dolbeault groups of Z . Notice however, that the Dolbeault groups appear in *reversed order* compared to their order in the graded pieces of the Hodge filtration on the cohomology ring $H^*(Z, \mathbb{C})$.

The construction of the 2-extension class $y(\alpha)$ and hence of the generalized Mukai pairing (44) can be carried out for all coherent sheaves parametrized by $\mathcal{M}^{s,sm}$. We obtain:

Theorem 8.18 [Ma2] *The symplectic structure $\sigma_{\mathcal{M}}$ on the relative Picard \mathcal{M} extends to an algebraic 2-form over the smooth locus $\mathcal{M}^{s,sm}$ of the closure of \mathcal{M} in the moduli space of stable sheaves on X . It is identified by the pairing (44).*

The non-degeneracy of $\sigma_{\mathcal{M}}$ at a point $[L] \in \mathcal{M}$ parametrizing a line bundle on a smooth Lagrangian subvariety Z follows from the Hard Lefschetz theorem. We expect $\sigma_{\mathcal{M}}$ to be non degenerate everywhere on $\mathcal{M}^{s,sm}$.

Finally we remark that the pairing (44) can be used to define a 2-form on other components of the moduli space of stable sheaves on X . This 2-form will, in general, be degenerate. In fact, some components are odd dimensional:

Example 8.19 Consider an odd dimensional complete linear system $|Z|$ whose generic element is a smooth ample divisor on an abelian variety X of even dimension ≥ 4 , with a symplectic structure σ . The dimension of the component of the Hilbert scheme parametrizing deformations of Z is $\dim(\text{Pic } X) + \dim |Z| = \dim X + \dim |Z|$. Since $h^{1,0}(Z) = h^{1,0}(X)$, the component of the moduli space of sheaves parameterizing deformations of the structure sheaf \mathcal{O}_Z , as an \mathcal{O}_X -module, is of dimension $2 \cdot \dim X + \dim |Z|$ which is odd.

□

It is the Hodge theoretic interpretation of the graded pieces of the spectral sequence of $\text{Ext}_{\mathcal{O}_X}^k(L, L)$ for *Lagrangian* line bundles which assures the non degeneracy of $\sigma_{\mathcal{M}}$.

8.5 Examples

8.5.1 Higgs Pairs

In chapter 9 we define the notion of a 1-form valued Higgs pair (E, φ) over a smooth n -dimensional projective algebraic variety X . It consists of a torsion free sheaf E over X and a homomorphism $\varphi : E \rightarrow E \otimes \Omega_X^1$ satisfying the symmetry condition $\varphi \wedge \varphi = 0$.

The moduli space Higgs_X of semistable Higgs pairs of rank r with vanishing first and second Chern classes may be viewed as the Dolbeault non-abelian first $GL_r(\mathbb{C})$ -cohomology group of X (cf. [Sim2] and theorem 4.7 when X is a curve):

Non-abelian Hodge theory introduces a hyperkahler structure on the smooth locus of the space $\mathcal{M}_{\text{Betti}}$ of isomorphism classes of semisimple $GL_r(\mathbb{C})$ -representations of the fundamental group $\pi_1(X)$ of X [De2, H1, Sim3]. The hyperkahler structure consists of a Riemannian metric and an action of the quaternion algebra \mathbb{H} on the real tangent bundle with respect to which

- (i) the (purely imaginary) unit vectors $\{a|a\bar{a} = 1\}$ in \mathbb{H} correspond to a (holomorphic) \mathbb{P}^1 -family of integrable complex structures,
- (ii) the metric is Kahler with respect to these complex structures.

All but two of the complex structures are isomorphic to that of $\mathcal{M}_{\text{Betti}}$, the two special ones are that of Higgs_X and its conjugate ($\mathcal{M}_{\text{Betti}}$ and Higgs_X are diffeomorphic).

The hyperkahler structure introduces a holomorphic symplectic structure σ on the smooth locus of Higgs_X . In case X is a Riemann surface, that symplectic structure is the one giving rise to the Hitchin integrable system of spectral Jacobians.

Our aim is to interpret the symplectic structure on Higgs_X as an example of a Lagrangian structure over the relative Picard of a Lagrangian component of the Hilbert scheme of the cotangent bundle T_X^* of X . This interpretation will apply to the Hitchin system (where $\dim X = 1$). For higher dimensional base varieties X it will apply only to certain particularly nice cases. See also [Bi] for a deformation theoretic study of the holomorphic symplectic structure.

The spectral construction (proposition 4.5) can be carried out also for Higgs pairs over a higher dimensional smooth projective variety X (cf. [Sim1]). We have a one to one correspondence between

- (i) (Stable) Higgs pairs (E, φ) on X (allowing E to be a rank r torsion free sheaf) and
- (ii) (Stable) sheaves F on the cotangent bundle T_X^* which are supported on (pure) n -dimensional projective subschemes of T_X^* which are finite, degree r , branched coverings (in a scheme theoretic sense) of X .

Projective subvarieties of T_X^* which are finite over X are called *spectral coverings*. Spectral coverings \tilde{X} are necessarily Lagrangian since the symplectic form σ on T_X^* , which restricts to a global exact 2-form on \tilde{X} , must vanish on \tilde{X} .

Let B be the open subset of a component of the Hilbert scheme of $\mathbb{P}(T_X^* \oplus \mathcal{O}_X)$ parametrizing degree r smooth spectral coverings (closed subvarieties of $\mathbb{P}(T_X^* \oplus \mathcal{O}_X)$ which are contained in T_X^*). The above correspondence embeds components of the relative Picard $\mathcal{M} \rightarrow B$ as open subsets of components of the moduli spaces of stable rank r Higgs pairs over X . Theorem 8.1 of this chapter implies

Corollary 8.20 (i) *The open subset \mathcal{M} of the moduli Higgs_X of Higgs pairs over X which, under the spectral construction, parametrizes line bundles on smooth spectral covers, has a canonical symplectic structure $\sigma_{\mathcal{M}}$ (we do not require the Chern classes of the Higgs pairs to vanish).*

(ii) *The support morphism $h : \mathcal{M} \rightarrow B$ is a Lagrangian fibration.*

Remark 8.21 In general, when $\dim X > 1$, there could be components of the moduli spaces of Higgs pairs for which the open set \mathcal{M} above is empty, i.e.,

1. the spectral coverings of all Higgs pairs in this component are singular, or
2. the corresponding sheaves on the spectral coverings are torsion free but not locally free.

8.5.2 Fano Varieties of Lines on Cubic Fourfolds

We will use theorem 8.1 to prove:

Example 8.22 *Let Y be a smooth cubic hypersurface in \mathbb{P}^5 . The relative intermediate Jacobian $\mathcal{J} \rightarrow B$ over the family $B \subset |\mathcal{O}_{\mathbb{P}^5}(1)|$ of smooth cubic hyperplane sections of Y is an algebraically completely integrable Hamiltonian system.*

The statement follows from a description of the family $\mathcal{J} \rightarrow B$ as an open subset of the moduli space of Lagrangian sheaves on the Fano variety X of lines on Y .

A. Beauville and R. Donagi proved [BD] that X is symplectic (fourfold). Clemens and Griffiths proved in [CG] that the intermediate jacobian J_b of a smooth hyperplane section $Y \cap H_b$ is isomorphic to the Picard $Pic^0 Z_b$ of the 2-dimensional Fano variety Z_b of lines on the cubic 3-fold $Y \cap H_b$. C. Voisin observed that Z_b is a Lagrangian subvariety of X [V]. Since $h^{1,0}(Y \cap H_b) = 5$, B is isomorphic to a dense open subset of a component of the Hilbert scheme. In fact, using results of Altman and Kleiman, one can show that the corresponding component is isomorphic to $|\mathcal{O}_{\mathbb{P}^5}(1)|$ (see [AK1] Theorem 3.3 (iv)). Theorem 8.1 implies that the relative Picard $\mathcal{M} \rightarrow B$ has a completely integrable Hamiltonian system structure.

The symplectic structure $\sigma_{\mathcal{M}}$ is defined also at the fiber of the relative Picard corresponding to a Fano variety Z_b of lines on a hyperplane section $Y \cap H_b$ with an ordinary double point $x_b \in Y \cap H_b$ (Theorem 8.18). In that case, we have a genus 4 curve C_b in Z_b parametrizing lines through x_b . Z_b is isomorphic to the quotient $S^2 C_b / (C_1 \sim C_2)$ of the second symmetric product of C_b modulo the identification of two disjoint copies of C_b [CG]. It is not difficult to check that $\sigma_{\mathcal{M}}$ is *non-degenerate* also on the fiber $Pic^0(Z_b)$ of the relative Picard which is a \mathbb{C}^\times -extension of the Jacobian of genus 4. The non-degeneracy of the symplectic structure implies that we get an *induced boundary integrable system* on the relative Picard of the family of genus 4 curves

$$Pic(\mathcal{C}) \rightarrow (Y^* - \Delta) \tag{45}$$

over the complement of the singular locus Δ of the dual variety of the cubic fourfold.

It is interesting to note that the boundary integrable system (45) can not be realized as the relative Picard of a family of curves on a symplectic surface. If this were the case, the generic rank of the pullback $a^*(\sigma_{\mathcal{P}ic^1(\mathcal{C})})$ of the symplectic structure from $\mathcal{P}ic^1(\mathcal{C}) \rightarrow (Y^* - \Delta)$ to $\mathcal{C} \rightarrow (Y^* - \Delta)$ via the Abel-Jacobi map would be 2. On the other hand, $a^*(\sigma_{\mathcal{P}ic^1(\mathcal{C})})$ is equal to the pullback of the symplectic structure σ_X on X via the natural dominant map $\mathcal{C} \rightarrow X$ (corollary 8.13). Thus, its generic rank is 4. The importance of this rank as an invariant of integrable systems supported by families of Jacobians is illustrated in an interesting recent study of J. Hurtubise [Hur].

More examples of nonrigid Lagrangian subvarieties can be found in [V, Ye].

9 Spectral covers

9.1 Algebraic extensions

We have seen that Hitchin's system, the geodesic flow on an ellipsoid, the polynomial matrices system of Chapter 4, the elliptic solitons, and so on, all fit as special cases of the spectral system on a curve. In this final chapter, we consider some algebraic properties of the general spectral system. We are still considering families of Higgs pairs $(E, \varphi : E \rightarrow E \otimes K)$, but we generalize in three separate directions:

1. The base curve C is replaced by an arbitrary complex algebraic variety S . The spectral curve \tilde{C} then becomes a spectral cover $\tilde{S} \rightarrow S$.
2. The line bundle K in which the endomorphism φ takes its values is replaced by a vector bundle, which we still denote by K . (this requires an integrability condition on φ .) Equivalently, \tilde{S} is now contained in the total space \mathbb{K} of a vector bundle over S .
3. Instead of the vector bundle E we consider a principal G -bundle \mathcal{G} , for an arbitrary complex reductive group G . The G -vector bundle E is then recovered as $E := \mathcal{G} \times^G V$, given the choice of a representation $\rho : G \rightarrow \text{Aut}(V)$. The twisted endomorphism φ is replaced by a section of $K \otimes \text{ad}(\mathcal{G})$. Even in the original case of $G = GL(n)$ one encounters interesting phenomena in studying the dependence of $\tilde{S} := \tilde{S}_V$, for a given (\mathcal{G}, φ) , on the representation V of G .

We will see that essentially all *algebraic* properties (but not the *symplectic* structure) of the Hitchin system, or of the (line-bundle valued, $G = GL(n)$) spectral system on a curve, survive in this new context. In fact, the added generality forces the discovery of some symmetries which were not apparent in the original:

- Spectral curves are replaced by spectral covers. These come in several flavors: $\tilde{S}_V, \tilde{S}_\lambda, \tilde{S}_P$, indexed by representations of G , weights, and parabolic subgroups. The most basic object is clearly the *cameral* cover \tilde{S} ; all the others can be considered as associated objects. In case $G = GL(n)$, the cameral cover specializes not to our previous spectral cover, which has degree n over S , but roughly to its Galois closure, of degree $n!$ over S .
- The spectral Picards, $\text{Pic}(\tilde{S}_V)$ etc., can all be written directly in terms of the decomposition of $\text{Pic}(\tilde{S})$ into Prym-type components under the action of the Weyl group W . In particular, there is a distinguished Prym component common to all the nontrivial $\text{Pic}(\tilde{S}_V)$. The identification of this component combines and unifies many interesting constructions in Prym theory.
- The Higgs bundle too can be relieved of its excess baggage. Stripping away the representation V as well as the values bundle K , one arrives (in subsection 9.3.1) at the notion of abstract, principal Higgs bundle. The abelianization procedure assigns to this a spectral datum, consisting of a cameral cover with an equivariant bundle on it.

- There is a Hitchin map (59) which is algebraically completely integrable in the sense that its fibers can be naturally identified, up to a "shift" and a "twist", with the distinguished Pryms (Theorem 9.10).
- The "shift" is a property of the group G , and is often nonzero even when \tilde{S} is etale over S , cf. Proposition 9.8. The "twist", on the other hand, arises from the ramification of \tilde{S} over S , cf. formula (57).
- The resulting abelianization procedure is local in the base S , and does not require particularly nice behavior near the ramification, cf. example 9.11. It does require that φ be regular (this means that its centralizer has the smallest possible dimension; for $GL(n)$, this means that each eigenvalue may have arbitrary multiplicity, but the eigen-*space* must be 1-dimensional), at least over the generic point of S . At present we can only guess at the situation for irregular Higgs bundles.

The consideration of general spectral systems is motivated in part by work of Hitchin [H2] and Simpson [Sim1]. In the remainder of this section we briefly recall those works. Our exposition in the following sections closely follows that of [D4], which in turn is based on [D2], for the group-theoretic approach to spectral decomposition used in section (9.2), and on [D3] for the Abelianization procedure, or equivalence of Higgs and spectral data, in section(9.3). Some of these results, especially in the case of a base curve, can also be found in [AvM, BK, F1, K, Me, MS, Sc].

Reductive groups

Principal G -bundles \mathcal{G} for arbitrary reductive G were considered already in Hitchin's original paper [H2]. Fix a curve C and a line bundle K . There is a moduli space $\mathcal{M}_{G,K}$ parametrizing equivalence classes of semistable K -valued G -Higgs bundles, i.e. pairs (\mathcal{G}, φ) with $\varphi \in K \otimes ad(\mathcal{G})$. The Hitchin map goes to

$$B := \oplus_i H^0(K^{\otimes d_i}),$$

where the d_i are the degrees of the f_i , a basis for the G -invariant polynomials on the Lie algebra \mathfrak{g} . It is:

$$h : (\mathcal{G}, \varphi) \longrightarrow (f_i(\varphi))_i.$$

When $K = \omega_C$, Hitchin showed [H2] that one still gets a completely integrable system, and that this system is algebraically completely integrable for the classical groups $GL(n), SL(n), SP(n), SO(n)$. The generic fibers are in each case (not quite canonically; one must choose various square roots! cf. sections 9.3.2 and 9.3.3) isomorphic to abelian varieties given in terms of the spectral curves \tilde{C} :

- $GL(n)$ \tilde{C} has degree n over \mathbb{C} , the AV is $\text{Jac}(\tilde{C})$.
 $SL(n)$ \tilde{C} has degree n over \mathbb{C} , the AV is $\text{Prym}(\tilde{C}/\mathbb{C})$.
 $SP(n)$ \tilde{C} has degree $2n$ over \mathbb{C} and an involution $x \mapsto -x$.
The map factors through the quotient \overline{C} .
The AV is $\text{Prym}(\tilde{C}/\overline{C})$.
 $SO(n)$ \tilde{C} has degree n and an involution, with: (46)
 - a fixed component, when n is odd.
 - some fixed double points, when n is even.
One must desingularize \tilde{C} and the quotient \overline{C} ,
and ends up with the Prym of the
desingularized double cover.

For the exceptional group G_2 , the algebraic complete integrability was verified in [KP1]. A sketch of the argument for any reductive G is in [BK], and a complete proof was given in [F1]. We will outline a proof in section 9.3 below.

Higher dimensions

A sweeping extension of the notion of Higgs bundle is suggested by the work of Simpson [Sim1], which was already discussed in Chapter 8. To him, a Higgs bundle on a projective variety S is a vector bundle (or principal G -bundle ...) E with a *symmetric*, Ω_S^1 -valued endomorphism

$$\varphi : E \longrightarrow E \otimes \Omega_S^1.$$

Here *symmetric* means the vanishing of:

$$\varphi \wedge \varphi : E \longrightarrow E \otimes \Omega_S^2,$$

a condition which is obviously vacuous on curves. Simpson constructs a moduli space for such Higgs bundles (satisfying appropriate stability conditions), and establishes diffeomorphisms to corresponding moduli spaces of connections and of representations of $\pi_1(S)$.

In our approach, the Ω^1 -valued Higgs bundle will be considered as a particular realization of an abstract Higgs bundle, given by a subalgebra of $ad(\mathcal{G})$. The symmetry condition will be expressed in the definition 9.5 of an abstract Higgs bundle by requiring the abelian subalgebras of $ad(\mathcal{G})$ to be abelian.

9.2 Decomposition of spectral Picards

9.2.1 The question

Throughout this section we fix a vector bundle K on a complex variety S , and a pair (\mathcal{G}, φ) where \mathcal{G} is a principal G -bundle on S and φ is a regular section of $K \otimes ad(\mathcal{G})$. (This data is equivalent to the regular case of what we call in section 9.3.1 a K -valued principal Higgs bundle.) Each representation

$$\rho : G \longrightarrow \text{Aut}(V)$$

determines an associated K -valued Higgs (vector) bundle

$$(\mathcal{V} := \mathcal{G} \times^G V, \quad \rho(\varphi)),$$

which in turn determines a spectral cover $\tilde{S}_V \longrightarrow S$.

The question, raised first in [AvM] when $S = \mathbf{P}^1$, is to relate the Picard varieties of the \tilde{S}_V as V varies, and in particular to find pieces common to all of them. For Adler and van Moerbeke, the motivation was that many evolution DEs (of Lax type) can be *linearized* on the Jacobians of spectral curves. This means that the "Liouville tori", which live naturally in the complexified domain of the DE (and hence are independent of the representation V) are mapped isogenously to their image in $\text{Pic}(\tilde{S}_V)$ for each nontrivial V ; so one should be able to locate these tori among the pieces which occur in an isogeny decomposition of each of the $\text{Pic}(\tilde{S}_V)$. There are many specific examples where a pair of abelian varieties constructed from related covers of curves are known to be isomorphic or isogenous, and some of these lead to important identities among theta functions.

Example 9.1 Take $G = SL(4)$. The standard representation V gives a branched cover $\tilde{S}_V \longrightarrow S$ of degree 4. On the other hand, the 6-dimensional representation $\wedge^2 V$ (=the standard representation of the isogenous group $SO(6)$) gives a cover $\tilde{\tilde{S}} \longrightarrow S$ of degree 6, which factors through an involution:

$$\tilde{\tilde{S}} \longrightarrow \bar{S} \longrightarrow S.$$

One has the isogeny decompositions:

$$\text{Pic}(\tilde{S}) \sim \text{Prym}(\tilde{S}/S) \oplus \text{Pic}(S)$$

$$\text{Pic}(\tilde{\tilde{S}}) \sim \text{Prym}(\tilde{\tilde{S}}/\bar{S}) \oplus \text{Prym}(\bar{S}/S) \oplus \text{Pic}(S).$$

It turns out that

$$\text{Prym}(\tilde{S}/S) \sim \text{Prym}(\tilde{\tilde{S}}/\bar{S}).$$

For $S = \mathbf{P}^1$, this is Recillas' *trigonal construction* [Rec]. It says that every Jacobian of a trigonal curve is the Prym of a double cover of a tetragonal curve, and vice versa.

Example 9.2 Take $G = SO(8)$ with its standard 8-dimensional representation V . The spectral cover has degree 8 and factors through an involution, $\tilde{\tilde{S}} \longrightarrow \bar{S} \longrightarrow S$. The two half-spin representations V_1, V_2 yield similar covers

$$\tilde{\tilde{S}}_i \longrightarrow \bar{S}_i \longrightarrow S, \quad i = 1, 2.$$

The *tetragonal construction* [D1] says that the three Pryms of the double covers are isomorphic. (These examples, as well as Pantazis' *bigonal construction* and constructions based on some exceptional groups, are discussed in the context of spectral covers in [K] and [D2].)

It turns out in general that there is indeed a distinguished, Prym-like isogeny component common to all the spectral Picards, on which the solutions to Lax-type DEs evolve linearly. This was noticed in some cases already in [AvM], and was greatly extended by Kanev's construction of Prym-Tyurin varieties. (He still needs S to be \mathbf{P}^1 and the spectral cover to have generic ramification; some of his results apply only to *minuscule representations*.) Various parts of the general story have been worked out recently by a number of authors, based on either of two approaches: one, pursued in [D2, Me, MS], is to decompose everything according to the action of the Weyl group W and to look for common pieces; the other, used in [BK, D3, F1, Sc], relies on the correspondence of spectral data and Higgs bundles. The group-theoretic approach is described in the rest of this section. We take up the second method, known as *abelianization*, in section 9.3.

9.2.2 Decomposition of spectral covers

The decomposition of spectral Picards arises from three sources. First, the spectral cover for a sum of representations is the union of the individual covers \tilde{S}_V . Next, the cover \tilde{S}_V for an irreducible representation is still the union of subcovers \tilde{S}_λ indexed by weight orbits. And finally, the Picard of \tilde{S}_λ decomposes into Pryms. We start with a few observations about the dependence of the covers themselves on the representation. The decomposition of the Picards is taken up in the next subsection.

Spectral covers

Whenever a representation space V of G decomposes,

$$V = \oplus V_i,$$

there is a corresponding decomposition

$$\tilde{S}_V = \cup \tilde{S}_{V_i},$$

so we may restrict attention to irreducible representations V . There is an *infinite* collection (of irreducible representations $V := V_\mu$, hence) of spectral covers \tilde{S}_V , which can be parametrized by their highest weights μ in the dominant Weyl chamber \overline{C} , or equivalently by the W -orbit of extremal weights, in Λ/W . Here T is a maximal torus in G , $\Lambda := \text{Hom}(T, \mathbf{C}^*)$ is the *weight lattice* (also called *character lattice*) for G , and W is the Weyl group. Now V_μ decomposes as the sum of its weight subspaces V_μ^λ , indexed by certain weights λ in the convex hull in Λ of the W -orbit of μ . We conclude that each \tilde{S}_{V_μ} itself decomposes as the union of its subcovers \tilde{S}_λ , each of which involves eigenvalues in a given W -orbit $W\lambda$. (λ runs over the weight-orbits in V_μ .)

Parabolic covers

There is a *finite* collection of covers \tilde{S}_P , parametrized by the conjugacy classes in G of parabolic subgroups (or equivalently by arbitrary dimensional faces F_P of the chamber \overline{C}) such that (for general S) each eigenvalue cover \tilde{S}_λ is birational to some parabolic cover \tilde{S}_P , the one whose open face F_P contains λ .

The cameral cover

There is a W -Galois cover $\tilde{S} \rightarrow S$ such that each \tilde{S}_P is isomorphic to \tilde{S}/W_P , where W_P is the Weyl subgroup of W which stabilizes F_P . We call \tilde{S} the *cameral cover*, since, at least generically, it parametrizes the chambers determined by φ (in the duals of the Cartans). Informally, we think of $\tilde{S} \rightarrow S$ as the cover which associates to a point $s \in S$ the set of Borel subalgebras of $ad(\mathcal{G})_s$ containing $\phi(s)$. More carefully, this is constructed as follows: There is a morphism $\mathfrak{g} \rightarrow \mathfrak{t}/W$ sending $g \in \mathfrak{g}$ to the conjugacy class of its semisimple part g_{ss} . (More precisely, this is $Spec$ of the composed ring homomorphism $\mathbf{C}[\mathfrak{t}]^W \xrightarrow{\cong} \mathbf{C}[\mathfrak{g}]^G \hookrightarrow \mathbf{C}[\mathfrak{g}]$.) Taking fiber product with the quotient map $\mathfrak{t} \rightarrow \mathfrak{t}/W$, we get the cameral cover $\tilde{\mathfrak{g}}$ of \mathfrak{g} . The cameral cover $\tilde{S} \rightarrow S$ of a K -valued principal Higgs bundle on S is glued from covers of open subsets in S (on which K and \mathcal{G} are trivialized) which in turn are pullbacks by φ of $\tilde{\mathfrak{g}} \rightarrow \mathfrak{g}$.

9.2.3 Decomposition of spectral Picards

The decomposition of the Picard varieties of spectral covers can be described as follows:

The cameral Picard

From each isomorphism class of irreducible W -representations, choose an integral representative Λ_i . (This can always be done, for Weyl groups.) The group ring $\mathbf{Z}[W]$ which acts on $Pic(\tilde{S})$ has an isogeny decomposition:

$$\mathbf{Z}[W] \sim \oplus_i \Lambda_i \otimes_{\mathbf{Z}} \Lambda_i^*, \quad (47)$$

which is just the decomposition of the regular representation. There is a corresponding isotypic decomposition:

$$Pic(\tilde{S}) \sim \oplus_i \Lambda_i \otimes_{\mathbf{Z}} Prym_{\Lambda_i}(\tilde{S}), \quad (48)$$

where

$$Prym_{\Lambda_i}(\tilde{S}) := Hom_W(\Lambda_i, Pic(\tilde{S})). \quad (49)$$

Parabolic Picards

There are at least three reasonable ways of obtaining an isogeny decomposition of $Pic(\tilde{S}_P)$, for a parabolic subgroup $P \subset G$:

- The ‘Hecke’ ring $Corr_P$ of correspondences on \tilde{S}_P over S acts on $Pic(\tilde{S}_P)$, so every irreducible integral representation M of $Corr_P$ determines a generalized Prym

$$Hom_{Corr_P}(M, Pic(\tilde{S}_P)),$$

and we obtain an isotypic decomposition of $Pic(\tilde{S}_P)$ as before.

- $Pic(\tilde{S}_P)$ maps, with torsion kernel, to $Pic(\tilde{S})$, so we obtain a decomposition of the former by intersecting its image with the isotypic components $\Lambda_i \otimes_{\mathbf{Z}} Prym_{\Lambda_i}(\tilde{S})$ of the latter.

- Since \tilde{S}_P is the cover of S associated to the W -cover \tilde{S} via the permutation representation $\mathbf{Z}[W_P \backslash W]$ of W , we get an isogeny decomposition of $Pic(\tilde{S}_P)$ indexed by the irreducible representations in $\mathbf{Z}[W_P \backslash W]$.

It turns out ([D2], section 6) that all three decompositions agree and can be given explicitly as

$$\oplus_i M_i \otimes Prym_{\Lambda_i}(\tilde{S}) \subset \oplus_i \Lambda_i \otimes Prym_{\Lambda_i}(\tilde{S}), \quad M_i := (\Lambda_i)^{W_P}. \quad (50)$$

Spectral Picards

To obtain the decomposition of the Picards of the original covers \tilde{S}_V or \tilde{S}_λ , we need, in addition to the decomposition of $Pic(\tilde{S}_P)$, some information on the singularities. These can arise from two separate sources:

Accidental singularities of the \tilde{S}_λ . For a sufficiently general Higgs bundle, and for a weight λ in the interior of the face F_P of the Weyl chamber \overline{C} , the natural map:

$$i_\lambda : \tilde{S}_P \longrightarrow \tilde{S}_\lambda$$

is birational. For the *standard* representations of the classical groups of types A_n, B_n or C_n , this *is* an isomorphism. But for general λ it is *not*. In order for i_λ to be an isomorphism, λ must be a multiple of a fundamental weight, cf. [D2], lemma 4.2. In fact, the list of fundamental weights for which this happens is quite short; for the classical groups we have only: ω_1 for A_n, B_n and C_n , ω_n (the dual representation) for A_n , and ω_2 for B_2 . Note that for D_n the list is *empty*. In particular, the covers produced by the standard representation of $SO(2n)$ are singular; this fact, noticed by Hitchin in [H2], explains the need for desingularization in his result (46).

Gluing the \tilde{S}_V . In addition to the singularities of each i_λ , there are the singularities created by the gluing map $\Pi_\lambda \tilde{S}_\lambda \longrightarrow \tilde{S}_V$. This makes explicit formulas somewhat simpler in the case, studied by Kanev [K], of *minuscule* representations, i.e. representations whose weights form a single W -orbit. These singularities account, for instance, for the desingularization required in the $SO(2n+1)$ case in (46).

9.2.4 The distinguished Prym

Combining much of the above, the Adler–van Moerbeke problem of finding a component common to the $Pic(\tilde{S}_V)$ for all non-trivial V translates into:

Find the non trivial irreducible representations Λ_i of W which occur in $\mathbf{Z}[W_P \backslash W]$ with positive multiplicity for all proper Weyl subgroups $W_P \subsetneq W$.

It is easy to see that for arbitrary finite groups W , or even for Weyl groups W if we allow arbitrary rather than Weyl subgroups W_P , there may be no common factors [D2]. For example, when W is the symmetric group S_3 (=the Weyl group of $GL(3)$) and W_P is S_2 or A_3 , the representations $\mathbf{Z}[W_P \backslash W]$ are 3 or 2 dimensional, respectively, and have only the trivial representation as common component. In any case, our problem is equivalent (by Frobenius reciprocity or (50)) to

Find the irreducible representations Λ_i of W such that for every proper Weyl subgroup $W_P \subsetneq W$, the space of invariants $M_i := (\Lambda_i)^{W_P}$ is non-zero.

One solution is now obvious: the *reflection representation* of W acting on the weight lattice Λ has this property. In fact, Λ^{W_P} in this case is just the face F_P of \overline{C} . The corresponding component $\text{Prym}_\Lambda(\tilde{S})$, is called *the distinguished Prym*. We will see in section 9.3 that its points correspond, modulo some corrections, to Higgs bundles.

For the classical groups, this turns out to be the only common component. For G_2 and E_6 it turns out ([D2], section 6) that a second common component exists. The geometric significance of points in these extra components is not known. As far as we know, the only component other than the distinguished Prym which has arisen ‘in nature’ is the one associated to the 1-dimensional sign representation of W , cf. [KP2].

9.3 Abelianization

9.3.1 Abstract vs. K -valued objects

We want to describe the abelianization procedure in a somewhat abstract setting, as an equivalence between *principal Higgs bundles* and certain *spectral data*. Once we fix a *values* vector bundle K , we obtain an equivalence between *K -valued principal Higgs bundles* and *K -valued spectral data*. Similarly, the choice of a representation V of G will determine an equivalence of *K -valued Higgs bundles* (of a given representation type) with *K -valued spectral data*.

As our model of a W -cover we take the natural quotient map

$$G/T \longrightarrow G/N$$

and its partial compactification

$$\overline{G/T} \longrightarrow \overline{G/N}. \tag{51}$$

Here $T \subset G$ is a maximal torus, and N is its normalizer in G . The quotient G/N parametrizes maximal tori (=Cartan subalgebras) \mathfrak{t} in \mathfrak{g} , while G/T parametrizes pairs $\mathfrak{t} \subset \mathfrak{b}$ with $\mathfrak{b} \subset \mathfrak{g}$ a Borel subalgebra. An element $x \in \mathfrak{g}$ is *regular* if the dimension of its centralizer $\mathfrak{c} \subset \mathfrak{g}$ equals $\dim T$ (=the rank of \mathfrak{g}). The partial compactifications $\overline{G/N}$ and $\overline{G/T}$ parametrize regular centralizers \mathfrak{c} and pairs $\mathfrak{c} \subset \mathfrak{b}$, respectively.

In constructing the cameral cover in section 9.2.2, we used the W -cover $\mathfrak{t} \longrightarrow \mathfrak{t}/W$ and its pullback cover $\tilde{\mathfrak{g}} \longrightarrow \mathfrak{g}$. Over the open subset \mathfrak{g}_{reg} of regular elements, the same cover is obtained by pulling back (51) via the map $\alpha : \mathfrak{g}_{reg} \longrightarrow \overline{G/N}$ sending an element to its centralizer:

$$\begin{array}{ccccc} \mathfrak{t} & \longleftarrow & \tilde{\mathfrak{g}}_{reg} & \longrightarrow & \overline{G/T} \\ \downarrow & & \downarrow & & \downarrow \\ \mathfrak{t}/W & \longleftarrow & \mathfrak{g}_{reg} & \xrightarrow{\alpha} & \overline{G/N} \end{array} . \tag{52}$$

When working with K -valued objects, it is usually more convenient to work with the left hand side of (52), i.e. with *eigenvalues*. When working with the abstract objects, this is unavailable, so we are forced to work with the *eigenvectors*, or the right hand side of (52). Thus:

Definition 9.3 An abstract cameral cover of S is a finite morphism $\tilde{S} \rightarrow S$ with W -action, which locally (etale) in S is a pullback of (51).

Definition 9.4 A K -valued cameral cover (K is a vector bundle on S) consists of a cameral cover $\pi : \tilde{S} \rightarrow S$ together with an S -morphism

$$\tilde{S} \times \Lambda \rightarrow \mathbb{K} \tag{53}$$

which is W -invariant (W acts on \tilde{S}, Λ , hence diagonally on $\tilde{S} \times \Lambda$) and linear in Λ .

We note that a cameral cover \tilde{S} determines quotients \tilde{S}_P for parabolic subgroups $P \subset G$. A K -valued cameral cover determines additionally the \tilde{S}_λ for $\lambda \in \Lambda$, as images in \mathbb{K} of $\tilde{S} \times \{\lambda\}$. The data of (53) is equivalent to a W -equivariant map $\tilde{S} \rightarrow \mathfrak{t} \otimes_{\mathbb{C}} K$.

Definition 9.5 A G -principal Higgs bundle on S is a pair (\mathcal{G}, c) with \mathcal{G} a principal G -bundle and $c \subset ad(\mathcal{G})$ a subbundle of regular centralizers.

Definition 9.6 A K -valued G -principal Higgs bundle consists of (\mathcal{G}, c) as above together with a section φ of $c \otimes K$.

A principal Higgs bundle (\mathcal{G}, c) determines a cameral cover $\tilde{S} \rightarrow S$ and a homomorphism $\Lambda \rightarrow \text{Pic}(\tilde{S})$. Let F be a parameter space for Higgs bundles with a given \tilde{S} . Each non-zero $\lambda \in \Lambda$ gives a non-trivial map $F \rightarrow \text{Pic}(\tilde{S})$. For λ in a face F_P of \bar{C} , this factors through $\text{Pic}(\tilde{S}_P)$. The discussion in section 9.2.4 now suggests that F itself should be given roughly by the distinguished Prym,

$$\text{Hom}_W(\Lambda, \text{Pic}(\tilde{S})).$$

It turns out that this guess needs two corrections. The first correction involves restricting to a coset of a subgroup; the need for this is visible even in the simplest case where \tilde{S} is etale over S , so (\mathcal{G}, c) is everywhere regular and semisimple (i.e. c is a bundle of Cartans.) The second correction involves a twist along the ramification of \tilde{S} over S . We explain these in the next two subsections.

9.3.2 The regular semisimple case: the shift

Example 9.7 Fix a smooth projective curve C and a line bundle $K \in \text{Pic}(C)$ such that $K^{\otimes 2} \approx \mathcal{O}_C$. This determines an etale double cover $\pi : \tilde{C} \rightarrow C$ with involution i , and homomorphisms

$$\begin{aligned} \pi^* & : \text{Pic}(C) \rightarrow \text{Pic}(\tilde{C}) \quad , \\ \text{Nm} & : \text{Pic}(\tilde{C}) \rightarrow \text{Pic}(C) \quad , \\ i^* & : \text{Pic}(\tilde{C}) \rightarrow \text{Pic}(\tilde{C}) \quad , \end{aligned}$$

satisfying

$$1 + i^* = \pi^* \circ \text{Nm}.$$

- For $G = GL(2)$ we have $\Lambda = \mathbf{Z} \oplus \mathbf{Z}$, and $W = \mathcal{S}_2$ permutes the summands, so

$$\text{Hom}_W(\Lambda, \text{Pic}(\tilde{C})) \approx \text{Pic}(\tilde{C}).$$

And indeed, the Higgs bundles corresponding to \tilde{C} are parametrized by $\text{Pic}(\tilde{C})$: send $L \in \text{Pic}(\tilde{C})$ to $(\mathcal{G}, \mathfrak{c})$, where \mathcal{G} has associated rank-2 vector bundle $\mathcal{V} := \pi_* L$, and $\mathfrak{c} \subset \underline{\text{End}}(\mathcal{V})$ is $\pi_* \mathcal{O}_{\tilde{C}}$.

- On the other hand, for $G = SL(2)$ we have $\Lambda = \mathbf{Z}$ and $W = \mathcal{S}_2$ acts by ± 1 , so

$$\text{Hom}_W(\Lambda, \text{Pic}(\tilde{S})) \approx \{L \in \text{Pic}(\tilde{C}) \mid i^* L \approx L^{-1}\} = \ker(1 + i^*).$$

This group has 4 connected components. The subgroup $\ker(\text{Nm})$ consists of 2 of these. The connected component of 0 is the classical Prym variety, cf. [Mum3]. Now the Higgs bundles correspond, via the above bijection $L \mapsto \pi_* L$, to

$$\{L \in \text{Pic}(\tilde{C}) \mid \det(\pi_* L) \approx \mathcal{O}_C\} = \text{Nm}^{-1}(K).$$

Thus they form the *non-zero* coset of the subgroup $\ker(\text{Nm})$. (If we return to a higher dimensional S , there is no change in the $GL(2)$ story, but it is possible for K not to be in the image of Nm , so there may be *no* $SL(2)$ -Higgs bundles corresponding to such a cover.)

This example generalizes to all G , as follows. The equivalence classes of extensions

$$1 \longrightarrow T \longrightarrow N' \longrightarrow W \longrightarrow 1$$

(in which the action of W on T is the standard one) are parametrized by the group cohomology $H^2(W, T)$. Here the 0 element corresponds to the semidirect product. The class $[N] \in H^2(W, T)$ of the normalizer N of T in G may be 0, as it is for $G = GL(n), \mathbf{PGL}(n), SL(2n+1)$; or not, as for $G = SL(2n)$.

Assume first, for simplicity, that S, \tilde{S} are connected and projective. There is then a natural group homomorphism

$$c : \text{Hom}_W(\Lambda, \text{Pic}(\tilde{S})) \longrightarrow H^2(W, T). \quad (54)$$

Algebraically, this is an edge homomorphism for the Grothendieck spectral sequence of equivariant cohomology, which gives the exact sequence

$$0 \longrightarrow H^1(W, T) \longrightarrow H^1(S, \mathcal{C}) \longrightarrow \text{Hom}_W(\Lambda, \text{Pic}(\tilde{S})) \xrightarrow{c} H^2(W, T). \quad (55)$$

where $\mathcal{C} := \tilde{S} \times_W T$. Geometrically, this expresses a *Mumford group* construction: giving $\mathcal{L} \in \text{Hom}(\Lambda, \text{Pic}(\tilde{S}))$ is equivalent to giving a principal T -bundle \mathcal{T} over \tilde{S} ; for $\mathcal{L} \in \text{Hom}_W(\Lambda, \text{Pic}(\tilde{S}))$, $c(\mathcal{L})$ is the class in $H^2(W, T)$ of the group N' of automorphisms of \mathcal{T} which commute with the action on \tilde{S} of some $w \in W$.

To remove the restriction on S, \tilde{S} , we need to replace each occurrence of T in (54,55) by $\Gamma(\tilde{S}, T)$, the global sections of the trivial bundle on \tilde{S} with fiber T . The natural map $H^2(W, T) \longrightarrow H^2(W, \Gamma(\tilde{S}, T))$ allows us to think of $[N]$ as an element of $H^2(W, \Gamma(\tilde{S}, T))$.

Proposition 9.8 [D3] Fix an étale W -cover $\pi : \tilde{S} \rightarrow S$. The following data are equivalent:

1. Principal G -Higgs bundles $(\mathcal{G}, \mathfrak{c})$ with cameral cover \tilde{S} .
2. Principal N -bundles \mathcal{N} over S whose quotient by T is \tilde{S} .
3. W -equivariant homomorphisms $\mathcal{L} : \Lambda \rightarrow \text{Pic}(\tilde{S})$ with $c(\mathcal{L}) = [N] \in H^2(W, \Gamma(\tilde{S}, T))$.

We observe that while the shifted objects correspond to Higgs bundles, the unshifted objects

$$\mathcal{L} \in \text{Hom}_W(\Lambda, \text{Pic}(\tilde{S})), \quad c(\mathcal{L}) = 0$$

come from the \mathcal{C} -torsors in $H^1(S, \mathcal{C})$.

9.3.3 The regular case: the twist along the ramification

Example 9.9 Modify example 9.7 by letting $K \in \text{Pic}(C)$ be arbitrary, and choose a section b of $K^{\otimes 2}$ which vanishes on a simple divisor $B \subset C$. We get a double cover $\pi : \tilde{C} \rightarrow C$ branched along B , ramified along a divisor

$$R \subset \tilde{C}, \quad \pi(R) = B.$$

Via $L \mapsto \pi_* L$, the $SL(2)$ -Higgs bundles still correspond to

$$\{L \in \text{Pic}(\tilde{C}) \mid \det(\pi_* L) \approx \mathcal{O}_C\} = \text{Nm}^{-1}(K).$$

But this is no longer in $\text{Hom}_W(\Lambda, \text{Pic}(\tilde{S}))$; rather, the line bundles in question satisfy

$$i^* L \approx L^{-1}(R). \tag{56}$$

For arbitrary G , let Φ denote the root system and Φ^+ the set of positive roots. There is a decomposition

$$\overline{G/T} \setminus G/T = \bigcup_{\alpha \in \Phi^+} R_\alpha$$

of the boundary into components, with R_α the fixed locus of the reflection σ_α in α . (Via (52), these correspond to the complexified walls in \mathfrak{t} .) Thus each cameral cover $\tilde{S} \rightarrow S$ comes with a natural set of (Cartier) *ramification divisors*, which we still denote R_α , $\alpha \in \Phi^+$.

For $w \in W$, set

$$F_w := \{\alpha \in \Phi^+ \mid w^{-1}\alpha \in \Phi^-\} = \Phi^+ \cap w\Phi^-,$$

and choose a W -invariant form \langle, \rangle on Λ . We consider the variety

$$\text{Hom}_{W,R}(\Lambda, \text{Pic}(\tilde{S}))$$

of R -twisted W -equivariant homomorphisms, i.e. homomorphisms \mathcal{L} satisfying

$$w^*\mathcal{L}(\lambda) \approx \mathcal{L}(w\lambda) \left(\sum_{\alpha \in F_w} \frac{\langle -2\alpha, w\lambda \rangle}{\langle \alpha, \alpha \rangle} R_\alpha \right), \quad \lambda \in \Lambda, \quad w \in W. \quad (57)$$

This turns out to be the correct analogue of (56). (E.g. for a reflection $w = \sigma_\alpha$, F_w is $\{\alpha\}$, so this gives $w^*\mathcal{L}(\lambda) \approx \mathcal{L}(w\lambda) \left(\frac{\langle \alpha, 2\lambda \rangle}{\langle \alpha, \alpha \rangle} R_\alpha \right)$, which specializes to (56).) As before, there is a class map

$$c : \text{Hom}_{W,R}(\Lambda, \text{Pic}(\tilde{S})) \longrightarrow H^2(W, \Gamma(\tilde{S}, T)) \quad (58)$$

which can be described via a Mumford-group construction.

To understand this twist, consider the formal object

$$\begin{aligned} \frac{1}{2}\text{Ram} : \Lambda &\longrightarrow \mathbf{Q} \otimes \text{Pic}\tilde{S}, \\ \lambda &\longmapsto \sum_{(\alpha \in \Phi^+)} \frac{\langle \alpha, \lambda \rangle}{\langle \alpha, \alpha \rangle} R_\alpha. \end{aligned}$$

In an obvious sense, a principal T -bundle \mathcal{T} on \tilde{S} (or a homomorphism $\mathcal{L} : \Lambda \longrightarrow \text{Pic}(\tilde{S})$) is R -twisted W -equivariant if and only if $\mathcal{T}(-\frac{1}{2}\text{Ram})$ is W -equivariant, i.e. if \mathcal{T} and $\frac{1}{2}\text{Ram}$ transform the same way under W . The problem with this is that $\frac{1}{2}\text{Ram}$ itself does not make sense as a T -bundle, because the coefficients $\frac{\langle \alpha, \lambda \rangle}{\langle \alpha, \alpha \rangle}$ are not integers. (This argument shows that if $\text{Hom}_{W,R}(\Lambda, \text{Pic}(\tilde{S}))$ is non-empty, it is a torser over the untwisted $\text{Hom}_W(\Lambda, \text{Pic}(\tilde{S}))$.)

Theorem 9.10 [D3] *For a cameral cover $\tilde{S} \longrightarrow S$, the following data are equivalent:*

- (1) G -principal Higgs bundles with cameral cover \tilde{S} .
- (2) R -twisted W -equivariant homomorphisms $\mathcal{L} \in c^{-1}([N])$.

The theorem has an essentially local nature, as there is no requirement that S be, say, projective. We also do not need the condition of generic behavior near the ramification, which appears in [F1, Me, Sc]. Thus we may consider an extreme case, where \tilde{S} is ‘everywhere ramified’:

Example 9.11 In example 9.9, take the section $b = 0$. The resulting cover \tilde{C} is a ‘ribbon’, or length-2 non-reduced structure on C : it is the length-2 neighborhood of C in \mathbb{K} . The $\text{SL}(2)$ -Higgs bundles $(\mathcal{G}, \mathfrak{c})$ for this \tilde{C} have an everywhere nilpotent \mathfrak{c} , so the vector bundle $\mathcal{V} := \mathcal{G} \times^{SL(2)} V \approx \pi_* L$ (where V is the standard 2-dimensional representation) fits in an exact sequence

$$0 \longrightarrow \mathcal{S} \longrightarrow \mathcal{V} \longrightarrow \mathcal{Q} \longrightarrow 0$$

with $\mathcal{S} \otimes K \approx \mathcal{Q}$. Such data are specified by the line bundle \mathcal{Q} , satisfying $\mathcal{Q}^{\otimes 2} \approx K$, and an extension class in $\text{Ext}^1(\mathcal{Q}, \mathcal{S}) \approx H^1(K^{-1})$. The kernel of the restriction map $\text{Pic}(\tilde{C}) \longrightarrow \text{Pic}(C)$ is also given by $H^1(K^{-1})$ (use the exact sequence $0 \longrightarrow K^{-1} \longrightarrow \pi_* \mathcal{O}_{\tilde{C}}^\times \longrightarrow \mathcal{O}_C^\times \longrightarrow 0$), and the R -twist produces the required square roots of K . (For more details on the nilpotent locus, cf. [L] and [DEL].)

9.3.4 Adding values and representations

Fix a vector bundle K , and consider the moduli space $\mathcal{M}_{S,G,K}$ of K -valued G -principal Higgs bundles on S . (It can be constructed as in Simpson's [Sim1], even though the objects we need to parametrize are slightly different than his. In this subsection we sketch a direct construction.) It comes with a Hitchin map:

$$h : \mathcal{M}_{S,G,K} \longrightarrow B_K \tag{59}$$

where $B := B_K$ parametrizes all possible Hitchin data. Theorem 9.10 gives a precise description of the fibers of this map, independent of the values bundle K . This leaves us with the relatively minor task of describing, for each K , the corresponding base, i.e. the closed subvariety B_s of B parametrizing *split* Hitchin data, or K -valued cameral covers. The point is that Higgs bundles satisfy a symmetry condition, which in Simpson's setup is

$$\varphi \wedge \varphi = 0,$$

and is built into our definition 9.5 through the assumption that \mathfrak{c} is a bundle of regular centralizers, hence is abelian. Since commuting operators have common eigenvectors, this is translated into a splitness condition on the Hitchin data, which we describe below. (When K is a line bundle, the condition is vacuous, $B_s = B$.) The upshot is:

Lemma 9.12 *The following data are equivalent:*

- (a) A K -valued cameral cover of S .
- (b) A split, graded homomorphism $R^\bullet \longrightarrow \text{Sym}^\bullet K$.
- (c) A split Hitchin datum $b \in B_s$.

Here R^\bullet is the graded ring of W -invariant polynomials on \mathfrak{t} :

$$R^\bullet := (\text{Sym}^\bullet \mathfrak{t}^*)^W \approx \mathbf{C}[\sigma_1, \dots, \sigma_l], \quad \deg(\sigma_i) = d_i \tag{60}$$

where $l := \text{Rank}(\mathfrak{g})$ and the σ_i form a basis for the W -invariant polynomials. The Hitchin base is the vector space

$$B := B_K := \bigoplus_{i=1}^l H^0(S, \text{Sym}^{d_i} K) \approx \text{Hom}(R^\bullet, \text{Sym}^\bullet K).$$

For each $\lambda \in \Lambda$ (or $\lambda \in \mathfrak{t}^*$, for that matter), the expression in an indeterminate x :

$$q_\lambda(x, t) := \prod_{w \in W} (x - w\lambda(t)), \quad t \in \mathfrak{t}, \tag{61}$$

is W -invariant (as a function of t), so it defines an element $q_\lambda(x) \in R^\bullet[x]$. A Hitchin datum $b \in B \approx \text{Hom}(R^\bullet, \text{Sym}^\bullet K)$ sends this to

$$q_{\lambda,b}(x) \in \text{Sym}^\bullet(K)[x].$$

We say that b is *split* if, at each point of S and for each λ , the polynomial $q_{\lambda,b}(x)$ factors completely, into terms linear in x .

We note that, for λ in the interior of C (the positive Weyl chamber), $q_{\lambda,b}$ gives the equation in \mathbb{K} of the spectral cover \tilde{S}_λ of section (9.2.2): $q_{\lambda,b}$ gives a morphism $\mathbb{K} \longrightarrow$

$\text{Sym}^N \mathbb{K}$, where $N := \#W$, and \tilde{S}_λ is the inverse image of the zero-section. (When λ is in a face F_P of $\overline{\mathcal{C}}$, we define analogous polynomials $q_\lambda^P(x, t)$ and $q_{\lambda, b}^P(x)$ by taking the product in (61) to be over $w \in W_P \setminus W$. These give the reduced equations in this case, and q_λ is an appropriate power.)

Over B_s there is a universal K -valued cameral cover

$$\tilde{\mathcal{S}} \longrightarrow B_s$$

with ramification divisor $R \subset \tilde{\mathcal{S}}$. From the relative Picard,

$$\text{Pic}(\tilde{\mathcal{S}}/B_s)$$

we concoct the relative N -shifted, R -twisted Prym

$$\text{Prym}_{\Lambda, R}(\tilde{\mathcal{S}}/B_s).$$

By Theorem (9.10), this can then be considered as a parameter space $\mathcal{M}_{S, G, K}$ for all K -valued G -principal Higgs bundles on S . (Recall that our objects are assumed to be everywhere *regular!*) It comes with a ‘Hitchin map’, namely the projection to B_s , and the fibers corresponding to smooth projective \tilde{S} are abelian varieties. When S is a smooth, projective curve, we recover this way the algebraic complete integrability of Hitchin’s system and its generalizations. More generally, for any S , one obtains an ACIHS (with symplectic, respectively Poisson structures) when the values bundle has the same (symplectic, respectively Poisson) structure, by a slight modification of the construction in Chapter 8. One considers only Lagrangian supports which retain a W -action, and only equivariant sheaves on them (with the numerical invariants of a line bundle). These two restrictions are symplectically dual, so the moduli space of Lagrangian sheaves with these invariance properties is a symplectic (respectively, Poisson) subsp! ace of the total moduli space, and the fibers of the Hitchin map are Lagrangian as expected.

9.3.5 Irregulars?

The Higgs bundles we consider in this survey are assumed to be everywhere regular. This is a reasonable assumption for line-bundle valued Higgs bundles on a curve or surface, but *not* in $\dim \geq 3$. This is because the complement of \mathfrak{g}_{reg} has codimension 3 in \mathfrak{g} . The source of the difficulty is that the analogue of (52) fails over \mathfrak{g} . There are two candidates for the universal cameral cover: $\tilde{\mathfrak{g}}$, defined by the left hand side of (52), is finite over \mathfrak{g} with W action, but does not have a family of line bundles parametrized by Λ . These live instead on $\tilde{\tilde{\mathfrak{g}}}$, the object defined by the right hand side, which parametrizes pairs (x, \mathfrak{b}) , $x \in \mathfrak{b} \subset \mathfrak{g}$. This suggests that the right way to analyze irregular Higgs bundles may involve spectral data consisting of a tower

$$\tilde{\tilde{\mathcal{S}}} \xrightarrow{\sigma} \tilde{\mathcal{S}} \longrightarrow S$$

together with a homomorphism $\mathcal{L} : \Lambda \longrightarrow \text{Pic}(\tilde{\tilde{\mathcal{S}}})$ such that the collection of sheaves

$$\sigma_*(\mathcal{L}(\lambda)), \quad \lambda \in \Lambda$$

on \tilde{S} is R -twisted W -equivariant in an appropriate sense. As a first step, one may wish to understand the direct images $R^i \sigma_*(\mathcal{L}(\lambda))$ and in particular the cohomologies $H^i(F, \mathcal{L}(\lambda))$ where F , usually called a *Springer fiber*, is a fiber of σ . For regular x , this fiber is a single point. For $x = 0$, the fiber is all of G/B , so the fiber cohomology is given by the Borel-Weil-Bott theorem. The question may thus be considered as a desired extension of BWB to general Springer fibers.

References

- [AB] Adams, M. R., Bergvelt, M. J.: *The Krichever map, vector bundles over algebraic curves, and Heisenberg algebras*, Comm. Math. Phys. 154 (1993), 265-305.
- [AdC] Arbarello, E., De Concini, C.: *Another proof of a conjecture of S. P. Novikov on periods and abelian integrals on Riemann surfaces*, Duke Math. J. 54 (1987) 163-178.
- [AG] Arnol'd, V. I., Givental', A. B.: *Symplectic Geometry*. In: Arnol'd, V.I., Novikov, S.P. (eds.) *Dynamical systems IV*, (EMS, vol. 4, pp. 1-136) Berlin Heidelberg New York: Springer 1988
- [AHH] Adams, M. R., Harnad, J., Hurtubise J.: *Isospectral Hamiltonian flows in finite and infinite dimensions, II. Integration of flows*. Commun. Math. Phys. 134, 555-585 (1990)
- [AIK] Altman, A., Iarrobino, A., Kleiman, S.: *Irreducibility of the Compactified Jacobian*. Proc. Nordic Summer School, 1-12 (1976)
- [AK1] Altman, A. B., Kleiman S. L.: *Foundation of the theory of Fano schemes*, Comp. Math. Vol 34, Fasc. 1, 1977, pp 3-47
- [AK2] Altman, A. B., Kleiman S. L.: *Compactifying the Picard Scheme*, Advances in Math. 35, 50-112 (1980)
- [At] Atiyah, M.: *Vector bundles over an elliptic curve*. Proc. Lond. Math. Soc. 7, 414-452 (1957)
- [AvM] Adler, M., van Moerbeke, P.: *Completely integrable systems, Euclidean Lie algebras, and curves*, Advances in Math. 38, (1980), 267-379.
- [B1] Beauville, A.: Beauville, A.: *Jacobiennes des courbes spectrales et systèmes hamiltoniens complètement intégrables*. Acta Math. 164, 211-235 (1990)
- [B2] Beauville, A.: *Variétés Kähleriennes dont la première classe de Chern est nulle*. J. Diff. Geom. 18, p. 755-782 (1983).
- [Bi] Biswas, I.: *A remark on a deformation theory of Green and Lazarsfeld*, J. reine angew. Math. 449 (1994) 103-124.
- [Bo] Bogomolov, F.: *Hamiltonian Kähler manifolds*, Soviet Math Dokl. **19** (1978), 1462-1465.
- [Bn] Bottacin, F.: Thesis, Orsay, 1992.
- [BD] Beauville, A., Donagi, R.: *La variété des droites d'une hypersurface cubique de dimension 4*. C. R. Acad. Sci. Paris Ser. I t. 301, 703-706 (1985)
- [BG] Bryant, R., Griffiths, P.: *Some observations on the infinitesimal period relations for regular threefolds with trivial canonical bundle*, in Arithmetic and Geometry, Papers dedicated to I.R. Shafarevich, vol. II, Birkhäuser, Boston (1983), 77-102.

- [BK] Beilinson, A., Kazhdan, D.: *Flat Projective Connections*, unpublished(1990).
- [BL] Beauville, A., Laszlo, Y.: *Conformal blocks and generalized theta functions*, Comm. Math. Phys., to appear.
- [BNR] Beauville, A., Narasimhan, M. S., Ramanan, S.: *Spectral curves and the generalized theta divisor*. J. Reine Angew. Math. 398, 169-179 (1989)
- [CdOGP] Candelas, P., de la Ossa, X. C., Green, P. S., and Parkes L.: *A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory*, Phys. Lett. **B 258** (1991), 118-126; Nuclear Phys. **B 359** (1991), 21-74.
- [CG] Clemens, H., Griffiths, P.: *The intermediate Jacobian of the cubic threefold*. Annals of Math. 95 281-356 (1972)
- [De1] Deligne P.: *Theoreme de Lefschetz et criteres de degenerescence de suites spectrales*, Publ. Math. IHES, 35(1968) 107-126.
- [De2] Deligne, P.: letter to C. Simpson, March 20, 1989
- [D1] Donagi, R.: *The tetragonal construction*, Bull. Amer. Math. Soc. (N.S.) **4** (1981), 181-185.
- [D2] Donagi, R.: *Decomposition of spectral covers*, in: Journees de Geometrie Algebrique D'Orsay, Asterisque 218 (1993), 145-175.
- [D3] Donagi, R.: *Abelianization of Higgs bundles*, preprint.
- [D4] Donagi, R.: *Spectral covers* , To appear in: Proceedings of special year in algebraic geometry 1992–1993, MSRI. (Alg. Geom. eprint no. 9505009).
- [D5] Donagi, R.: *Group law on the intersection of two quadrics*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **7** (1980), no. 2, 217-239.
- [DEL] Donagi, R., Ein, L. and Lazarsfeld, R.: *A non-linear deformation of the Hitchin dynamical system*, preprint, (Alg. Geom. eprint no. 9504017).
- [DM] Donagi, R. and Markman, E.: *Cubics, integrable systems, and Calabi-Yau threefolds*, to appear in the proceedings of the Algebraic Geometry Workshop on the occasion of the 65th birthday of F. Hirzebruch, May 1993. (Alg. Geom. eprint no. 9408004).
- [DR] Desale, U. V., Ramanan, S.: *Classification of Vector Bundles of Rank 2 on Hyperelliptic Curves*. Invent. Math. 38, 161-185 (1976)
- [EZ] El Zein F. and Zucker S.: *Extendability of normal functions associated to algebraic cycles*. In *Topics in transcendental Algebraic Geometry*, ed. by P. Griffiths, Annals of Mathematics
- [F1] Faltings, G.: *Stable G-bundles and Projective Connections*, Jour. Alg. Geo **2** (1993), 507-568.

- [F2] Faltings, G.: *A proof of the Verlinde formula*, to appear in Jour. Alg. Geo.
- [Gr] Griffiths, P. A.: *Infinitesimal variations of Hodge structure III: determinantal varieties and the infinitesimal invariant of normal functions*. Compositio Math. **50** (1983), 267-324.
- [GH] Griffiths, P. A., Harris, J.: *Principles of algebraic geometry*. Pure and Applied Mathematics. Wiley-Interscience 1978
- [Gro] Grothendieck, A.: *Techniques de construction et théorèmes d'existence en géométrie algébrique, IV : Les schémas de Hilbert*. Séminaire Bourbaki 13e année (1960/61) Exposés 221
- [HS] Hilton P., Stammach U.: *A course in homological algebra*, Graduate Texts in Mathematics, Vol. 4, Springer-Verlag, New York-Berlin,
- [H1] Hitchin, N.J.: *The self-duality equations on a Riemann surface*, Proc. Lond. Math. Soc. **55** (1987) 59–126.
- [H2] Hitchin, N. J.: *Stable bundles and integrable systems*. Duke Math. J. 54, No 1 91-114 (1987)
- [H3] Hitchin, N. J.: *Flat Connections and Geometric Quantization*. Comm. Math. Phys. 131, 347-380 (1990)
- [Hur] Hurtubise J.: *Integrable systems and Algebraic Surfaces*, Preprint 1994
- [IVHS] Carlson, J., Green, M., Griffiths, P., and Harris, J.: *Infinitesimal variations of Hodge structure (I)*. Compositio Math. **50** (1983), 109-205.
- [K] Kanev, V.: *Spectral curves, simple Lie algebras and Prym-Tjurin varieties*, Proc. Symp. Pure Math. **49** (1989), Part I, 627-645.
- [Ka] Kawamata, Y.: *Unobstructed deformations - a remark on a paper of Z. Ran* J. Alg. Geom. 1 (1992) 183-190
- [Kn] Knörrer, H.: *Geodesics on the Ellipsoid*. Invent. Math. 59, 119-143 (1980)
- [Ko] Kobayashi, S.: *Simple vector bundles over symplectic Kähler manifolds*. Proc. Japan Acad. Ser. A Math. Sci. 62, 21-24 (1986)
- [Ko-P] Kouvidakis, A., Pantev, T.: *automorphisms of the moduli spaces of stable bundles on a curve.*, Math. Ann., to appear.
- [KP1] Katzarkov, L., Pantev, T.: *Stable G_2 bundles and algebraically completely integrable systems*, Comp. Math. **92** (1994), 43-60.
- [KP2] Katzarkov, L., Pantev, T.: *Representations of Fundamental groups whose Higgs bundles are pullbacks*, J. Diff. Geo. 39 (1994), 103-121.
- [L] Laumon, G.: *Un analogue global du cône nilpotent*, Duke Math. J. **57**(1988), 647-671.

- [LM1] Li, Y., Mulase, M.: *Category of morphisms of algebraic curves and a characterization of Prym varieties*. Preprint
- [LM2] Li, Y., Mulase, M.: *Hitchin systems and KP equations*. Preprint
- [Ma1] Markman, E.: *Spectral curves and integrable systems*, Compositio Math. 93, 255-290, (1994).
- [Ma2] Markman, E.: In preparation.
- [Me] Merindol, J. Y.: *Varietes de Prym d'un Revetement Galoisien*, prep.(1993).
- [Mor] Morrison, D.: *Mirror Symmetry and Rational Curves on Quintic Threefolds: A Guide for Mathematicians*, J. Am. Math. Soc. **6** (1993), 223.
- [Mo] Moser, J.: *Various aspects of integrable Hamiltonian systems*, in: Dynamical Systems 1978, Prog. Math. Vol. 8, 233-290.
- [MS] McDaniel, A., Smolinsky, L.: *A Lie theoretic Galois Theory for the spectral curves of an integrable system II*, prep. (1994).
- [Mu1] Mukai S.: *Symplectic structure of the moduli space of sheaves on an abelian or K3 surface*, Invent. math. 77,101-116 (1984)
- [Mu2] Mukai S.: *On the moduli space of bundles on K3 surfaces I*, Vector bundles on algebraic varieties, Proc. Bombay Conference, 1984, Tata Institute of Fundamental Research Studies, no. 11, Oxford University Press, 1987, pp. 341-413.
- [Mul] Mulase, M.: *Cohomological structure in soliton equations and Jacobian varieties*, J. Diff. Geom. 19 (1984), 403-430.
- [Mum1] Mumford, D.: *Lectures on curves on an algebraic surface*, Princeton University Press, 1966.
- [Mum2] Mumford, D.: *Tata Lectures on Theta II*, Birkhaeuser-Verlag, Basel, Switzerland, and Cambridge, MA 1984
- [Mum3] Mumford, D.: *Prym varieties I*, in:Contribution to analysis, Acad. Press (1974), 325-350.
- [Nit] Nitsure, N.: *Moduli space of semistable pairs on a curve*. Proc. London Math. Soc. (3) 62 (1991) 275-300
- [NR] Narasimhan, M. S., Ramanan, M. S.: *2θ -linear systems on abelian varieties*, in Vector bundles and algebraic varieties, Oxford University Press, 415-427 (1987)
- [Ra] Ran, Ziv: *Lifting of cohomology and unobstructedness of certain holomorphic maps*. Bull. Amer. Math. Soc. (N.S.) 26 (1992), no. 1, 113-117.
- [Re] Reid, M.: *The complete intersection of two or more quadrics*, Thesis, Cambridge (GB) 1972.

- [Rec] Recillas, S.: *Jacobians of curves with g_4^1 's are Pryms of trigonal curves*, Bol. Soc. Mat. Mexicana **19** (1974) no.1.
- [Se] Seshadri, C. S.: *Fibrés vectoriels sur les courbes algébriques*. asterisque 96 1982
- [Sc] Scognamillo, R.: *Prym-Tjurin Varieties and the Hitchin Map*, preprint (1993).
- [Sch] Schmid, W.: *Variation of Hodge structure: the singularities of the period mapping*. Invent. Math. 22 (1973), 211-319.
- [Sh] Shiota, T.: *Characterization of Jacobian varieties in terms of soliton equations*, Inv. Math. 83 (1986) 333-382.
- [Sim1] Simpson, C.: *Moduli of representations of the fundamental group of a smooth projective variety I, II*. Preprint
- [Sim2] Simpson, C.: *Higgs bundles and local systems*. Inst. Hautes Etudes Sci. Publ. Math. (1992), No. 75 5-95.
- [Sim3] Simpson, C.: *Nonabelian Hodge Theory*, Proceedings of the International Congress of Mathematics, Kyoto, Japan, 1990.
- [SW] Segal, G., Wilson, G.: *Loop groups and equations of KdV type*. Inst. Hautes Etudes Sci. Publ. Math. (1985), No. 61 5-65.
- [Ti] Tian, G.: *Smoothness of the universal deformation space of compact Calabi-Yau manifolds and its Petersson-Weil metric*, in Mathematical Aspects of String Theory (S.T. Yau, ed.), World Scientific, Singapore (1987), 629-646.
- [To] Todorov, A.: *The Weil-Petersson geometry of the moduli space of $SU(n \geq 3)$ (Calabi-Yau) manifolds, I*, Comm. Math. Phys. **126** (1989), 325-346.
- [TV1] Treibich, A., J.-L. Verdier *Solitons elliptiques*. The Grothendieck Festschrift, Vol. III, 437-480, Birkhauser Boston, 1990.
- [TV2] Treibich, A., J.-L. Verdier *Varietes de Krichever des solitons elliptiques de KP*. Proceedings of the Indo-French Conference on Geometry (Bombay, 1989), 187-232, 1993.
- [Ty1] Tyurin, A. N.: *Symplectic structures on the varieties of moduli of vector bundles on algebraic surfaces with $p_g > 0$* . Math. USSR Izvestiya Vol. 33(1989), No. 1
- [Ty2] Tyurin, A. N.: *The geometry of moduli of vector bundles*. Russian Math. Surveys 29:6 57-88 (1974)
- [V] Voisin, C.: *Sur la stabilite des sous-varietes lagrangiennes des varietes symplectiques holomorphes*. Complex projective geometry (Trieste, 1989/Bergen, 1989), 294-303, Cambridge Univ. Press, Cambridge, 1992.
- [We] Weinstein, A.: *The Local Structure of Poisson Manifolds*. J. Diff. Geom. 18, 523-557 (1983)

- [Ye] Ye, Yun-Gang: *Lagrangian subvarieties of the moduli space of stable vector bundles on a regular algebraic surface with $p_g > 0$* . Math. Ann. 295, 411-425 (1993)